

W , Z and exclusive $\mu\mu$ at LHCb

Ronan McNulty
University College Dublin




Workshop on Heavy Particles at the LHC
Pauli Centre, ETH, Zurich
05.01.2011

Outline

- Overview
- W, Z production ($x \sim 10^{-4}$)
- Sensitivity to PDFs
- Sensitivity to $\sin^2\theta_W$
- γ^* production ($x \sim 10^{-6}$)
- Exclusive processes


EW physics motivation





Test the SM at highest energies


Check out that DGLAP evolution works (test QCD)


Push theory into interesting regions with very soft gluons















LHCb: a forward spectrometer








Effect of PDF uncertainties on cross-sections

- Region where the most precise EW tests can be made.
- At highest rapidities, PDFs can be constrained.

- Experimental statistical error <<1%.
- Systematic error likely to be $\sim 1\%$



But you can do better !

$$R_{\mp}(y_W) \equiv \frac{d\sigma/dy_W(W^-)}{d\sigma/dy_W(W^+)} \approx \frac{d(x_1)\bar{u}(x_2)}{u(x_1)\bar{d}(x_2)} = \frac{d(x_1)}{u(x_1)} \cdot \frac{\bar{u}(x_2)}{\bar{d}(x_2)} \approx \frac{d(x_1)}{u(x_1)}$$

So ratio of Ws is sensitive to d to u ratio. (For LHCb d_v/u_v)

$$A_{\pm}(y_W) = \frac{d\sigma/dy_W(W^+) - d\sigma/dy_W(W^-)}{d\sigma/dy_W(W^+) + d\sigma/dy_W(W^-)} \approx \frac{u(x_1)\bar{d}(x_2) - d(x_1)\bar{u}(x_2)}{u(x_1)\bar{d}(x_2) + d(x_1)\bar{u}(x_2)} \approx \frac{u(x_1) - d(x_1)}{u(x_1) + d(x_1)}$$

W asymmetry is sensitive to difference in u and d. (For LHCb u_v-d_v)

$$R_{Z/W}(y) \equiv \frac{d\sigma/dy(Z^0)}{d\sigma/dy(W^+) + d\sigma/dy(W^-)} \approx \frac{\kappa_u u(x'_1)\bar{u}(x'_2) + \kappa_d d(x'_1)\bar{d}(x'_2)}{|V_{ud}|^2 \{u(x_1)\bar{d}(x_2) + d(x_1)\bar{u}(x_2)\}}$$


Ratio of Z to W is almost insensitive to PDFs!

Gold plated test of SM at the highest energies

Effect of PDF uncertainties on cross-sections



- R_{WZ} precise test of SM everywhere.
- Difference in u and d quarks can be significantly improved by all experiments at the LHC.
- Going forward, you increasingly constrain the u-valence to d-valence ratio.

- Even nicer, most experimental systematics (especially luminosity) cancel in the ratio.



Z Cross-section Measurement at LHCb

$\sigma(Z \rightarrow \mu\mu : 2 < \eta_\mu < 4.5, P_{T\mu} > 20 \text{ GeV}, 81 < M_{\mu\mu} < 101 \text{ GeV})$





**8% of Z within
LHCb acceptance**

**17% (16%) of W⁺ (W⁻)
within LHCb acceptance**



Z: Characteristic signature of two high transverse momentum muons with invariant mass m_Z

Require two muons: $p_t > 20\text{ GeV}$, $2 < \eta < 4.5$. $81 < M_{\mu\mu} < 101\text{ GeV}$



Z: Characteristic signature of two high transverse momentum muons with invariant mass m_Z

Require two muons: $p_t > 20 \text{ GeV}$, $2 < \eta < 4.5$. $81 < M_{\mu\mu} < 101 \text{ GeV}$



To turn into cross-section:


$$\sigma_{Z \rightarrow \mu\mu}(\Delta y) = \frac{N_{tot}^Z - N_{bkg}^Z}{\epsilon_Z L}$$

$$\epsilon_Z = A_z \epsilon_Z^{trig} \epsilon_Z^{track} \epsilon_Z^{muon} \epsilon_Z^{selection} = 1 = 1$$


Insomuch as possible, take everything from data.

Muon identification efficiency

$$\epsilon_Z = A_z \epsilon_Z^{trig} \epsilon_Z^{track} \epsilon_Z^{muon} \epsilon_Z^{selection}$$




Efficiency flat in p_T, η, ϕ
No evidence for charge bias





Track Efficiency from Tag and Probe

$$\epsilon_Z = A_z \epsilon_Z^{\text{trig}} \epsilon_Z^{\text{track}} \epsilon_Z^{\text{muon}} \epsilon_Z^{\text{selection}}$$



$\epsilon_{W+} = 73 \pm 3\%$
$\epsilon_{W-} = 78 \pm 3\%$
$\epsilon_Z = 83 \pm 3\%$

(W+/W- differ due to lepton η distribution)




(flat in p_T and ϕ)

η


Trigger Efficiency from Tag and Probe

$$\epsilon_Z = A \cdot \epsilon_Z^{trig} \epsilon_Z^{track} \epsilon_Z^{muon} \epsilon_Z^{selection}$$



$$\begin{aligned}\epsilon_W &= 0.9 * (80.5 \pm 1.4)\% \\ \epsilon_Z &= 0.9 * (95.7 \pm 0.5)\%\end{aligned}$$

(Global cuts:
Very large events
rejected by trigger)




Z analysis

$$\sigma_{Z \rightarrow \mu\mu}(\Delta y) = \frac{N_{tot}^Z - N_{bkg}^Z}{\epsilon_Z L}$$

N_Z^{tot}	833
$Z \rightarrow \tau\tau$	0.2 ± 0.2
Heavy flavours	1 ± 1
Misidentified π/K	$<< 1$
N_Z^{bkg}	1.2 ± 1.2
ϵ_{trig}^Z	0.86 ± 0.01
ϵ_{track}^Z	0.83 ± 0.03
ϵ_{muon}^Z	0.97 ± 0.01
ϵ_{sel}^Z	1.
A^Z	1.
ϵ_Z	0.69 ± 0.03
L	$16.5 \pm 1.7 pb^{-1}$
$\sigma_Z(2. < \eta_1, \eta_2 < 4.5, 81 < m_Z < 101)$	$73 \pm 4 \pm 7 pb.$

↑
Phase space for measurement ↑



Differential distribution (Z)



Measuring A_{FB} in $pp \rightarrow Z \rightarrow \mu\mu$?

$$A_{FB}^0 = \frac{3}{4} A_\mu (u A_u + d A_d + s A_s)$$

- LHC Problems: 1. PDF uncertainties
2. which is quark direction?



Statistically, a forward detector at high luminosity could measure A_{FB} with better precision than current WA.



How confident are we of theory? of PDF uncertainties? of detector systematics?

W Cross-section Measurement at LHCb

$$\sigma(W \rightarrow \mu\nu : 2 < \eta_\mu < 4.5, P_{T\mu} > 20 \text{ GeV})$$

W/One high transverse momentum muon with little else. Background much larger than for Z

Require muon $p_t > 20\text{GeV}$ + little activity in 'rest of event'




Can be found
using Z events
in data


Selecting W events

- pt of muon (>20GeV)
- ip significance of muon (<2)
- Mass of *rest of event* (<20 GeV)
- Pt of *rest of event* (<10 GeV)
- Charged transverse momentum in cone of 0.5 units of $\sqrt{(\Delta\eta)^2 + (\Delta\Phi)^2}$ around muon. (<2 GeV)

W analysis



W analysis



$$\sigma_{W \rightarrow \mu\nu}(\Delta\eta) = \frac{N_{tot}^W - N_{bkg}^W}{\epsilon_W L} \rightarrow \frac{N_{tot}^W}{L} \left(\frac{p_W}{\epsilon_W} \right) \quad \text{and all found from data.}$$


W analysis

$$\sigma_{W \rightarrow \mu\nu}(\Delta\eta) = \frac{N_{tot}^W - N_{bkg}^W}{\epsilon_W L}$$

	$W+$	$W-$
N_W^{tot}	7624	5732
$W \rightarrow \tau\nu$	151	90
$Z \rightarrow \tau\tau$	2	2
$Z \rightarrow \mu\mu$	460	506
QCD	2194 ± 150	1654 ± 150
N_W	4817 ± 165	3480 ± 161
ϵ_{trig}^W	0.725 ± 0.03	
ϵ_{track}^W	0.73 ± 0.03	0.78 ± 0.03
ϵ_{muon}^W	0.982 ± 0.005	
ϵ_{sel}^W	0.55 ± 0.01	
A^W	1	1
ϵ_W	0.29 ± 0.01	0.31 ± 0.01
N_W^{tot}	16610 ± 800	11226 ± 650
L	$16.5 \pm 1.7 \text{ pb}^{-1}$	$16.5 \pm 1.7 \text{ pb}^{-1}$
$\sigma_W(2.0 < y < 4.5)$	$1007 \pm 48 \pm 100 \text{ pb}$	$682 \pm 40 \pm 68 \text{ pb}$

(Measurement as function of lepton rapidity)

W asymmetry




W,Z Summary

$$\frac{\sigma_{W+}(2.0 < \eta_\mu < 4.5) + \sigma_{W-}(2.0 < \eta_\mu < 4.5)}{\sigma_Z(2.0 < y < 4.5)}$$

Generator	Order	PDF Set	Z	W+	W-	$(W^+ + W^-)/Z$	W^+/W^-
FEWZ	NLO	MSTW08NLO CTEQ66NLO NNPDF2.0	$65.7^{+2.9}_{-2.5}$				
MCFM		MSTW08NLO CTEQ66NLO NNPDF2.0	$65.5^{+2.8}_{-2.5}$	855 ± 43	656 ± 39	23.1 ± 0.1	1.30 ± 0.05
FEWZ	NNLO	MSTW08NNLO					
Data			$73 \pm 4 \pm 7.5$	$1007 \pm 48 \pm 101$	$682 \pm 40 \pm 68$	23.1 ± 1.5	1.48 ± 0.11


LHCb preliminary

LHCb Preliminary using 16.5pb^{-1} of data.
 Theory: FEWZ at NLO for Z; MCFM at NLO for W.
 Kinematic cuts: charged leptons $p_T > 20\text{ GeV}$, $2 < \eta < 4.5$.
 Uncertainty band combines NLO and MSTW2008 90% uncertainties.




How well can W,Z measurements constrain the PDFs?

Improvement to **MSTW08 PDFs** with 0.1fb^{-1} of high mass vector bosons at 7TeV




Modest improvement with small amount of data

Comparison with different PDFs using 0.1fb-1 of high mass vector bosons at 7TeV



Similar sensitivity. Ability to distinguish models

Improvement to CTEQ66 PDFs with 1fb-1 of high mass vector bosons at 14 TeV





CTEQ66 1 fb⁻¹

- reduced with Z
- reduced with W^+
- reduced with W^-
- reduced with WWZ

More data and higher energy lead to larger improvements.

Using γ^* to go to very low-x.






Unexplored region
Very little HERA
data at this x

More theoretical
work required to
make predictions


Sensitivity to
saturation?

Effect of PDF uncertainties on cross-sections



PDF uncertainties at low- x , low- Q^2

MSTW08. (Thanks to Graeme Watt)




Different behaviour and uncertainty with order of calculation.


Gluon essentially unconstrained by data below 10^{-4}

DGLAP evolution not trustworthy in this region. Gluon re-summarization effects. Possibly entering saturation regime.

LHCb preliminary




Improvement to MSTW08 PDFs with 0.1 fb⁻¹ of low mass vector bosons at 7 TeV



Similar improvements to NNPDF, CTEQ and Alekhin PDFs.

Sensitivity exists to distinguish between models.

Current uncertainty on MSTW08 PDFs and projections with 0.1fb^{-1} , 1fb^{-1} of very low invariant mass muons at 7TeV



— Significant improvements possible with modest amount of data

Exclusive dimuon final states


Exclusive particle production (2μ and nothing else)

1

Exclusive JPsis, Psi' ($\rightarrow \mu^+\mu^-$)

Produced by photon pomeron fusion

Starlight: Models diphoton and photon pomeron fusion
 (S.R.Klein and J.Nystrand, Phys. Rev. Lett. 92 (2004) 142003).




2

Exclusive ChiC ($\rightarrow \mu^+\mu^- + \gamma$)

Produced by double pomeron exchange

SuperChiC: MC for central exclusive production
 (L.A. Harland-Lang, V.A. Khoze, M.G. Ryskin, W.J. Stirling,
 arXiv:0909.4748 [hep-ph].).




3

Exclusive diphoton dimuon


Produced by diphoton fusion

LPAIR: Models EM production of lepton pairs
 (A.G.Shamov and V.I.Telnov, NIM A {\bf 494} (2002) 51).

For LHC
Luminosity




But is it exclusive ?



#J/Psi as Fn of #tracks


Assuming Linear background ?

Extrapolate Background to determine number of experimentally exclusive events



817 Exclusive JPsi candidates
77 Background Events

Purity of Events = 91%




76 Exclusive ChiC candidates
22 Background Events

Purity of Events = 78%

Exclusive J/Psi (compared to Starlight)

Require 2μ consistent with Jpsi + no other charged or neutral activity




Number of exclusive candidates: 817 (Calculated from extrapolation)

Purity of Events = 91%

2

Exclusive Chi_c (compared to SuperCHIC)


Require 2m consistent with Jpsi, 1 photon, and nothing else



3

Diphoton fusion (compared to LPAIR)

Require 2 back-to-back muons and no other charged or neutral activity.
Remove resonances.



As in other cases, feed-down from non-exclusive processes needs to be evaluated.

Precision on luminosity given by uncertainty on backgrounds and trigger efficiency.

Summary

- LHCb EW programme complementary to ATLAS/CMS
- Tests SM in different region with similar precision
- Possible future precision measurement of $\sin^2\theta_w$
- Constrain PDFs at low x for W, Z and very low x for γ^* .
- Potential for luminosity measurement at few % level.