

J/ψ Production at NLO in NRQCD

Bernd Kniehl

II. Institut für Theoretische Physik
Universität Hamburg

LHC Physics Day: Charm and bottom quark production at the
LHC
CERN, 3 December 2010



In collaboration with Mathias Butenschön
Phys. Rev. Lett. **104** (2010) 072001, arXiv:1009.5662 [hep-ph]

Production and Decay Rates of Heavy Quarkonia

Heavy quarkonia: Bound states of heavy quark and its antiquark.

- Charmonia ($c\bar{c}$) and Bottomonia ($b\bar{b}$)
 - Top decays too fast for bound state

The classic approach: Color-singlet model

- Calculate cross section for heavy quark pair in physical color-singlet (= color neutral) state. In case of J/ψ : $c\bar{c}[{}^3S_1]$ ^[1]
 - Multiply by quarkonium wave function (or its derivative) at origin
 - Strong disagreement with Tevatron data

Nonrelativistic QCD (NRQCD):

- 1995: Rigorous effective field theory by Bodwin, Braaten, Lepage
 - Based on **factorization of soft and hard scales**
(Scale hierarchy: $Mv^2, Mv \ll \Lambda_{\text{QCD}} \ll M$)
 - Could explain hadroproduction at Tevatron

J/ψ Production with NRQCD

Factorization theorem: $\sigma_{J/\psi} = \sum_n \sigma_{c\bar{c}[n]} \cdot \langle O^{J/\psi}[n] \rangle$

- n : Every possible Fock state, including color-octet states.
 - $\sigma_{c\bar{c}[n]}$: Production rate of $c\bar{c}[n]$, calculated in perturbative QCD.
 - $\langle O^{J/\psi}[n] \rangle$: Long distance matrix elements (LDMEs):
describe $c\bar{c}[n] \rightarrow J/\psi$, universal, extracted from experiment.


Scaling rules: MEs scale with relative velocity v ($v^2 \approx 0.2$):

scaling	v^3	v^7	v^{11}
n	${}^3S_1^{[1]}$	${}^1S_0^{[8]}, {}^3S_1^{[8]}, {}^3P_{0/1/2}^{[8]}$	\dots

- Double expansion in v and α_s .
 - Leading term in v ($n = {}^3S_1^{[1]}$) equals color-singlet model.


Production of J/ψ : NRQCD vs. Experiment (History)

Hadroproduction at Tevatron:



- Color octet states important
⇒ Great success for NRQCD

Photoproduction at HERA:



- MEs from fits to Tevatron data.
 - Importance of color octet **unclear**

This work: NLO NRQCD calculation for photo- and hadroproduction
⇒ Aim: Establish universality of long distance matrix elements.

Production of J/ψ : Summary of Calculations

Hadroproduction:

	$^3S_1^{[1]}$	$^1S_0^{[8]}, ^3S_1^{[8]}, ^3P_{0/1/2}^{[8]}$
Born	Baier, Rückl (1980)	Cacciari, Krämer (1996)
NLO	Campbell et al. (2007)	Butenschön, BK (2010) Ma et al. (2010)

Photoproduction:


	$^3S_1^{[1]}$	$^1S_0^{[8]}, ^3S_1^{[8]}, ^3P_{0/1/2}^{[8]}$
Born	Berger, Jones (1981)	Ko, Lee, Song (1996)
NLO	Krämer (1995)	Butenschön, BK (2009)

Open question of ME universality:

- NLO NRQCD calculation: after **14 years!**
- Difficulty: virtual corrections to **P states**

Direct J/ψ Production

Factorization formulas: (e.g. photoproduction)



- Convolute partonic cross sections with **proton PDFs**:

$$\sigma_{\text{hadr}} = \sum_i \int dx f_{i/p}(x) \cdot \sigma_{\text{part},i}$$

- NRQCD factorization:**

$$\sigma_{\text{part},i} = \sum_n \sigma(\gamma i \rightarrow c\bar{c}[n] + X) \cdot \langle O^{J/\psi}[n] \rangle$$


Amplitudes for $c\bar{c}[n]$ production by projector application, e.g.:

$$A_{c\bar{c}[3S_1^{1/8}]} = \varepsilon_\alpha \text{Tr} [C \Pi^\alpha A_{c\bar{c}}] |_{q=0}$$

$$A_{c\bar{c}[3P_J^{[8]}]} = \varepsilon_{\alpha\beta} \frac{d}{dq_\beta} \text{Tr} [C \Pi^\alpha A_{c\bar{c}}] |_{q=0}$$


- $A_{c\bar{c}}$:** Amputated pQCD amplitude for open $c\bar{c}$ production.
- q :** Relative momentum between c and \bar{c} .

Overview of IR Singularity Structure



Structure of Soft Singularities

Soft limits of the real corrections:



S and P states: Soft #1 + Soft #2 + Soft #3 terms:

$$A_{\text{soft,s}} = A_{\text{soft}}(0) = A_{\text{Born,s}} \cdot E(0)$$


$$A_{\text{soft,p}} = A'_{\text{soft}}(0) = A_{\text{Born,p}} \cdot E(0) + A_{\text{Born,s}} \cdot E'(0)$$

$$|A_{\text{soft,s}}|^2 = |A_{\text{Born,s}}|^2 \cdot E(0)^2$$

$$|A_{\text{soft,p}}|^2 = |A_{\text{Born,p}}|^2 \cdot E(0)^2 + 2 \operatorname{Re} A_{\text{Born,s}}^* A_{\text{Born,p}} \cdot E(0) E'(0) + |A_{\text{Born,s}}|^2 \cdot E'(0)^2$$

Radiative Corrections to Long Distance MEs

In NRQCD: Long distance MEs = $c\bar{c}$ scattering amplitudes:



$O[n]$ = 4-fermion operators
($n = {}^3S_1^{[1]}, {}^1S_0^{[8]}, {}^3S_1^{[8]}, {}^3P_{0/1/2}^{[8]}, \dots$)



Corrections to $\langle O^{J/\psi}[{}^3S_1^{[1/8]}] \rangle$ with NRQCD Feynman rules:

+ similar diagrams $\propto \frac{4\alpha_s}{3\pi m_c^2} \left(\frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}} \right) \cdot$

- UV singularity cancelled by renormalization of 4-fermion operat.
- IR singularity cancels soft #3 terms of P states.

Combined Fit to Tevatron and HERA (1)



Fit CO LDMEs to p_T distributions from Tevatron II and HERA II:



- Best fit values: $\langle \mathcal{O}(^1S_0^{[8]}) \rangle = (0.0476 \pm 0.0071) \text{ GeV}^3$,
 $\langle \mathcal{O}(^3S_1^{[8]}) \rangle = (0.00265 \pm 0.00091) \text{ GeV}^3$, $\langle \mathcal{O}(^3P_0^{[8]}) \rangle = -(0.0132 \pm 0.0035) \text{ GeV}^5$
 - $\propto v^4 \langle \mathcal{O}_1(^3S_1) \rangle \rightsquigarrow \text{NRQCD velocity scaling rules } \checkmark$

Combined Fit to Tevatron and HERA (2)



Contribution of individual states:



- Hadroprod.: Short-distance $\sigma(c\bar{c}[{}^3P_J^{[8]}])$ negative for $p_T \gtrsim 7$ GeV
 - But: Short-distance cross sections and LDMEs unphysical
(NRQCD scale and scheme dependence) \implies No problem!

Further Predictions for HERA


Use LDMEs to make W and z distribution predictions:



- Proton rest frame: $z =$ fraction of photon energy going to J/ψ .
 - $z \lesssim 0.45$: **Resolved** photoproduction important (not yet included).
 - W distribution also well described. Data not part of the fit.

Predictions for RHIC



Use LDMEs to make prediction for RHIC:



- Also RHIC data well described by CS+CO.
- Like at Tevatron: CS orders of magnitudes below the data.
- These data not part of the fit, outcome not trivial.

Predictions for LHC



Use LDMEs to make predictions for CMS:



- Also CMS data **well described** by CS+CO.
- Like at Tevatron: **CS** orders of magnitudes **below** the data.
- These data **not part** of the fit, outcome not trivial.

Predictions for LHC (2)



Use LDMEs to make predictions for ATLAS:



- Also ATLAS data **well described** by CS+CO.
- Like at Tevatron: **CS** orders of magnitudes **below** the data.
- These data **not part** of the fit, outcome not trivial.

Predictions for LHC (3)



Use LDMEs to make predictions for ATLAS and LHCb:



- Also ATLAS and LHCb data **well described** by CS+CO.
- Like at Tevatron: **CS** orders of magnitudes **below** the data.
- These data **not part** of the fit, outcome not trivial.

Comparison with Ma, Wang, Chao arXiv:1009.3655


Fit only to Tevatron data with $p_T > 7$ GeV, including feed-down



- Observe that $d\hat{\sigma}({}^3P_J^{[8]}) \approx r_0 d\hat{\sigma}({}^1S_0^{[8]}) + r_1 d\hat{\sigma}({}^3S_1^{[8]})$ with $r_0 = 3.9$ and $r_1 = -0.56$
- Define $M_0 = \langle \mathcal{O}({}^1S_0^{[8]}) \rangle + \frac{r_0}{m_c^2} \langle \mathcal{O}({}^3P_0^{[8]}) \rangle$ and $M_1 = \langle \mathcal{O}({}^3S_1^{[8]}) \rangle + \frac{r_1}{m_c^2} \langle \mathcal{O}({}^3P_0^{[8]}) \rangle$
- Substitute $\langle \mathcal{O}({}^1S_0^{[8]}) \rangle \rightarrow M_0$ and $\langle \mathcal{O}({}^3S_1^{[8]}) \rangle \rightarrow M_1$ and discard $d\hat{\sigma}({}^3P_J^{[8]})$.
- Fit yields $M_0 = (7.4 \pm 1.9) \times 10^{-2}$ GeV 3 and $M_1 = (0.05 \pm 0.02) \times 10^{-2}$ GeV 3 with $\chi^2/\text{d.o.f.} = 0.33$
- Cf. $M_0 = (2.47 \pm 0.93) \times 10^{-2}$ GeV 3 and $M_1 = (0.59 \pm 0.13) \times 10^{-2}$ GeV 3 from BK

Significance of HERA data

Fit only to Tevatron data with $p_T > 3$ GeV, excluding feed-down




10^{-2} GeV $^{3+2L}$	BK default	Tevatron only
$\langle \mathcal{O}(^1S_0^{[8]}) \rangle$	4.76 ± 0.71	59
$\langle \mathcal{O}(^3S_1^{[8]}) \rangle$	0.265 ± 0.091	-6.6
$\langle \mathcal{O}(^3P_0^{[8]}) \rangle$	-1.32 ± 0.35	-29

- CO MEs too large in magnitude \rightsquigarrow NRQCD velocity scaling rules violated
- Substantial fine tuning \rightsquigarrow unnatural

Significance of p_T cut on Tevatron data

Fit to Tevatron $p_T > 7$ GeV and HERA data, excluding feed-down





	$10^{-2} \text{ GeV}^{3+2L}$	BK default	Tevatron $p_T > 7$ GeV
• $\langle \mathcal{O}(^1S_0^{[8]}) \rangle$	4.76 ± 0.71	2.9	
• $\langle \mathcal{O}(^3S_1^{[8]}) \rangle$	0.265 ± 0.091	0.4	
• $\langle \mathcal{O}(^3P_0^{[8]}) \rangle$	-1.32 ± 0.35	-0.58	

- Shift of p_T cut on Tevatron data from 3 GeV to 7 GeV less dramatic
- Thanks to stabilizing influence of HERA data

Significance of feed-down

Fit to HERA and Tevatron data w/ feed-down subtracted



- feed-down/prompt $\approx (32 - 0.62 \frac{p_T}{\text{GeV}})\%$ from Tevatron I, $\approx 15\%$ at HERA from MC

	10^{-2} GeV $^{3+2L}$	BK default	feed-down subtracted
$\langle \mathcal{O}(^1S_0^{[8]}) \rangle$	4.76 ± 0.71	3.1	
$\langle \mathcal{O}(^3S_1^{[8]}) \rangle$	0.265 ± 0.091	0.23	
$\langle \mathcal{O}(^3P_0^{[8]}) \rangle$	-1.32 ± 0.35	-0.82	

- Feed-down corrections \ll theoretical uncertainties

Comparison with MWC arXiv:1009.3655 (2)

Summary of M_0 and M_1 in (10^{-2} GeV 3) from BK and MWC

authors	data	feed-down	M_0	M_1
BK	default	—	2.5	0.59
BK	HERA $p_T > \sqrt{5}$ GeV	—	2.5	0.60
BK	Tevatron $p_T > 7$ GeV	—	1.9	0.54
BK	default	✓	1.7	0.43
BK	only Tevatron	—	8.7	0.62
BK	only Tevatron $p_T > 7$ GeV	—	9.3	0.30
MWC	only Tevatron $p_T > 5$ GeV	✓	5.2	0.16
MWC	only Tevatron $p_T > 7$ GeV	✓	7.4	0.05

- Lower p_T cuts on HERA or Tevatron data marginal in joint fit
- Feed-down corrections moderate \leadsto no qualitative change
- Exclusion of HERA data \leadsto fit greatly underconstrained

Comparison with MWC arXiv:1009.3655 (3)

Errors on M_0 and M_1 in 3-parameter fit to only Tevatron $p_T > 7$ GeV

Covariance matrix V_{ij} defined through

$$(V^{-1})_{ij} = \frac{1}{2} \frac{\partial^2 \chi^2}{\partial x_i \partial x_j}, \quad (x_1, x_2, x_3) = (\langle \mathcal{O}(^1S_0^{[8]}) \rangle, \langle \mathcal{O}(^3S_1^{[8]}) \rangle, \langle \mathcal{O}(^3P_0^{[8]}) \rangle)$$

$$\lambda_1 = 0.127: \quad \vec{v}_1 = (0.864, -0.114, -0.491)$$

has eigenvalues and eigenvectors: $\lambda_2 = 1.78 \times 10^{-6}: \quad \vec{v}_2 = (0.502, 0.0968, 0.860)$

$$\lambda_3 = 4.69 \times 10^{-8}: \quad \vec{v}_3 = (0.0507, 0.989, -0.141)$$

Cf. $\vec{v}_{M_0} = (0.500, 0, 0.866)$ and $\vec{v}_{M_1} = (0, 0.970, -0.242)$ from 2-parameter fit of MWC

$$\Delta M_0 = (\vec{v}_{M_0}^T V \vec{v}_{M_0})^{1/2} / (\vec{v}_{M_0})_1 = 0.95 \times 10^{-2}$$

$$\Delta M_1 = (\vec{v}_{M_1}^T V \vec{v}_{M_1})^{1/2} / (\vec{v}_{M_1})_2 = 0.29 \times 10^{-2}$$

i.e. 3-parameter fit yields: $M_0 = (9.3 \pm 0.95)$ and $M_1 = (0.30 \pm 0.29)$ in 10^{-2} GeV 3

~~ almost 100% error on M_1 in 3-parameter fit

Cf. $M_0 = (7.4 \pm 1.9)$ and $M_1 = (0.05 \pm 0.02)$ from 2-parameter fit by MWC

- M_1 corresponds mostly to \vec{v}_3 , but contains small admixtures of \vec{v}_1 and \vec{v}_2
- \vec{v}_1 very badly constrained ~~ large error on M_1
- Not exhibited in 2-parameter fit, where variations orthogonal to M_0 and M_1 are forbidden

Comparison with MWC arXiv:1009.3655 (4)

Errors on $\langle \mathcal{O}(^1S_0^{[8]}) \rangle$, $\langle \mathcal{O}(^3S_1^{[8]}) \rangle$ and $\langle \mathcal{O}(^3P_0^{[8]}) \rangle$ in 3-parameter fits

$10^{-2} \text{ GeV}^{3+2L}$	BK default	Tevatron $p_T > 7 \text{ GeV}$ only
$\langle \mathcal{O}(^1S_0^{[8]}) \rangle$	4.76 ± 0.71	25.1 ± 30.7
$\langle \mathcal{O}(^3S_1^{[8]}) \rangle$	0.265 ± 0.091	-1.96 ± 4.07
$\langle \mathcal{O}(^3P_0^{[8]}) \rangle$	-1.32 ± 0.35	-9.09 ± 17.48

Correlation matrix V_{ij} of default fit has eigenvalues and eigenvectors:

$$\lambda_1 = 6.39 \times 10^{-5}: \quad \vec{v}_1 = (0.893, -0.111, -0.435)$$

$$\lambda_2 = 1.92 \times 10^{-7}: \quad \vec{v}_2 = (0.440, 0.0193, 0.898)$$

$$\lambda_3 = 4.00 \times 10^{-8}: \quad \vec{v}_3 = 0.0910, 0.994, -0.0659$$

- No such strong hierarchy among eigenvalues
- Fit results meaningful

Summary (1)

Our project: Test NRQCD

- NRQCD provides rigorous **factorization theorem** for production and decay of heavy quarkonia: Include **color-octet** (CO) states.
 - But: Need to proof **universality** of CO LDMEs.
 - **This work:** Technological breakthrough: After **14 years** finally NLO NRQCD photo- and hadroproduction calculation.

Our Results:

- **CSM predictions:** Could verify all previous results:
CS contributions **far below data** in all considered experiments.
 - **Fitted** CO LDMEs to p_T distributions at Tevatron and HERA.
 - Used LDMEs for RHIC, LHC and HERA W and z distributions
 \Rightarrow CS+CO: Good **agreement** with data.

Summary (2)

Discussion of Fit:

- NLO hadroproduction P states: Shape changes, even negative.
 \Rightarrow Fit all three CO LDMEs (not linear combination like at LO).
 - Negative unphysical quantities no problem.
 - Photo- and hadroproduction consistently described by NRQCD.

Still to be done:

- Include resolved photoproduction.
 - Extend analysis to e^+e^- collisions (LEP, B factories).
 - Include feed-down processes.
 - Do polarization analysis.