


# $F_2^{bb}$ from muon+jet final states at ZEUS


B. Kahle, DESY Hamburg

- Introduction
- Beauty measurements
- Beauty contribution to  $F_2$
- Summary and Conclusions



# Motivation


- Heavy flavour production in DIS is a test of pQCD providing an additional hard scale  $M$  to the momentum transfer of the boson  $Q$  and  $p_t$
- Beauty contribution to  $F_2$  is directly sensitive to the gluon distribution in the proton
- Beauty production is of increasing interest for higher energies -> LHC




# Beauty contribution to the proton structure function $F_2$

$$\frac{d^2\sigma}{dx dQ^2} = \frac{2\pi\alpha^2}{Q^4x} \left\{ \left[ 1 + (1-y)^2 \right] F_2(x, Q^2) - y^2 F_L(x, Q^2) + \dots xF_3 \right\}$$

~negligible  
at high  $Q^2$




$$\frac{d^2\sigma^{ep}}{dQ^2 dx} \propto F_2(x, Q^2)$$



$$\frac{d^2\sigma^{ep \rightarrow b\bar{b}X}}{dQ^2 dx} \propto F_2^{b\bar{b}}(x, Q^2)$$

# Heavy Flavour production mechanism

Dominant process in  $ep$ -collisions: **Boson-Gluon-Fusion**



## Kinematic variables:

$Q^2 = -q^2$  photon virtuality, squared momentum transfer

$x = \frac{Q^2}{2Pq}$  Bjorken scaling variable, for  $Q^2 \gg (2m_Q)^2$ : momentum fraction of  $p$  constituent

## Kinematic regime:

- Deep inelastic scattering (**DIS**):  $Q^2 > 1 \text{ GeV}^2$

## Multiple scales:

$m_b \sim 5 \text{ GeV}$

$p_t^b \sim \text{typically few GeV}$

$Q^2 \gtrsim 1 \text{ GeV}^2$  in DIS

→ different pQCD approaches


good testing ground for pQCD

# pQCD approximations

## Massive scheme:

- $b$  massive
- neglects  $[\alpha_s \ln(Q^2/m_b^2)]^n$
- scale  $m_b, p_t$


↳  $b$  produced perturbatively  
(not part of the Proton or Photon)



## Massless scheme:

- $b$  massless
- resumes  $[\alpha_s \ln(Q^2/m_b^2)]^n$
- scale:  $Q^2, p_t$

↳  $b$  also in Proton and Photon




## Variable flavour number scheme (VFNS):

- massive at small  $Q^2$
- massless at large  $Q^2$

# Beauty identification

Process :  $e p \rightarrow e b \bar{b} X \rightarrow e \mu \text{ jet } X'$

$p_t^{\text{rel}}$  method:




$p_t^{\text{rel}}$  is the momentum of the muon transverse to the axis of the associated jet (including the muon)

$p_t^{\text{rel}}$  spectrum is **harder** for **b** than for **c**

→ statistical separation using MC

**ZEUS**



$\chi^2$  fit of b MC against c+l f MC to the data in  $p_t^{\text{rel}}$

resulting beauty fraction of about 21%

→ scale Rapgap-b MC up by a factor 2.49

# Cross section

Beauty cross section for the DIS process:

$$e p \rightarrow e b \bar{b} X \rightarrow e \mu \text{ jet } X'$$

Cuts on  $MC_{\text{true}}$  quantities:

$$Q^2 > 4 \text{ GeV}^2$$

$$0.05 < y < 0.7$$

$$E_{t,\text{jet}}^{\text{lab}} > 5 \text{ GeV}$$

$$-2 > \eta_{\text{jet}} > 2.5$$

$$p_{t,\mu} > 1.5 \text{ GeV}$$

$$-1.6 > \eta_{\mu}$$

$$\text{Data: } \sigma_{b\bar{b}} = 77.1 \pm 7.8 \text{ (stat.) } \pm_{14.9}^{9.6} \text{ (sys.) pb}$$

$$\text{MC (LO+PS): } \sigma_{b\bar{b}} = 31.0 \text{ pb}$$

$$\text{NLO (HVQDIS): } \sigma_{b\bar{b}} = 32.9 \pm 3.3 \text{ (sys.) pb}$$





$$\mu_{R,F}^2 = p_t^2 + 4m_b^2 \quad (\text{simultaneous variation: factor } 1/4 \text{ to } 4)$$

$$m_b = 4.75 \text{ GeV} \quad (\text{variation: } 4.5 \text{ GeV to } 5 \text{ GeV})$$

$$\text{Peterson frag. function } \epsilon = 0.0035$$





Proton parton distribution: CTEQ5F4

# Cross sections in $p_t$ and $\eta$





# Cross sections in $Q^2$ and $Q^2x$



# Calculation of

## $F_2^{b\bar{b}}$ = beauty contribution to $F_2$

„Reduced cross section“ is defined as:

$$\tilde{\sigma}_{NLO}^{b\bar{b}}(x, Q^2) = \frac{d^2 \sigma_{NLO}^{b\bar{b}}}{d x d Q^2} \frac{x Q^4}{2 \pi \alpha^2 (1 + (1 - y)^2)}$$

Cross section calculated using `hvgdis (CTEQ5F4)` for a tiny bin around  $(x, Q^2)$

Expected to be compatible with calculations by Riemersma *et al.* used for  $F_2^{cc}$

Neglecting the small contribution from  $F_L$ , the reduced cross section is equal to  $F_2$ :

$$\tilde{\sigma}^{b\bar{b}}(x, Q^2) = F_2^{b\bar{b}} - \frac{y^2}{1 + (1 - y)^2} F_L^{b\bar{b}}$$

*neglected*

# $F_2^{b\bar{b}}$ measurement

The reduced cross section for data is the reduced cross section of the NLO multiplied by the ratio of data to NLO in a  $x, Q^2$  bin:

$$\tilde{\sigma}_{data}^{b\bar{b}}(x, Q^2) = \tilde{\sigma}_{NLO}^{b\bar{b}}(x, Q^2) \frac{d^2 \sigma_{data}^{b\bar{b} \rightarrow \mu}}{dx dQ^2} / \frac{d^2 \sigma_{NLO}^{b\bar{b} \rightarrow \mu}}{dx dQ^2}$$


Cross section for  
 $e p \rightarrow e b\bar{b} X \rightarrow e \mu jet X'$

NLO using  
hvqdis with same  
settings as for  $\tilde{\sigma}^{b\bar{b}}$   
(but requiring SL  
decay to  $\mu$  and jet)

H1 uses the impact parameter method to measure  $F_2^{b\bar{b}}$  and  $F_2^{c\bar{c}}$  with an inclusive charm and beauty sample of  $57\text{pb}^{-1}$ :


H1 Collab., A. Aktas et al., Eur. Phys. J. C45 (2006) 23-33

# Kinematic plane (ZEUS)



●  $(x, Q^2)$  values  
chosen for  $F_2^{bb}$   
to compare with  
H1's results

# $F_2^{bb}$ at ZEUS and H1



ZEUS data lie above H1 data but compatible within errors.


HVQDIS+CTEQ5F4 agrees with similar predictions by H1

■ ZEUS: 39 pb<sup>-1</sup>  
● ▼ H1: 57.4 pb<sup>-1</sup>

Theory predictions except HVQDIS+CTEQ5F4 provided by P.D.Thompson, hep-ph/0703103

ZEUS data point at  $Q^2=200\text{GeV}^2$ ;  $x=0.13$  is shifted to lower  $x$  value to be separated from the H1 point

# $F_2^{bb}$ vs. $Q^2$



Same ZEUS data. Some points are recalculated for other values of  $x$  to be comparable with H1 data and theory curves.

ZEUS data are derived using HVQDIS+CTEQ5F4 and should strictly compared only with this curve.

For comparison with other curves, model dependencies have to be taken into account.


Some ZEUS data point are shifted to higher  $Q^2$  values to be separated from the H1 point

# PDF Schemes and Parameters

PDF	Order	Scheme	$\mu_M^2$	$M_b(\text{GeV})$
- · MRST04	$\alpha_s^2$	VFNS	$Q^2$	4.3
— MRST NNLO	$\alpha_s^3$	VFNS	$Q^2$	4.3
..... CTEQ6HQ	$\alpha_s^2$	VFNS	$Q^2$	4.5
- - HVQDIS+CTEQ5F4	$\alpha_s^2$	FFNS	$p_t^2+4M^2$	4.75
CTEQ5F3	$\alpha_s^2$	FFNS	$Q^2$	4.5
MRST FF3	$\alpha_s^2$	FFNS	$Q^2$	4.3
CTEQ6.5	$\alpha_s^2$	VFNS	$Q^2+M^2$	4.5

Theory predictions except HVQDIS+CTEQ5F4  
provided by P.D.Thompson, hep-ph/0703103

# $F_2^{bb}$ at ZEUS and H1



Same plot as before, but with additional theory curves.

■ ZEUS: 39 pb<sup>-1</sup>  
● ▼ H1: 57.4 pb<sup>-1</sup>

Theory predictions except HVQDIS+CTEQ5F4 provided by P.D.Thompson, hep-ph/0703103

ZEUS data point at  $Q^2=200\text{GeV}^2$ ;  $x=0.13$  is shifted to lower  $x$  value to be separated from the H1 point



# Summary and Outlook

---

- First measurement of  $F_2^{bb}$  at ZEUS ( $39 \text{ pb}^{-1}$ ),  
~10 times more data to come -> much reduced errors
- Results agree with H1's using a very different method to obtain  $F_2^{bb}$  but similar uncertainties (both statistical and systematical)
- NLO predictions agree with data within large spread.

Questions to this meeting:

What are the reasons for the large NLO spread?

Does this affect the PDF extraction from inclusive  $F_2$ ?

# BACKUP

---


# NLO calculations

---

The calculation of the NLO QCD visible cross section predictions proceeds in three steps:

- **HVQDIS** (B.Harris, J.Smith, hep-ph/9503484):  
 $\gamma^*g \rightarrow bb$ ,  $\gamma^*g \rightarrow bbg$ ,  $\gamma^*q \rightarrow bbq$ , etc. (pointlike only)  
using **CTEQ5F4** (FFNS) PDF
- **Fragmentation of the **b-quark** into a **B-meson****  
(Peterson function with  $\epsilon=0.0035$ )
- **Semileptonic decay of the B-meson**  
(Muon momentum spectrum extracted from RAPGAP, including primary and secondary muons)

# CTEQ5F4 with different scales



$F_2^{bb}$  using different scales for CTEQ5F4:

$Q^2$

$Q^2+4M^2$  (M=4.5 GeV)

$Q^2+4M^2$  (M=4.75 GeV)

large differences at low  $Q^2$  for different masses

# Extrapolation to full phase-space

$Q^2$	Extrapolation factor:
25 GeV <sup>2</sup>	~6
110 GeV <sup>2</sup>	~4
200 GeV <sup>2</sup>	~3

Similar to extrapolations for  $F_2^{cc}$

Extrapolation factor excluding branching fraction to  $\mu$  of 0.3924

Extrapolation factor includes  $p_t^b$  and  $\eta^b$  spectrum, fragmentation, and decay kinematics (jet and  $\mu$ )

# Inclusive lifetime tags

most significant impact parameter  $S_1$

2<sup>nd</sup> most significant impact parameter  $S_2$

H1

