



# Prospects for the observation of new physics at the LHC



Tracey Berry




University of Royal Holloway, London  
Flavour in the era of the LHC, CERN  
27<sup>th</sup> March 2007



# Overview



- Beyond the Standard Model
- Fundamental Symmetries
  - Heavy Gauge Bosons  $W'$  and  $Z'$
- Electroweak Symmetry Breaking
  - SUSY, Little Higgs, Technicolor
- Leptons & Quarks, Other New Particles
  - Left-right symmetry, E6 quarks
- Extra Dimensions
- Summary and Outlook





## Questions remaining with the Standard Model



→ Reasons to search for new physics...

- **Fundamental symmetries:**
  - Are there more symmetries beyond  $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ ?  
→ GUTs with larger symmetry group? Left-right symmetry?
- **ElectroWeak Symmetry Breaking (EWSB):**
  - Unitarity violation in longitudinal WW scattering at high E  
*solution:* Higgs boson or other new particle with mass  $< 1$  TeV
  - If Higgs → *hierarchy problem*: fine tuning in rad corr to Higgs mass  
*solution:* new physics at TeV scale (SUSY, Little Higgs, etc...)
  - If NO Higgs  
*solution:* new strong interactions (Technicolor, etc...)
- **Quark and lepton generations:**
  - Why are there 3 generations? → Fermions composite?
  - Is there a lepto(n)-quark symmetry?
  - More than 3 generations of quarks & leptons?





# Heavy Gauge Bosons



- Many extensions of the SM rely on larger symmetry groups (GUTs, string-inspired, left-right, little Higgs models, etc...)
  - predict existence of new gauge bosons  $W'$  and  $Z'$  (or KK modes)

- Production: s-channel

- Clean decay channels:


$$W' \rightarrow e \nu_e \text{ or } \mu \nu_\mu$$

$$Z' \rightarrow e^+e^- \text{ or } \mu^+\mu^-$$

- Tevatron searches:  $M$  up to  $\sim 1$  TeV


- $Z'$  models considered:


- Sequential SM (SSM) with same  $Z'$  couplings to fermions as for  $Z$
- Models based on different patterns of  $E_6$  symmetry breaking ( $\psi$ ,  $\chi$  and  $\eta$ )
- Left-right (LR) symmetry models



$Z' 1.5\text{TeV}$

T. Rizzo, hep-ph/0610104v1







# Heavy New Gauge Bosons $Z'$




- **Selection:** pairs of isolated  $e$  or  $\mu$
- **Bkg:** dominated by dileptons from Drell-Yan
- $5\sigma$  discovery up to  $\sim 5$  TeV (model dependent) for both ATLAS and CMS


CMS PTDR  $Z' \rightarrow ee$  (SSM)



A very low luminosity,  $< 0.1 \text{ fb}^{-1}$ , should be sufficient to discover  $Z'$  bosons at 1 TeV For all models and in a single channel ( $e/m$ )

$Z'$   $5\sigma$  reach  $ee$  vs.  $\mu\mu$  channels









# Z' ATLAS reaches

ATLAS discovery reaches in CDDT models of Z'





L=400pb<sup>-1</sup> : first months of running at LHC

L=100fb<sup>-1</sup>




$g_{Z'}$  = global coupling strength

x= describes relative coupling strength to different fermions



With only 400 pb<sup>-1</sup> : can go beyond LEP exclusion limits and probe regions of parameter space not yet excluded by CDF.

ATL-PHYS-PUB-2006-024



# Discrimination between $Z'$ models



- Models differ in the  $Z'$  couplings to fermions especially parity-violating couplings to leptons + couplings to initial u/d
- Distinguish by measuring:

- $\sigma \times \Gamma$ : Decay width (ee only due to worse  $\mu\mu$  resolution)
- Forward-backward asymmetry
- $Z'$  rapidity

ATL-PHYS-PUB-2005-010


→ Also provides discrimination against other models like extra-D, little Higgs, ...

In one year at  $10^{34}$ : reach 4-5 TeV & discriminating between models possible up to  $m \sim 2.5$  TeV


$M = 1.5$  TeV  $100 \text{ fb}^{-1}$

Model	$\sigma_{ll} \times \Gamma_{ll}$ (fb x GeV)	Corrected $A_{FB}$ at $Z'$ peak
SSM	$3668 \pm 138$	$+0.108 \pm 0.027$
$\chi$	$828 \pm 48$	$-0.361 \pm 0.030$
LR	$1515 \pm 75$	$+0.186 \pm 0.032$


Dilepton invariant mass spectrum




Forward backward asymmetry measurement



Rapidity distribution








# Heavy Gauge Bosons: $W'$



- General Model by Altarelli, Mele, Ruiz-Altaba with same  $W'$  couplings to fermions as for  $W$
- Selection: one-muon event with track isolation req<sup>t</sup> around  $\mu$  + missing transverse energy
- Background: mostly  $W \rightarrow \mu \nu$









# SUSY Spectrum

- SUSY gives rise to partners of SM states with opposite spin-statistics but otherwise same Quantum Numbers.  
 spin-1/2 matter particles (fermions)  $\Leftrightarrow$  spin-1 force carriers (bosons)
- Expect SUSY partners to have same masses as SM states
  - Not observed
  - SUSY must be a broken symmetry
- Different mechanisms of SUSY breaking lead to different models  
 MSSM, mSugra, MSSM, GMSB, AMSB
  - R-Parity  $R_p = (-1)^{3B+2S+L}$
  - Conservation of  $R_p$  causes LSP to be stable






## LHC SUSY Searches




Typical Signature/decay chain:



- Strongly interacting sparticles (squarks, gluinos) dominate production.
- Heavier than sleptons, gauginos etc. g cascade decays to LSP.
- Potentially long decay chains and large mass differences
  - **Many high  $p_T$  objects observed (leptons, jets, b-jets).**
- If R-Parity conserved LSP (lightest neutralino in mSUGRA) stable and sparticles pair produced.
  - **Large  $E_T^{\text{miss}}$  signature (c.f.  $W_{\text{glv}}$ ).**

### SUSY Searches

- **Inclusive** searches to detect SUSY with first data
- **Exclusive** studies – performed with more data to determine model parameters e.g. masses etc from end point measurements...





# SUSY discovery with first data




**Inclusive** searches to detect SUSY with first data

Look for deviations from SM predictions – requires good knowledge of SM processes at that energy scale and on an understanding of the detectors' performances

Best reach obtained with the most inclusive channels:




The range of gluino and squark masses up to about 1.5 TeV can be probed with  $L=1\text{fb}^{-1}$  and up to about 2 TeV with  $10\text{fb}^{-1}$




# LHC: Jets + $E_T^{\text{miss}}$




- Inclusive searches with Jets + n leptons +  $E_T^{\text{miss}}$  channel.
- Map statistical discovery reach in mSUGRA  $m_0$ - $m_{1/2}$  parameter space.
- Sensitivity only weakly dependent on  $A_0$ ,  $\tan(\beta)$  and  $\text{sign}(\mu)$ .



“Golden channel at the LHC”







# Exclusive SUSY Searches




- **Exclusive** studies – reconstruct specific decay channels in order to estimate physical parameters characterising the decay:
  - e.g. reconstructing kinematic endpoints (edges & thresholds) in invariant mass distributions to extract masses

Simultaneous observation of a signal in various topologies would allow measurements of cross-sections of sub-processes and their ratios and help to determine the underlying physics.



Dilepton channel is the "golden channel" for exclusive studies

Cleanest signature is the opposite sign same flavour invariant mass distributions of lepton (e/ $\mu$ ) pairs. **From its endpoints can extract precise constraints on difference in masses between s-particles in decay chain with a statistical precision of 0.1% or better with  $L=100\text{fb}^{-1}$**



# EWSB: Little Higgs




- Models with Higgs as pseudo-Goldstone boson from a broken global symmetry (SU(5) in "littlest Higgs model")
  - Extra Q=2/3 heavy quark (T) and heavy gauge bosons ( $A_H, W_H, Z_H$ )
  - Quadratic divergences cancel top and VB divergences to Higgs mass


## Heavy Quark T

- Production: via QCD ( $gg \rightarrow T \bar{T}, q\bar{q} \rightarrow T \bar{T}$ )  
via W exchange ( $qb \rightarrow q' T$ ) dominant for  $M_T > 700$  GeV
- Decays:  $T \rightarrow t Z, T \rightarrow t H, T \rightarrow b W$

Parameters:  $\lambda_1, \lambda_2$  & f determine masses of T, t-quark & their couplings

- cleanest is  $T \rightarrow t Z \rightarrow b l \nu l^+ l^-$   $\longrightarrow$   
main bkg is tbZ  
5 $\sigma$  signal up to  $\sim 1.0-1.4$  TeV  $\lambda_1/\lambda_2=1(2)$
- $T \rightarrow t H \rightarrow b l \nu \bar{b} b < 5\sigma$
- $T \rightarrow b W \rightarrow b l \nu$   
main bkg is t t  
5 $\sigma$  signal up to  $\sim 2.0-2.5$  TeV  $\lambda_1/\lambda_2=1(2)$







# Little Higgs




- $A_H$  and  $Z_H$  discovery in lepton modes

Signal: dilepton resonance






- Reach up to  $M \sim 6$  TeV (depending on mixing angle  $\cot \theta$ )
- New gauge bosons observable over almost the entire mass range




# Little Higgs

- Discrimination against other models predicting dilepton resonances via observation of decay modes like  $W_H \rightarrow W H$ ,  $Z_H \rightarrow Z H$ , and  $W_H \rightarrow t \bar{b}$  (important at  $\cot \theta \approx 1$ , where  $\text{BR}(W_H/Z_H \rightarrow W/Z H)$  vanishes)









# Dynamical EWSB: Technicolor



- Dynamical EWSB via new strong interaction
  - No need for Higgs boson → removes fine tuning problem
  - Predict new technifermions, technihadrons
- Study  $\rho_{TC} \rightarrow W Z$  process
- Select isolated leptons, measure missing  $E_T$  & apply W and Z kinematical constraints
- Bkg: WZ, ZZ, Zbb, t $\bar{t}$




5 $\sigma$  sensitivity contours




Flavour Workshop, CERN

Tracey Berry (RHUL)



CMS PTDR 2006



# Doubly Charged Higgs in the L-R Symmetric Model

- Left-Right Symmetric Model based on  $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ 
  - Features triplet of Higgs fields ( $\Delta_R^0, \Delta_R^+, \Delta_R^{++}$ ) + two doublets  $\phi$
  - Predicts new gauge bosons ( $W_R$  and  $Z_R$ ) & new fermions (heavy Majorana  $\nu, \nu_R$ )
  - Addresses origin of pure left-handed charged weak interaction + origin of light neutrino masses (via see-saw mech. & heavy  $\nu_R$ )



- Production:  $q\bar{q} \rightarrow q'q' W_{R,L}^+ W_{R,L}^+ \rightarrow q'q' \Delta_{R,L}^{++}$   
 $q\bar{q} \rightarrow \gamma^*/Z/Z_{R,L} \rightarrow \Delta_{R,L}^{++} \Delta_{R,L}^{--}$


$$g_R = g_L$$

$$m(W_R) = g_R v_R / \sqrt{2}$$

SN-ATLAS-2005-049

- Decay:  $\Delta_{R,L}^{++} \rightarrow l^+ l^+$
- Selection (WW fusion):  
2 like-sign leptons ( $e, \mu, \tau$ )  
+ "forward" jets
- Bkg:  $W^+ W^+ q q, W t \bar{\tau}$







# Doubly Charged Higgs

## in the L-R Symmetric Model




- Selection ( $\Delta_{R,L}^{++} \Delta_{R,L}^{--} \rightarrow 4l$  process):  
2 pairs of like-sign leptons ( $e, \mu, \tau$ )
- Bkg: negligible
- Contours for 10 signal events

SN-ATLAS-2005-049



Reach improves if only 3 leptons are required (solid lines)



→  $\Delta_R^{++}$  Mass reach 0.8 - 1.2 TeV (100 fb<sup>-1</sup>)  
0.9 - 1.4 TeV (300 fb<sup>-1</sup>)




# $W_R$ and Majorana Neutrinos




- Left-right Symmetric Model
- Signature:  
 lepton + 2 jets for heavy neutrino  $N_1$   
 dilepton + 2 jets for  $W_R$



CMS NOTE 2006/098



Even after just 1 month of running at low lum. can achieve  $W_R=2$  TeV  $N_e$  500 GeV/c<sup>2</sup>



# Leptoquarks



SM extension: lepton-quark symmetry

Generic prediction of GUTs, composite models, technicolor schemes, superstring-inspired  $E_6$  models & SUSY with R-parity violation

⇒ LQ: color triplets with couplings to quarks and leptons

Study: **Pair Production of Scalar LQ**

$LQ\bar{L}\bar{Q} \rightarrow l^+q l^-q$

**Signature:** 2jets+2 leptons: 1<sup>st</sup> and 2<sup>nd</sup> gen.

Sensitivity:  $M_{LQ} \sim 1.3 \text{ TeV } \beta=1$

$M_{LQ} \sim 1 \text{ TeV } \beta=0.5 \text{ L}=30\text{fb}^{-1}$



$\beta = \text{BR}(LQ \rightarrow lq)$


$LQ\bar{L}\bar{Q} \rightarrow \nu_\tau b \bar{\nu}_\tau \bar{b}$

**Signature** 2 jets +  $E_t$  : 3<sup>rd</sup> generation

Sensitivity:  $M_{LQ} \sim 1 \text{ TeV } \beta=0.5$

$M_{LQ} \sim 1.3 \text{ TeV } \beta=0$





# Excited Quarks



If quarks & leptons have substructure...  
 there should exist new interactions among them at the scale of  
 the constituents binding energies

→ a possible consequence is the existence of excited quarks/leptons  
 with masses of order the compositeness scale  $\Lambda$ .


## Excited Quarks

**Production:** quark-gluon fusion

**Decay Signature:** photons+jets  
 quarks + gauge bosons  
 quarks + gluons


$L=300\text{fb}^{-1}$ ,  $\Lambda=m^*$ ,  $f=f'=1$

- Reach limit for  $q^* \rightarrow q\gamma$ : 6.5 TeV
- Reach limit for  $q^* \rightarrow qW$ : 7 TeV
- Reach limit for  $q^* \rightarrow qZ$ : 4.5 TeV
- Reach limit for  $q^* \rightarrow gq$ : 6 TeV



$f$  = Coupling determined by the compositeness dynamics

ATL-PHYS-PUB-2006-027






# Excited Leptons



## Excited Electrons

At  $E \ll \Lambda$  compositeness scale  $\Lambda$ , quark-lepton interactions can be effectively approximated by contact interactions


- Production and decay:  $qq \rightarrow e^*e \rightarrow Zee$   
&  $Z \rightarrow ee$  or  $jj$
- Primary backgrounds:  $ZZ, Z + \text{photon}$



$f = f' = 1$

Reach  $\sim 3\text{-}4 \text{ TeV}$  for  $L = 6 \text{ TeV } 300 \text{ fb}^{-1}$


ATLAS-PHYS-2002-014





# Heavy Quarks




- Symmetry group  $E_6$  favored by string-inspired GUTs (supergravity)
  - Predicts new  $Q=-1/3$  quark
- Production:  $gg \rightarrow D\bar{D}$  (dominant for  $M_D < 1.1$  TeV)  
 $\bar{q}q \rightarrow D\bar{D}$  (dominant for  $M_D > 1.1$  TeV)
- Decay:  $D \rightarrow W u$  or  $D \rightarrow Z d$  (for this study)
- Selection: 4 leptons (from Z)  
 + 2 jets



SN-ATLAS-2006-056







# Extra Dimensions: ADD Model



Arkani-Hamed, Dimopoulos, Dvali, Phys Lett B429 (98), Nuc.Phys.B544(1999)



(Many) Large flat Extra-Dimensions (LED),

could be as large as a few  $\mu\text{m}$

G can propagate in ED

SM particles restricted to 3D brane

The fundamental scale is not planckian:  $M_D = M_{\text{Pl}(4+\delta)} \sim \text{TeV}$



Model parameters are:

- $\delta$  = number of ED

$$M_{\text{Pl}}^2 \sim R^\delta M_{\text{Pl}(4+\delta)}^{(2+\delta)}$$


- $M_{\text{Pl}(4+\delta)}$  = Planck mass in the  $4+\delta$  dimensions


Signature:

- ❖ Real graviton emission: **in association with a vector-boson**

**jets + missing ET, V+missing ET**

- ❖ Or deviations in virtual graviton exchange  
**e.g. Excess above di-lepton continuum**





# ADD Discovery Limit: Real G Emission



J. Weng et al. CMS NOTE 2006/129

$$pp \rightarrow \gamma + G^{KK}$$


**Signature:** high- $p_T$  photon + high missing  $E_T$

**Main Bkgd:** irreducible  $Z\gamma \rightarrow \nu\nu\gamma$ ,  
Also  $W \rightarrow e(\mu, \tau)\nu$ ,  $W\gamma \rightarrow e\nu, \gamma + \text{jets}$ ,  
QCD, di- $\gamma$ ,  $Z^0 + \text{jets}$

$M_D =$ 1 – 1.5 TeV for 1 fb <sup>-1</sup>
2 – 2.5 TeV for 10 fb <sup>-1</sup>
3 – 3.5 TeV for 60 fb <sup>-1</sup>



Rates for  $M_D \geq 3.5$  TeV are very low – too low for 5 $\sigma$  discovery



$M_D^{\text{MAX}}$ (TeV)	$\delta=2$
HL 100fb <sup>-1</sup>	4





$$pp \rightarrow \text{jet} + G^{KK}$$

**Signature:** high  $E_T$  jet + large missing  $E_T$   
**Bkgd:** irreducible jet+Z/W  $\rightarrow$  jet+ $\nu\nu$  /jet+l $\nu$   
vetos leptons: to reduce jet+W bkgd mainly

Discovery limits

$M_{\text{Pl}(4+d)}^{\text{MAX}}$ (TeV)	$\delta=2$	$\delta=3$	$\delta=4$
LL 30fb <sup>-1</sup>	7.7	6.2	5.2
HL 100fb <sup>-1</sup>	9.1	7.0	6.0






# ADD Discovery Limit: G Exchange



$$pp \rightarrow G^{KK} \rightarrow \mu\mu$$



- Two opposite sign muons &  $M_{\mu\mu} > 1$  TeV
- **Bkg**: Irreducible Drell-Yan, also ZZ, WW, WW, tt (suppressed after selection cuts)




1 fb<sup>-1</sup>: 3.9-5.5 TeV for n=6..3  
 10 fb<sup>-1</sup>: 4.8-7.2 TeV for n=6..3  
 100 fb<sup>-1</sup>: 5.7-8.3 TeV for n=6..3  
 300 fb<sup>-1</sup>: 5.9-8.8 TeV for n=6..3

channel	n		2	3	4	5
 $\gamma\gamma$	luminosity					
	10 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	6.3	5.6	5.1	4.9
		S/B	36/18	36/18	39/25	34/13
	100 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	7.9	7.3	6.7	6.3
		S/B	50/53	62/96	55/72	51/53
$l^+l^-$	10 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	6.6	5.9	5.4	5.1
		S/B	33/11	31/8	30/6	30/6
	100 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	7.9	7.5	7.0	6.6
		S/B	49/48	38/21	36/16	29/6
Fast MC	10 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	7.0	6.3	5.7	5.4
	$\gamma\gamma + l^+l^-$	100 fb <sup>-1</sup>	$M_S^{max}$ (TeV)	8.1	7.9	7.4

V. Kabachenko et al.  
ATL-PHYS-2001-012


Belotelov et al.,  
CMS NOTE 2006/076, CMS PTDR 2006




# RS Model



1 extra warped dimension




**Signature:**  
Narrow, high-mass resonance states in dilepton/dijet/diboson channels




## Model parameters:

- Gravity Scale:  $\Lambda_\pi = \overline{M}_{pl} e^{-kR_c\pi}$
  - 1<sup>st</sup> graviton excitation mass:  $m_1 \rightarrow$  **Resonance position**
  - $\Lambda_\pi = m_1 \overline{M}_{pl} / kx_1$ , &  $m_n = kx_n \overline{M}_{pl} (J_1(x_n) = 0)$
  - Coupling constant:  $c = k/\overline{M}_{pl}$
  - $\Gamma_1 = \rho m_1 x_1^2 (k/\overline{M}_{pl})^2 \rightarrow$  **width**
- $k =$  curvature,  $R =$  compactification radius

Best channels to search in are   $G(1) \rightarrow e^+e^-$  and  $G(1) \rightarrow \gamma\gamma$  due to the energy and angular resolutions of the LHC detectors


$G(1) \rightarrow e^+e^-$  best chance of discovery due to relatively small bkgd, from Drell-Yan

Allenach et al, hep-ph0211205 Allenach et al, hep-ph0006114



# RS Model



1 extra warped dimension



**Signature:**  
Narrow, high-mass resonance states in dilepton/dijet/diboson channels

### Model parameters:

- Gravity Scale:  $\Lambda_\pi = \overline{M}_{pl} e^{-kR_c\pi}$
  - 1<sup>st</sup> graviton excitation mass:  $m_1 \rightarrow$  **Resonance position**
  - $\Lambda_\pi = m_1 \overline{M}_{pl} / kx_1$ , &  $m_n = kx_n \overline{M}_{pl} e^{krc\pi} (J_1(x_n) = 0)$
  - Coupling constant:  $c = k/\overline{M}_{pl}$
  - $\Gamma_1 = \rho m_1 x_1^2 (k/\overline{M}_{pl})^2 \rightarrow$  **width**
- $k =$  curvature,  $R =$  compactification radius




Best channels to search in are  $G(1) \rightarrow e^+e^-$  and  $G(1) \rightarrow \gamma\gamma$  due to the energy and angular resolutions of the LHC detectors




$G(1) \rightarrow e^+e^-$  best chance of discovery due to relatively small bkgd, from Drell-Yan

I. Belotelov et al. , CMS NOTE 2006/104, CMS PTDR 2006  
Allenach et al, hep-ph0211205 Allenach et al, hep-ph0006114



# RS1 Discovery Limit




$$\text{BR}(G \rightarrow \gamma\gamma) = 2 * \text{BR}(G \rightarrow ee/\mu\mu)$$


Reach for e and  $\gamma$  is comparable for low coupling and G mass due to the QCD and prompt photon bkgds in the  $\gamma\gamma$  channel which are harder to efficiently suppress  $\mu\mu$  channel trails ee channel due to resolution

< 60 fb<sup>-1</sup> LHC completely covers the region of interest


$G_1 \rightarrow \mu^+\mu^-$




$G_1 \rightarrow e^+e^-$



$G_1 \rightarrow \gamma\gamma$



$c > 0.1$  disfavoured as bulk curvature becomes to large (larger than the 5-dim Planck scale)




# RS1 Model Determination



How could a RS G resonance be distinguished from a Z' resonance?



Potentially using Spin information:

G has spin 2:  $pp \rightarrow G \rightarrow ee$  has 2 components:  $gg \rightarrow G \rightarrow ee$  &  $q\bar{q} \rightarrow G \rightarrow ee$ : each with different angular distributions:



$$qq \rightarrow G \rightarrow ff: 1 - 3 \cos^2 \theta + 4 \cos^4 \theta$$


$$gg \rightarrow G \rightarrow ff: 1 - \cos^4 \theta$$



Spin-2 could be determined (spin-1 ruled out) with 90% C.L. up to  $M_G = 1720 \text{ GeV}$  with  $100 \text{ fb}^{-1}$

Note: acceptance at large pseudo-rapidities is essential for spin discrimination ( $1.5 < |\eta| < 2.5$ )

Allanach et al, hep-ph 0006114



# TeV<sup>-1</sup> Sized Extra Dimensions



I. Antoniadis, PLB246 377 (1990)


- One extra dimension compactified on a S<sup>1</sup>/Z<sup>2</sup> orbifold
- Radius of compactification small enough -> Gauge bosons can travel in the bulk
- Fermions (quarks/leptons) localized
  - at a fixed point (M1) or
  - opposite (M2) points
 => destructive (M1) or constructive (M2) interference of the KK excitations with SM model gauge bosons

## Signature:

KK excitations of the gauge bosons (Z<sup>(k)</sup>, W<sup>(k)</sup>) appear as resonances with masses :

$M_k = \sqrt{(M_0^2 + k^2/R^2)}$  where (k=1,2,...) & also interference effects!

- Look for l+l- decays of  $\gamma$  and Z<sup>0</sup> KK modes. Also in decays (m<sub>T</sub>) of W<sup>+/-</sup> KK modes. Or evidence of g\* via dijet  $\sigma$  or bb, tt s



## New Parameters


$R = M_C^{-1}$  : size of compact dimension

$M_C$  : compactification scale

$M_0$  : mass of the SM gauge boson

G. Azuelos, G. Polesello  
EPJ Direct 10.1140 (2004)







# TeV<sup>-1</sup> Sized Extra Dimensions



Look for resonances/deviations in the  $ll$  spectrum



## 1) Search for the resonance peak



$\mathcal{L}=30/80 \text{ fb}^{-1}$  CMS will be able to detect a peak in the  $e^+e^-$  invar.  $M_{ll}$  if  $M_C < 5.5/6 \text{ TeV}$ .

ATLAS:  $\mathcal{L}=100 \text{ fb}^{-1}$   
 $M_C (R^{-1}) < 5.8 \text{ TeV}$   
 $(ee+\mu\mu)$


## 2) Search for interference in a mass window



## 3) Fit to kinematics of signal

With  $300 \text{ fb}^{-1}$  can reach  $13.5 \text{ TeV}$  ( $ee+\mu\mu$ )

$ee+\mu\mu$ : ATLAS  $5\sigma$  reach is  $\sim 8 \text{ TeV}$  for  $\mathcal{L}=100 \text{ fb}^{-1}$  and  $\sim 10.5 \text{ TeV}$  for  $300 \text{ fb}^{-1}$



# Distinguishing $Z^{(1)}/\gamma^{(1)}$ from $Z'$ , RS G




Distinguish


- spin-1  $Z^{(1)}$  from spin-2 G: **angular distribution** of decay products
- spin-1  $Z^{(1)}$  from spin-1  $Z'$  with SM-like couplings: **forward-backward asymmetry**  
 due to contributions of the higher lying states, interference terms and additional  $\sqrt{2}$  factor in its coupling to SM fermions.

The  $Z^{(1)}$  can be discriminated for masses up to about 5 TeV with  $L=300\text{fb}^{-1}$ .

4 TeV  $Z^{(1)}/\gamma^{(1)}$  or  $Z'$  or RS Graviton?




Forward-backward asymmetries:



100 pb<sup>-1</sup>

G. Azuelos, G. Polesello  
 EPJ Direct 10.1140 (2004)



# TeV<sup>-1</sup> ED Discovery Limits



## W<sub>KK</sub> decays

Search for deviations in lepton-neutrino transverse invariant mass ( $m_T^{l\nu}$ ) spectra


1) Search for peak:

$L=100 \text{ fb}^{-1}$  detect a peak if compactification scale ( $M_C = R^{-1}$ )  $< 6 \text{ TeV}$   
Sum over 2 lepton flavours

2) Studying distribution below the peak: in  $m_T^{e\nu}$  spectra


$L=100 \text{ fb}^{-1}$  a limit of  $M_C > 11.7 \text{ TeV}$

-ve interference sizable even for  $M_C$  above the ones accessible to a direct detection of the mass peak.



If a peak is detected, a measurement of the couplings of the boson to the leptons and quarks can be performed for  $M_C$  up to  $\sim 5 \text{ TeV}$ .

G. Polesello, M. Patra  
EPJ Direct, ATLAS 2003-023  
G. Polesello, M. Patra  
EPJ Direct C 32 Sup.2 (2004) pp.55-67




# Universal Extra Dimensions

## Standard/Minimal UED



- ❑ All particles can travel into the bulk, so each SM particle has an infinite tower of KK partners
- ❑ Spin of the KK particles is the same as their SM partners
- ❑ In minimal UED: 1 ED compactified in an orbifold ( $S^1/Z_2$ ) of size  $R$ 
  - ❑ KK parity conservation  $\rightarrow$  the lightest massive KK particle (LKP) is stable (dark matter candidate).
  - ❑ Level one KK states must be pair produced
- ❑ Mass degeneration except if radiative corrections included

The model parameters: compactification radius  $R$ , cut-off scale  $\Lambda$ ,  $m_h$




### Thick/Fat brane

- ❑ SM brane is endowed with a finite thickness in the ED
- ❑ Gravity-matter interactions break KK number conservation:
  - 1st level KK states decay to  $G+SM$ .
  - If radiative corrections  $\rightarrow$  mass degeneracy is broken and  $\gamma$  and leptons are produced.




Standard UED



# UED Discovery Limit

- Geo accep
- L1,HLT
- 2 OSSF
- 4 ISO
- b-tag veto
- $p_T^l <$
- $E_T^{miss}$
- Z veto



$$pp \rightarrow g_1 g_1 \rightarrow 4l + 4q + 2LKP \rightarrow 4l + 4 jets + P_T$$


$$pp \rightarrow g_1 Q_1 \rightarrow 4l + 3q + 2LKP \rightarrow 4l + 3 jets + P_T$$


$$pp \rightarrow Q_1 Q_1 \rightarrow 4l + 2q + 2LKP \rightarrow 4l + 2 jets + P_T$$

**Signature:** 4 low- $p_T$  isolated leptons (2 pairs of opposite sign, same flavour leptons) + n jets + missing  $E_T$  (from 2 undetected  $\gamma_1$ )

**Irreducible Bckg:**  $t\bar{t}$  + n jets (n = 0,1,2), 4 b-quarks, ZZ, Zbbar

Studied for low lum run  $\sim 2 \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$






# UED Discovery Limit



Thick brane in UED with  $\text{TeV}^{-1}$  size

$$pp \rightarrow g_1 g_1 / q_1 g_1 / q_1 q_1 \rightarrow 2 \text{ jets} + E_T$$




Signature:


2 back-to back jets + missing  $E_T$  ( $>775$  GeV)

Irreducible Bckg:  $Z(\rightarrow \nu\nu)$   $jj$ ,  $W(\rightarrow lv)$   $jj$

Significance vs Mass of 1<sup>st</sup> KK excitation



**5 $\sigma$  discovery possible at ATLAS with  $100 \text{ fb}^{-1}$  if first KK excitation mass  $< 2.7 \text{ TeV}$**




# Summary & Outlook




- ATLAS & CMS have significant discovery potential related to fundamental symmetries, Electroweak symmetry breaking, quark-lepton family structure and extra dimensions.
  - Heavy gauge bosons up to  $\sim 5\text{-}6$  TeV
  - Little Higgs T quark up to  $\sim 2$  TeV
  - Vector boson resonances; Technihadron  $\rho_{\text{TC}}$  mass up to  $\sim 600$  GeV
  - Doubly-charged Higgs up to  $\sim 2$  TeV
  - Heavy neutrino up to  $\sim 2.5$  TeV,
  - RS Model ED up to  $\sim 4$  TeV,  $Z^{(\text{KK})}/\gamma^{(\text{KK})}$  up to  $\sim 13$  TeV
  - *Many more topics not covered*
- ATLAS & CMS increasing focus on first year of data taking
  - Understand/optimize detector performance (calibration, alignment, ...)
  - Understand/measure Standard Model processes (bkg sources)

**Once these are achieved ATLAS could potentially have new physics results within months!**
- **Eager to start exploration of TeV scale!**

# *Backup Slides*









# EWSB: Resonant Vector Boson Scattering



- SM cross section for  $W_{\text{long}} W_{\text{long}}$  scattering diverges at high energy if there is no Higgs → new physics via diboson resonances?
- Chiral Lagrangian Model
  - low-energy effective description of electroweak interactions  
→ yields interaction terms describing VB scattering with arb. coeffs.
  - respects chiral symmetry via  $SU(2)_L \otimes SU(2)_R$
  - choose parameters such that new resonance  $M = 1.15 \text{ TeV}$
- Study  $W Z$  scattering (cleaner than  $W W$  + to reconstruct mass):
  - $qq \rightarrow qqWZ \rightarrow qq \ell\nu \ell\ell$  ( $\sigma \times \text{BR} = 1.3 \text{ fb}$ )
  - $qq \rightarrow qqWZ \rightarrow qq jj \ell\ell$  ( $\sigma \times \text{BR} = 4.1 \text{ fb}$ )
  - $qq \rightarrow qqWZ \rightarrow qq \ell\nu jj$  ( $\sigma \times \text{BR} = 14 \text{ fb}$ )







# Resonant Vector Boson Scattering



- Selection:** 2 forward jets + central jets and/or leptons  
 + missing  $E_T$  (for  $W \rightarrow l \nu$ )  
 Require no additional central jet & b-jet veto (for jet modes)
- Bkg:** gluon and g/Z exchange with W and Z radiation  
 also  $t \bar{t}$  & W+4 jets (need more stats)
- Promising sensitivity for jet modes at  $100 \text{ fb}^{-1}$   
 (need  $300 \text{ fb}^{-1}$  for  $WZ \rightarrow l \nu ll$ )  
 → study is ongoing



ATL-COM-PHYS-2006-041



$WZ \rightarrow jj ll$



## UED Discovery Limit




$$pp \rightarrow g_1 g_1 \rightarrow 4l + 4q + 2LKP \rightarrow 4l + 4 \text{ jets} + P_T$$

$$pp \rightarrow g_1 Q_1 \rightarrow 4l + 3q + 2LKP \rightarrow 4l + 3 \text{ jets} + P_T$$

$$pp \rightarrow Q_1 Q_1 \rightarrow 4l + 2q + 2LKP \rightarrow 4l + 2 \text{ jets} + P_T$$


- ❑ 4 leptons in the final state + missing  $P_T$
  - ❑ Irreducible Bckg:  $t\bar{t}$  + n jets (n = 0,1,2), 4 b-quarks, ZZ,  $Z\bar{b}$
  - ❑ To improve bkgd rejection over signal: apply b-tagging and Z-tagging vetoes
  - ❑ CompHEP for signal and CompHEP, PYTHIA, Alpgen for bckg. with CTEQ5L
  - ❑ Full simulation/reco + L1 + HLT(rigger) cuts
  - ❑ Theoretical and experimental uncert.
- Studied for low lum run  $\sim 2 \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$







# LHC: Black Hole Signatures

Dimopoulos and Landsberg PRL87 (2001) 161602



- In large ED (ADD) scenario, when impact parameter smaller than Schwarzschild radius Black Hole produced with potentially large x-sec ( $\sim 100$  pb).
- Decays democratically through Black Body radiation of SM states – Boltzmann energy distribution.



- Discovery potential (preliminary)
  - $M_p < \sim 4$  TeV  $\rightarrow < \sim 1$  day
  - $M_p < \sim 6$  TeV  $\rightarrow < \sim 1$  year
- Studies continue ...



## TeV<sup>-1</sup> ED g\* Discovery Limits



This is more challenging than Z/W which have leptonic decay modes

Detect KK gluon excitations (g\*) by reconstructing their hadronic decays (no leptonic decays).

Detect g\* by (1) deviation in dijet  $\sigma$

(2) analysing its decays into heavy quarks

### Gluon excitation decays

$$q\bar{q} \rightarrow g^* \rightarrow b\bar{b}, q\bar{q} \rightarrow g^* \rightarrow t\bar{t}$$



- bbar or tbar jets
- For tbar one t is forced to decay leptonically
- Bckg: SM continuum bbar, tbar, 2 jets, W +jets
- PYTHIA
- Fast simulation/reco


With 300 fb<sup>-1</sup> Significance of 5 achieved for:

bbar channel: R<sup>-1</sup> = 2.7 TeV



tbar channel: R<sup>-1</sup> = 3.3 TeV

### Reconstructed mass peaks






# Excited Neutrino Discovery Potential



With  $300 \text{ fb}^{-1}$




# CMS SUSY



$\text{GeV}/c^2$

Point	$m_0$	$m_{1/2}$	$\tan \beta$	$\text{sgn}(\mu)$	$A_0$
LM1	60	250	10	+	0
LM2	185	350	35	+	0
LM3	330	240	20	+	0
LM4	210	285	10	+	0
LM5	230	360	10	+	0
LM6	85	400	10	+	0
LM7	3000	230	10	+	0
LM8	500	300	10	+	-300
LM9	1450	175	50	+	0
LM10	3000	500	10	+	0
HM1	180	850	10	+	0
HM2	350	800	35	+	0
HM3	700	800	10	+	0
HM4	1350	600	10	+	0



# A Toroidal LHC Apparatus (ATLAS) DETECTOR



EM Calorimeters,  $\sigma/E \approx 10\%/\sqrt{E(\text{GeV})} \oplus 0.7\%$   
 excellent electron/photon identification  
 Good  $E$  resolution (e.g.,  $H \rightarrow \gamma\gamma$ )

Precision Muon Spectrometer,  
 $\sigma/p_T \approx 10\%$  at 1 TeV/c  
 Fast response for trigger  
 Good  $p$  resolution  
 (e.g.,  $A/Z' \rightarrow \mu\mu$ ,  $H \rightarrow 4\mu$ )


**Full coverage for  $|\eta| < 2.5$**

Detector characteristics	
Width:	44m
Diameter:	22m
Weight:	7000t

CERN AC - ATLAS V1997

Hadron Calorimeters,  
 $\sigma/E \approx 50\% / \sqrt{E(\text{GeV})} \oplus 3\%$   
 Good jet and  $E_T$  miss performance  
 (e.g.,  $H \rightarrow \tau\tau$ )


Inner Detector:  
 Si Pixel and strips (SCT) &  
 Transition radiation tracker (TRT)  
 $\sigma/p_T \approx 5 \times 10^{-4} p_T \oplus 0.001$   
 Good impact parameter res.  
 $\sigma(d_0) = 15\mu\text{m}@20\text{GeV}$  (e.g.  $H \rightarrow b\bar{b}$ )



Magnets: solenoid (Inner Detector) 2T, air-core toroids (Muon Spectrometer) ~0.5T



# Compact Muon Solenoid (CMS) DETECTOR



## ATLAS Inclusive Trigger Selection Signatures


- To select an extremely broad spectrum of “expected” and “unexpected” Physics signals (hopefully!).
- The selection of Physics signals requires the identification of **objects** that can be **distinguished** from the high particle density environment.

Object	Examples of physics coverage	Nomenclature
Electrons	Higgs (SM, MSSM), new gauge bosons, extra dimensions, SUSY, W/Z, top	e25i, 2e15i
Photons	Higgs (SM, MSSM), extra dimensions, SUSY	$\gamma$ 60i, 2 $\gamma$ 20i
Muons	Higgs (SM, MSSM), new gauge bosons, extra dimensions, SUSY, W/Z, top	$\mu$ 20i, 2 $\mu$ 10
Jets	SUSY, compositeness, resonances	j360, 3j150, 4j100
Jet+missing $E_T$	SUSY, leptoquarks, “large” extra dimensions	j60 + xE60
Tau+missing $E_T$	Extended Higgs models (e.g. MSSM), SUSY	$\tau$ 30 + xE40


**also inclusive missing $E_T$ , Sum $E_T$ , Sum $E_T$ \_jet**

**& many prescaled and mixed triggers**

The list must be non-biasing, flexible, include some redundancy, extendable, to account for the “unexpected”.





# Experimental Signature for RS Model



**Signature:**  
Narrow, high-mass resonance states in dilepton/dijet/diboson channels


$$q\bar{q}, gg \rightarrow G_{KK} \rightarrow e^+e^-, \mu^+\mu^-, \gamma\gamma, jet + jet$$



## Model parameters:

- Gravity Scale:  $\Lambda_\pi = \overline{M}_{pl} e^{-kR_c\pi}$   
1<sup>st</sup> graviton excitation mass:  $m_1 \rightarrow$  **Resonance position**
  - $\Lambda_\pi = m_1 \overline{M}_{pl} / kx_1$ , &  $m_n = kx_n e^{krc\pi} (J_1(x_n) = 0)$
  - Coupling constant:  $c = k/M_{pl}$   
 $\Gamma_1 = \rho m_1 x_1^2 (k/M_{pl})^2 \rightarrow$  **width**
- $k =$  curvature,  $R =$  compactification radius


Davoudiasl, Hewett, Rizzo  
hep-ph0006041



# CMS SUSY



Best reach obtained with the most inclusive channels:  
jets+MET & muons+jet+MET



The range of gluino and squark masses up to about 1.5 TeV can be probed with  $L=1\text{fb}^{-1}$  and up to about 2 TeV with  $10\text{fb}^{-1}$


Simultaneous observation of a signal in various topologies would help determine the underlying physics. E.g. triangular dilepton mass distribution.

If discovered plan would be to do more exclusive analyses to reconstruct the sparticle masses and measure cross-sections of sub-processes and their ratios


# Heavy Gauge Bosons: $Z'$




■ CMS  $Z'$  studies (TDR): integrated luminosity needed for  $5\sigma$  signal



$Z'$   $5\sigma$  reach ee vs.  $\mu\mu$  channels



$Z'$   $5\sigma$  reach: impact of theory uncertainties




# The LHC Machine Schedule



<i>year</i>	<i>energy</i>	<i>luminosity</i>	<i>physics beam time</i>
<b>2007</b>	<b>450+450 GeV</b>	<b><math>5 \times 10^{30}</math></b>	<b>protons - 26 days at 30% overall efficiency → <math>0.7 \times 10^6</math> seconds</b>
<b>2008</b>	<b>7+7 TeV</b>	<b><math>0.5 \times 10^{33}</math></b>	<b>protons - starting beginning July <math>4 \times 10^6</math> seconds ions - end of run - 5 days at 50% overall efficiency → <math>0.2 \times 10^6</math> seconds</b>
<b>2009</b>	<b>7+7 TeV</b>	<b><math>1 \times 10^{33}</math></b>	<b>protons: 50% better than 2008 → <math>6 \times 10^6</math> seconds ions: 20 days of beam at 50% efficiency → <math>10^6</math> seconds</b>
<b>2010</b>	<b>7+7 TeV</b>	<b><math>1 \times 10^{34}</math></b>	<b>TDR targets: protons: → <math>10^7</math> seconds ions: → <math>2 \times 10^6</math> seconds</b>

See many other talks at this meeting for ATLAS experimental details





# ATLAS detector

- High energy electrons are detected by LAr calorimeter.
- Muons are detected by the Muon System.
- Expected electron energy resolution is:
  - $\sim 0.6\%$  for  $E=500\text{GeV}$ ,
  - $\sim 0.5\%$  for  $E=1000\text{GeV}$ .
- Muon transverse momentum ( $p_T$ ) resolution is:
  - $\sim 6\%$  for  $p_T=500\text{GeV}$ ,
  - $\sim 11\%$  for  $p_T=1000\text{GeV}$ .

Electron energy resolution

$$\frac{\sigma(E)}{E} = \frac{9.5\%}{\sqrt{E}} \oplus 0.45\%$$








# High $p_T$ leptons from $Z'$ decay



- The leptons  $p_T$  distribution from  $Z'$  decay has a Jacobian peak.
- At high  $p_T$  the muon momentum resolution degrades.
- For the muon  $p_T$  resolution, calibration and alignment are critical.







# Forward-backward asymmetry




- As a probe of the underlying model, one can measure the forward-backward asymmetry.
- The differential cross section of  $Z'$  depends on  $\cos\theta^*$ .
- And if  $Z'$  has spin 1, the differential cross section is given by:

$$A_{FB}(M_{l\nu}) \frac{d\sigma}{d\cos\theta^*} \propto \frac{3}{8}(1 + \cos^2\theta^*) + A_{FB}\cos\theta^* \quad \text{!}$$

$\theta^*$  is angle between  $l$  and quark in the CMS of the colliding partons.


- This quantity  $A_{FB}(M_{ll}) = \frac{N_+ - N_-}{N_+ + N_-}$  is independent.  $N_+$ : number of events with the lepton in the forward  
 $N_-$ : number of events with the lepton in the backward
- One can discriminate between the underlying models by measuring  $A_{FB}(M_{ll})$ .

Ref: ATL-PHYS-PUB-2005-010



# $A_{FB}(M_{ll})$ measurement(1)


- $Z' \rightarrow e^+e^-$  :
  - high discriminating power of the asymmetry.



- Correction:
  - Taking into account mis-estimation of quark direction.
  - Fractions of the mis-estimation of quark direction is parameterized by simulation.



Model	$\int \mathcal{L}(fb^{-1})$	Generation	Observed	Corrected
1.5 TeV				
<i>SSM</i>	100	$+0.088 \pm 0.013$	$+0.060 \pm 0.022$	$+0.108 \pm 0.027$
$\chi$	100	$-0.386 \pm 0.013$	$-0.144 \pm 0.025$	$-0.361 \pm 0.030$
$\eta$	100	$-0.112 \pm 0.019$	$-0.067 \pm 0.032$	$-0.204 \pm 0.039$
$\eta$	300	$-0.090 \pm 0.011$	$-0.050 \pm 0.018$	$-0.120 \pm 0.022$
$\psi$	100	$+0.008 \pm 0.020$	$-0.056 \pm 0.033$	$-0.079 \pm 0.042$
$\psi$	300	$+0.010 \pm 0.011$	$-0.019 \pm 0.019$	$-0.011 \pm 0.024$
<i>LR</i>	100	$+0.177 \pm 0.016$	$+0.100 \pm 0.026$	$+0.186 \pm 0.032$
4 TeV				
<i>SSM</i>	500	$+0.138 \pm 0.099$	$+0.006 \pm 0.183$	$+0.265 \pm 0.260$
<i>KK</i>	500	$+0.491 \pm 0.028$	$+0.189 \pm 0.057$	$+0.457 \pm 0.073$

Ref: ATL-PHYS-PUB-2005-010




# $A_{FB}(M_{\mu\mu})$ measurement(2)

- $Z' \rightarrow \mu^+ \mu^-$ 
  - With  $200\text{fb}^{-1}$ , the ATLAS can distinguish the underlying theories with accuracy better than 3% using the asymmetry for  $M(Z')$  less than 2TeV.
  - At higher masses, we need much more luminosity.




Ref: ATLAS Internal Note Muon-NO-161 23 May 1997



# Forward and backward


Forward

$$\cos \theta^* > 0$$




Backward

$$\cos \theta^* < 0$$



When  $\cos \theta^*$  is positive, we call forward, and when  $\cos \theta^*$  is negative we call backward. The quark direction is not directly accessible in the data. Therefore the  $Z'$  momentum defines the quark direction, because of the quark generally being at a higher momentum than the antiquark



# Specific models

- A popular model is:

- Effective  $SU(2) \times U(1)_Y \times U(1)'$ 
  - There are two additional neutral gauge bosons.
- The new gauge boson uniquely determined by:

$$Z' = \cos \theta Z'_\psi - \sin \theta Z'_\chi$$

*θ is a new mixing angle.*


*2 independent U(1) bosons.*

- There are 3 special cases:
  - $Z'_\psi$  model:  $\theta=0, E_6 \rightarrow SO(10) \times U(1)_\psi$
  - $Z'_\chi$  model:  $\theta=-\pi/2, E_6 \rightarrow SO(10) \times U(1)_\psi \rightarrow SU(5) \times U(1)_\chi \times U(1)_\psi$
  - $Z'_\eta$  model:  $\theta = \arctan(-\sqrt{5/3}) + \pi/2,$   
 $E_6 \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_\eta = SM \times U(1)_\eta$   
 ( $E_6$  breaks directly down to a rank 5 model.)

- Other popular models:

- The Left-Right model from the breaking of the  $SO(10)$  group,
- The Kaluza-Klein model (Extra Dimension).
- etc...

Ref: ATL-PHYS-PUB-2005-010




# ATLAS and CMS Experiments




## Large general-purpose particle physics detectors

### A Large Toroidal LHC Apparatus



<b>Total weight</b>	<b>7000 t</b>
<b>Overall diameter</b>	<b>25 m</b>
<b>Barrel toroid length</b>	<b>26 m</b>
<b>End-cap end-wall chamber span</b>	<b>46 m</b>
<b>Magnetic field</b>	<b>2 Tesla</b>

### Compact Muon Solenoid



<b>Total weight</b>	<b>12 500 t</b>
<b>Overall diameter</b>	<b>15.00 m</b>
<b>Overall length</b>	<b>21.6 m</b>
<b>Magnetic field</b>	<b>4 Tesla</b>

Detector subsystems are designed to measure:  
 energy and momentum of  $\gamma$ ,  $e$ ,  $\mu$ , jets, missing  $E_T$  up to a few TeV




## ADD Parameters: jet+G Emission




To characterise the model need to measure  $M_D$  and  $\delta$

Measuring  $\sigma(pp \rightarrow \text{jet} + G^{KK})$  gives ambiguous results



Use variation of  $\sigma$  on  $\sqrt{s}$   
 $\sigma$  at different  $\sqrt{s}$  almost independent of  $M_D$ , varies with  $\delta$

Run at two different  $\sqrt{s}$   
 e.g. 10 TeV and 14 TeV, need  $50 \text{ fb}^{-1}$



Rates at 14 TeV of  $\delta=2$   $M_D=6 \text{ TeV}$  very similar to  $\delta=3$   $M_D=5 \text{ TeV}$  whereas Rates at 10 TeV of ( $\delta=2$   $M_D=6 \text{ TeV}$ ) and ( $\delta=3$   $M_D=5 \text{ TeV}$ ) differ by  $\sim$  factor of 2




## RS1 Model Parameters



A resonance could be seen in many other channels:  $\mu\mu$ ,  $\gamma\gamma$ ,  $jj$ ,  $b\bar{b}$ ,  $t\bar{t}$ ,  $WW$ ,  $ZZ$ , hence allowing to **check universality of its couplings**:

Channel	Point $m_G, \Lambda_\pi$ (TeV)							
	1,10	1,20	1,30	2,10	2,20	2,30	3,10	3,20
$e^+e^-$	1.6	3.3	5.3	5.4	11.0	17.1	15.1	30.7
$\mu^+\mu^-$	1.9	4.5	8.2	6.2	15.2	28.2	15.1	32.7
$\gamma\gamma$	1.2	2.9	5.2	3.9	8.8	15.2	10.5	23.0
$WW$	11.6	44.9	-	38.2	-	-	-	-
$ZZ$	13.7	50.1	-	52.7	-	-	-	-
$jj$	19.0	77.0	-	31.0	-	-	59.0	-

Relative precision achievable (in %) for measurements of  $\sigma_B$  in each channel for fixed points in the  $M_G, \Lambda_\pi$  plane. Points with errors above 100% are not shown.




Also the **size (R) of the ED** could also be estimated from mass and cross-section measurements.

Allenach et al, hep-ph0211205

Allenach et al, JHEP 9 19 (2000), JHEP 0212 39 (2002)





# LHC Start-up Expectations



Model	Mass reach	Integrated Luminosity (fb <sup>-1</sup> )	Systematic uncertainties
<b>ADD</b> Direct G <sub>KK</sub>	M <sub>D</sub> ~ 1.5-1.0 TeV, n = 3-6	1	Theor.
<b>ADD</b> Virtual G <sub>KK</sub>	M <sub>D</sub> ~ 4.3 - 3 TeV, n = 3-6	0.1	Theor.+Exp.
	M <sub>D</sub> ~ 5 - 4 TeV, n = 3-6	1	
<b>RS1</b> di-electrons di-photons di-muons di-jets	M <sub>G1</sub> ~ 1.35- 3.3 TeV, c=0.01-0.1	10	Theor.+Exp.  (only stat. for di-jets)
	M <sub>G1</sub> ~ 1.31- 3.47 TeV, c=0.01-0.1	10	
	M <sub>G1</sub> ~ 0.8- 2.3 TeV, c=0.01-0.1	1	
	M <sub>G1</sub> ~ 0.7- 0.8 TeV, c=0.1	0.1	
<b>TeV<sup>-1</sup></b> (Z <sub>KK</sub> <sup>(1)</sup> )	M <sub>z1</sub> < 5 TeV	1	Theor.



# Spin-1/Spin-2 Discrimination





**Spin-1 States:**  $Z'$  from extended gauge models,  $Z_{KK}$

**Spin-2 States:** RS1-graviton

**Method:** unbinned likelihood ratio statistics incorporating the angles in of the decay products the Collins-Soper frame (R.Cousins et al. JHEP11 (2005) 046). The statistical technique has been applied to fully simu/reco events.

## Angular distributions

- $qq \rightarrow G \rightarrow ff: 1 - 3 \cos^2 \theta + 4 \cos^4 \theta$
- $gg \rightarrow G \rightarrow ff: 1 - \cos^4 \theta$
- $qq \rightarrow G \rightarrow VV: 1 - \cos^4 \theta$
- $gg \rightarrow G \rightarrow VV: 1 + 6 \cos^2 \theta + \cos^4 \theta$
- DY background:  $1 + \cos^2 \theta$




## Older results on spin discrimination from ATLAS

B. C. Allanach et al, JHEP 09 (2000) 019; ATL-PHYS-2000-029



## Distinguishing $Z^{(1)}$ from $Z'$ , RS G


How could a 4 TeV  $Z^{(1)}/\gamma^{(1)}$  resonance be distinguished from a 4 TeV  $Z'$  or Randall-Sundrum Model Graviton?



Select events around the peak of the resonance  $3750 \text{ GeV} < M_{ee} < 4250 \text{ GeV}$

Plot cosine of the angle of the lepton, w.r.t the beam direction, the frame of the decaying resonance.

(+ve direction was defined by the sign of reconstructed momentum in the dilepton system.)



Angular distributions are normalized to 116 events, the number predicted with a luminosity of  $100 \text{ fb}^{-1}$  for the  $Z^{(1)}/\gamma^{(1)}$  case