

Effective Field Theory Approach to Weak Boson Physics

Harrison Mebane

University of Illinois at Urbana-Champaign

May 10, 2011

The Challenge

- We believe there is new physics at the terascale
- Each BSM theory has its own set of parameters
- We would like a single, model-independent framework for parametrizing indirect effects of new physics

Effective Field Theory: An Introduction

- At sufficiently low energies, new physics can be integrated out and treated perturbatively
- The resulting Lagrangian will contain terms of dimension greater than four, suppressed by inverse powers of the energy scale of new physics

$$\mathcal{L}_{eff} = \mathcal{L}_{SM}^{(4)} + \frac{1}{\Lambda} \sum_i C_i \mathcal{O}_i^{(5)} + \frac{1}{\Lambda^2} \sum_j C_j \mathcal{O}_j^{(6)} + \dots$$

Effective Field Theory Properties

- Gauge-invariant
- Reduces to the standard model in the low-energy limit
- Renormalizable in the “modern sense”

Extending the Standard Model

- We need to extend the standard model in a way that will handle all possible low-energy effects of new physics
- A complete basis comprises 59 dimension-six operators (there is only one dimension-five operator)
- Here, we will focus on operators involving electroweak bosons

Operators Contributing to Vector Boson Pair Production

Triple-Boson

$$\mathcal{O}_W = \epsilon_{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$$

$$\mathcal{O}_{WB} = (\phi^\dagger \tau^I \phi) W_{\mu\nu}^I B^{\mu\nu}$$

$$\mathcal{O}_{\tilde{W}} = \epsilon_{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$$

$$\mathcal{O}_{\tilde{W}B} = (\phi^\dagger \tau^I \phi) \tilde{W}_{\mu\nu}^I B^{\mu\nu}$$

Fermion-Fermion-Boson

$$\mathcal{O}_{\phi q}^{(1)} = i(\phi^\dagger D_\mu \phi)(\bar{q}\gamma^\mu q)$$

$$\mathcal{O}_{\phi q}^{(3)} = i(\phi^\dagger \tau^I D_\mu \phi)(\bar{q}\gamma^\mu \tau^I q)$$

$$\mathcal{O}_{\phi u} = i(\phi^\dagger D_\mu \phi)(\bar{u}\gamma^\mu u)$$

$$\mathcal{O}_{\phi d} = i(\phi^\dagger D_\mu \phi)(\bar{d}\gamma^\mu d)$$

$$\mathcal{O}_{\phi\phi} = i(\tilde{\phi}^\dagger D_\mu \phi)(\bar{u}\gamma^\mu d)$$

Anomalous Couplings

- The anomalous couplings approach involves writing down interaction terms with no regard for gauge invariance

$$\mathcal{L} = ig_{WWV} \left(g_1^V (W_{\mu\nu}^+ W^{-\mu} - W^{+\mu} W_{\mu\nu}^-) V^\nu + \kappa_V W_\mu^+ W_\nu^- V^{\mu\nu} + \frac{\lambda_V}{m_W^2} W_\mu^{+\nu} W_\nu^{-\rho} V_\rho^\mu + \dots \right)$$

- Alternatively, we can move to momentum space and write down a vertex function

$$\Gamma^{\mu\nu\rho} = f_1^V(p^2, q^2) \{p^\nu \eta^{\mu\rho} - q^\mu \eta^{\nu\rho} + (q-p)^\rho \eta^{\mu\nu}\} + f_2^V(p^2, q^2) \{(p+q)^\nu \eta^{\mu\rho} - (p+q)^\mu \eta^{\nu\rho}\} + \dots$$

where f_1^V and f_2^V are functions of momentum and are called form factors

Unitarity

- S-matrix unitarity and partial wave analysis imply a bound on the amplitude of any 2-to-2 inelastic scattering process

$$\sum_{\lambda_3, \lambda_4} \int dPS_2 |T^{in}|^2 \leq 8\pi$$

- Higher-dimensional operators may add terms to $|T^{in}|^2$ which grow like s/Λ^2 , possibly violating unitarity at high energies

Unitarity, cont'd

- In the anomalous couplings approach, form factors are constructed so as prevent unitarity violation at arbitrarily high energies
- An effective field theory will not violate unitarity at energies less than Λ and has nothing to say about energies above Λ
- We are interested in energies for which there is data; data does not violate unitarity

Tree-Level Cross Section




Figure: Cross section versus invariant mass for the process $pp \rightarrow W^+W^-$ at the LHC. The pink curve includes the dimension-six operator O_W . $\frac{C_W}{\Lambda^2}$ is set to 1 TeV^{-2} . Generated with MadGraph.

Summary

- Effective field theory is a model-independent framework for studying indirect effects of physics beyond the standard model
- Unlike the anomalous couplings approach, it preserves $SU(3) \times SU(2) \times U(1)$ gauge symmetry and provides a framework for handling divergences
- Unitarity violation is not a concern in effective field theory, and form factors are not needed
- Soon all dimension-six operators will be incorporated into MadGraph