



Contribution ID: 112

Type: **Poster**

Investigating the ridge structure in #Delta#eta-#Delta#phi correlations at STAR

Thursday, 16 August 2012 16:00 (2 hours)

Triggered di-hadron correlation studies using Au+Au collisions at $\sqrt{S_{NN}} = 200\text{GeV}$ in STAR revealed a novel “ridge-like” structure in two dimensions ($\Delta\eta$, $\Delta\phi$) [1] for high p_T particles. Similar structure was also present in an inclusive un-triggered di-hadron correlation analysis [2]. We study the $\langle p_T \rangle$ evolution of un-triggered analysis by increasing the lower p_T cut of both charged particles. As smooth evolution of data is observed for $p_T \geq 2.7\text{GeV}/c$. We further quantify the correlation structure evolution by fitting a model function. The model function emphasizes the $\Delta\eta$ dependence for $p_T < 2.7\text{GeV}/c$ and the remainder is modeled via a asymmetric 2d Gaussian. Extracted harmonic parameters are compared to model predictions at $\sqrt{S_{NN}} = 200\text{GeV}$ and charged dependent studies.

- [1] B. I. Abelev et al. (STAR Collaboration), Phys. Rev. C 80, 064912 (2009)
- [2] M Daugherty et al. (STAR Collaboration), J. Phys. G 35, 104090 (2008)
- [3] B. Alver et al., Phys. Rev. C 81, 054905 (2010)
- [4] C. Gombeaud et al., Phys. Rev. C 81, 014901 (2010)

Primary author: Mr DE SILVA, L. Chanaka (University of Houston)

Presenter: Mr DE SILVA, L. Chanaka (University of Houston)

Session Classification: Poster Session Reception

Track Classification: Correlations and fluctuations