


Multi-parton interactions & the underlying event

Tim Namssoo - *DESY*

HERA LHC Workshop, 31st October 2007, DESY, Hamburg.

- *Introduction*
- *Motivation*
- *HERA: Underlying event in γp*
- *HERA: Underlying event in DIS*
- *Tevatron: Underlying event in $p\bar{p}$*
- *LHC?: Underlying event in pp*
- *Summary*




Introduction


- What is the underlying event? (*a working definition:*)
 - **all energy flow not associated with the primary process**
- What is the primary process?!
 - the ideal parton-parton collision as if the beams only contained free partons
 - (beyond PDFs) is insensitive to the incoming particles and beam remnants
 - includes all coherent radiation (to all orders) associated with that interaction
- What else could affect or contribute to the observable energy flow - i.e. constitutes the underlying event?
 - remnant-remnant interactions - i.e. multiple parton-parton interactions
 - multiple-scattering as a primary parton re-scatters off the remnants - i.e. one parton interacts with multiple partons from incoming hadron
 - such processes will be referred to as multi-parton interactions (MPIs)
- Although there are possibly other effects, I shall talk exclusively about MPIs

Introduction - MPIs

- The qualitative argument is, if one (or both) of the incoming particles has a hadron-like structure, multiple-scattering (and remnant-remnant interactions) can occur.
- There are more general arguments based on optical theorem & AGK cutting rules.
- The study of diffraction at HERA has provided evidence of pomeron exchange.
- Assuming one may describe the pomeron using a gluon ladder, the “squared” diagrams below should contribute to the inclusive cross section.
- The AGK cutting rules give a handle on the likelihood of various final states.



a) *diffractive interaction*




b) *multi-parton interaction*


- Therefore, MPIs really should be present at HERA at low- Q^2 and/or x

Introduction - MPIs

- MPIs may potentially range from being very soft upto hard (i.e. jet forming)



a) *soft remnant-remnant*



b) *hard remnant-remnant*


- thus possible MPI signatures (softest \Rightarrow hardest) are a low- E_T pedestal, increased production of (incoherent) mini-jets or an excess of (pair-wise) 4-jet events.
- experimentally, it's difficult to differentiate MPIs from HO pQCD effects

Motivation




- MPIs can interfere with many types of physics analysis so must be understood:
 - they reduce rapidity gap survival probability
 - they affect isolation criteria (e.g. for muons)
 - they lead to larger charged/particle multiplicities
 - affect jet profiles/pedestals, effectively increasing jet energy scale
 - potentially increase jet rates and (if jet forming) affect jet angular correlations
- And MPIs will be far more prevalent at the LHC
- to find (most) new physics must understand QCD background, including:
 - the primary interaction...
 - ...plus the secondary interactions...
 - ...from the multiple particle interactions per bunch crossing!
- MPIs affect LHC triggering strategies and what LHC analyses can be done
- Potentially though, MPIs may lead to a greater understanding of p - e.g.:
 - multi-parton correlated SFs from 4-jet (hard MPI) data?
 - low- x , saturation - are MPIs an inescapable consequence?

Underlying event in γp - multi-jets (ZEUS)

- Here we study: $\gamma p \rightarrow 3+$ or $4+$ jets ($E_T^{\text{jet}} > 6 \text{ GeV}$ & $|\eta^{\text{jet}}| < 2.4$)
- γ may act like a point-like (direct) or composite object (resolved)
- remnant-remnant interactions only present in resolved process
- Multi-jets generated by QCD processes
- ...and hard-MPIs? Note: soft underlying event changes jet energy scale and so, given some E_T^{jet} criteria, affects jet rates.
- Variables looked at:
 - M_{nj} : invariant mass of the n -jet system.
 - x_γ^{obs} : which approximates x_γ , the fraction of γ 's momentum transferred to the hard interaction (i.e. the jets). At LO, $x_\gamma = 1$ (direct) & $x_\gamma < 1$ (resolved).
- events studied in two M_{nj} regions: ($25 \leq M_{nj} < 50 \text{ GeV}$) & ($M_{nj} \geq 50 \text{ GeV}$)
- data compared to MCs with and without simulated MPIs and LO pQCD




Underlying event in γp - multi-jets (ZEUS)



- MC without MPIs fails to describe low M_{nj} regions
- adding MPIs helps description of M_{nj}
- highest order pQCD in γp only LO for 3-jet process
- shown here corrected for hadronisation and MPI effects
- largely describes M_{3j} data but theo. uncertainty large
- description greatly improved by MPI corrs.


Underlying event in γp - multi-jets (ZEUS)




- all MC models describe high mass data reasonably well.
- MCs without MPIs don't describe low x_γ^{obs} region at low mass.
- the discrepancy between the MC without MPIs and the data is larger for 4-jets.
- introducing MPIs into the MCs improves the description.
- note: predicted influence of MPIs very sensitive to tunable parameters in models.
- low mass 4-jet data some of the most MPI sensitive ZEUS data. **However...**
- ...always issue: really MPIs or HO effects not modelled by parton-showers?

Underlying event in DIS - mini-jets (H1)



- resolved processes suppressed by virtuality, Q^2 .
- are we even sensitive to MPIs/underlying event in DIS?
- Strategy:
 - select DIS events ($5 < Q^2 < 100 \text{ GeV}^2$)
 - define + select hardest jet in HCM ($P_T^{\text{jet}} > 5 \text{ GeV}$)
 - define 4 azimuthal regions
 - \sum of particle E_T defines low/high activity regions
 - measure average mini-jet multiplicity, $\langle N_{\text{minijet}} \rangle$.
 - where mini-jets have $P_T^{\text{jet}} > 3 \text{ GeV}$



- transverse regions - sensitive to incoherent energy flow.
- can further reduce coherent radiation by requiring back-to-back subleading jet. Strategy:
 - select dijet events ($P_T^{\text{jet}} > 5 \text{ GeV}$)
 - with subleading jet in "away region"
 - repeat procedure...




Underlying event in DIS - mini-jets (inclusive)



- $\langle N_{\text{minijet}} \rangle (P_T^{\text{jet}1})$ in the 4 regions. Shown for high $\eta^{\text{jet}1}$ region in 3 Q^2 bins.
- expect larger resolved contribution in high $\eta^{\text{jet}1}$ (forward) region.
- all MC models describe the “towards” and “away” regions reasonably well.
- MPIs improve description of “low” and “high” regions at low Q^2 but not at mid Q^2




Underlying event in DIS - mini-jets (dijets)



- $\langle N_{\text{minijet}} \rangle (P_T^{\text{jet}1})$ in the 4 regions in two x_γ regions.
- “towards” and “away” regions again largely described by all MC models
- more activity in “low” and “high” regions at low x_γ (resolved enriched)
- low x_γ description generally improved by the inclusion of MPIs

Underlying event in $p\bar{p}$ - transverse P_T



- Tevatron underlying event most relevant for LHC
- analysis of “transverse” regions
- plot hadronic P_T sums compared to MC models
- HERWIG (no MPIs) below the low-PT(jet#1) data
- best description by PYTHIA with MPIs (“Tune A”)



- R. Field [CDF Collab.], AIP Conf. Proc. 828 (2006) 163

Underlying event at the LHC?

- What will the underlying event be like at the LHC? Can we say anything presently?



- Clearly, LHC extrapolations based on tunes to current data disagree
- certainly first LHC data will provide an interesting test for the current models
- but beyond just being a background for physics it will be interesting if MPI events can be used constructively to gain further insight into e.g. proton structure.

Summary

- the topic of MPIs is presently very relevant. From practical considerations:
 - they interfere with triggering strategies & what physics analyses can be done

- at HERA, remnant-remnant interactions are possible in resolved γ processes.
- resolved processes suppressed with increasing Q^2 and x_γ .
- multi-jet γp (low- Q^2) data suggestive of large MPI contribution at low M_{nj} & low x_γ .
- furthermore, influence of MPIs predicted to grow with jet multiplicity.
- HERA DIS mini-jet data also suggestive of MPIs at Q^2 upto $\mathcal{O}(20)$ GeV², however this is beyond where MC predicts MPIs can have influence.
- however, always question whether MPIs or HO effects/soft physics?

- at the Tevatron, the picture is the same.
- particle P_T sums are in excess of MC prediction without MPIs
- description can be remedied by the inclusion of MPIs

- But as for the LHC, extrapolations to the relevant energies have large uncertainties
- LHC data will provide an interesting test of the models
- but beyond just being a background for physics, it will be interesting if MPI events can be used constructively to gain further physical insights