



# Experimental particle. physics

**esipap**...  
European School of Instrumentation  
in Particle & Astroparticle Physics



# Order of magnitudes



Optical microscope resolution

$$\Delta r \sim \frac{1}{\sin \theta}$$

with  $\theta$  = angular aperture of the light beam

De Broglie wave length

$$\lambda = \frac{h}{p} \quad \Delta r \sim \frac{h}{p}$$

with  $p$  = transferred momentum


# HEP, SI and “natural” units

Quantity	HEP units	SI units
length	1 fm	$10^{-15}$ m
charge	e	$1.602 \cdot 10^{-19}$ C
energy	1 GeV	$1.602 \times 10^{-10}$ J
mass	1 GeV/c <sup>2</sup>	$1.78 \times 10^{-27}$ kg
$\hbar = h/2\pi$	$6.588 \times 10^{-25}$ GeV s	$1.055 \times 10^{-34}$ Js
c	$2.988 \times 10^{23}$ fm/s	$2.988 \times 10^8$ m/s
$\hbar c$	197 MeV fm	...

## “natural” units ( $\hbar = c = 1$ )

mass	1 GeV
length	1 GeV <sup>-1</sup> = 0.1973 fm
time	1 GeV <sup>-1</sup> = $6.59 \times 10^{-25}$ s

# How much energy to probe these distances?



$$\lambda = \frac{h}{p} = \frac{2\pi\hbar c}{pc} = \frac{2\pi \times 197 \text{ MeV fm}}{pc}$$


What?	Dimension [m]	$p$ [GeV/c]
-------	---------------	-------------

Atom	$10^{-10}$	0.00001
------	------------	---------

Nucleus	$10^{-14}$	0.
---------	------------	----

Nucleon	$10^{-15}$	1
---------	------------	---


Quark	$10^{-18}$	100
-------	------------	-----




# What do we want to measure?

1968: SLAC <b><math>u</math></b> up quark	1974: Brookhaven & SLAC <b><math>c</math></b> charm quark	1995: Fermilab <b><math>t</math></b> top quark	1979: DESY <b><math>g</math></b> gluon
1968: SLAC <b><math>d</math></b> down quark	1947: Manchester University <b><math>s</math></b> strange quark	1977: Fermilab <b><math>b</math></b> bottom quark	1923: Washington University* <b><math>\gamma</math></b> photon
1956: Savannah River Plant <b><math>\nu_e</math></b> electron neutrino	1962: Brookhaven <b><math>\nu_\mu</math></b> muon neutrino	2000: Fermilab <b><math>\nu_\tau</math></b> tau neutrino	1983: CERN <b><math>W</math></b> $W$ boson
1897: Cavendish Laboratory <b><math>e</math></b> electron	1937: Caltech and Harvard <b><math>\mu</math></b> muon	1976: SLAC <b><math>\tau</math></b> tau	1983: CERN <b><math>Z</math></b> $Z$ boson
			2012: CERN <b><math>H</math></b> Higgs boson


# Baryons and Mesons



Meson octet



Baryon decuplet



Baryon octet

# Measuring particles

- Particles are characterized by
  - ✓ **Mass** [Unit: eV/c<sup>2</sup> or eV]
  - ✓ **Charge** [Unit: e]
  - ✓ **Energy** [Unit: eV]
  - ✓ **Momentum** [Unit: eV/c or eV]
  - ✓ (+ spin, lifetime, ...)

Particle identification via measurement of:

e.g. (E, p, Q) or (p, β, Q)  
(p, m, Q) ...

- ... and move at **relativistic speed**

$$\beta = \frac{v}{c} \quad \gamma = \frac{1}{\sqrt{1 - \beta^2}}$$

$$l = \frac{l_0}{\gamma} \quad \text{length contraction}$$

$$t = t_0 \gamma \quad \text{time dilatation}$$

$$E^2 = \vec{p}^2 c^2 + m^2 c^4$$

$$E = m \gamma c^2 = m c^2 + E_{\text{kin}}$$

$$\vec{\beta} = \frac{\vec{p}c}{E} \quad \vec{p} = m \gamma \vec{\beta} c$$

# Relativistic kinematics in a nutshell

$$E^2 = \vec{p}^2 + m^2$$

$$l = \frac{l_0}{\gamma}$$

$$E = m\gamma$$

$$t = t_0\gamma$$

$$\vec{p} = m\gamma\vec{\beta}$$

$$\vec{\beta} = \frac{\vec{p}}{E}$$




# Center of mass energy

- In the **center of mass frame** the total momentum is 0
- In **laboratory frame** center of mass energy can be computed as:

$$E_{\text{cm}} = \sqrt{s} = \sqrt{\left(\sum E_i\right)^2 - \left(\sum \vec{p}_i\right)^2}$$

# 2-bodies decay




$$E_1 = \frac{M^2 - m_2^2 + m_1^2}{2M},$$

$$|\mathbf{p}_1| = |\mathbf{p}_2|$$


$$= \frac{[(M^2 - (m_1 + m_2)^2)(M^2 - (m_1 - m_2)^2)]^{1/2}}{2M}$$

# Invariant mass

$$M = \sqrt{\left(\sum E_i\right)^2 - \left(\sum \vec{p}_i\right)^2}$$




# 3-bodies decay



$$|\mathbf{p}_3| = \frac{[(M^2 - (m_{12} + m_3)^2)(M^2 - (m_{12} - m_3)^2)]^{1/2}}{2M}$$


# 3-bodies decay



(a)

$$\max(|\vec{p}_3|)$$


$$\min(|\vec{p}_3|)$$



(b)

$$(m_{12})_{min} = m_1 + m_2$$

$$(m_{12})_{max} = M - m_3$$



(c)

# 3-bodies decay: pion decays


$$|\mathbf{p}_\mu| = \frac{m_\pi^2 - m_\mu^2}{2m_\pi} c \simeq 30 \text{ MeV}/c$$

$$m_\nu = 0$$


in most cases  
muon decays  
at rest

$$|\mathbf{p}_e|_{max} = \frac{m_\mu^2 - m_e^2}{2m_\mu} c \simeq 52 \text{ MeV}/c$$

$$|\mathbf{p}_e|_{min} = 0$$






# 3-bodies decay: Dalitz plot




**Figure 45.3:** Dalitz plot for a three-body final state. In this example, the state is  $\pi^+ \bar{K}^0 p$  at 3 GeV. Four-momentum conservation restricts events to the shaded region.

# Multi-bodies decay





# Reaction threshold




$$\sqrt{s} \geq \sum_i m_i c^2$$

$$\begin{aligned} s &= (p_\pi + p_p)^2 c^2 = (E_\pi + m_p c^2)^2 - |\mathbf{p}_\pi|^2 \\ &= (m_\pi c^2)^2 + (m_p c^2)^2 + 2E_\pi (m_p c^2) \end{aligned}$$

$$E_\pi \geq \frac{(\sum_i m_i c^2)^2 - (m_\pi c^2)^2 - (m_p c^2)^2}{2m_p c^2} \simeq 500 \text{ MeV}$$


# Fixed target vs. collider



How much energy should a fixed target experiment have to equal the center of mass energy of two colliding beam?

$$E_{\text{fix}} = 2 \frac{E_{\text{col}}^2}{m} - m$$

# Collider experiment coordinates



# Rapidity

Lorentz factor  $\gamma = \frac{1}{\sqrt{1 - \beta^2}} = \cosh \varphi$  Hyperbolic cosine of “rapidity”

$$\begin{aligned} E &= m \cosh \varphi & \varphi &= \tanh^{-1} \frac{E}{|\vec{p}|} = \frac{1}{2} \ln \frac{E + |\vec{p}|}{E - |\vec{p}|} \\ |\vec{p}| &= m \sinh \varphi \end{aligned}$$

- Particle physicists prefer to use modified rapidity along beam axis

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z}$$

# Pseudorapidity



$$\eta = \frac{1}{2} \ln \frac{|\vec{p}| + p_z}{|\vec{p}| - p_z}$$

$$\eta = \frac{1}{2} \ln \left( \tan \frac{\theta}{2} \right)$$

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z}$$

$$\eta \simeq y$$

if  $E \gg m$



# Transverse variables

- At hadron colliders, a significant and unknown fraction of the beam energy in each event escapes down the beam pipe.
- Net momentum can only be constrained in the plane transverse to the beam z-axis!

$$\sum p_T(i) = 0$$

$$p_T = \sqrt{p_x^2 + p_y^2}$$

$$p_x = p_T \cos \phi$$

$$p_y = p_T \sin \phi$$

$$p_z = p_T \sinh \eta$$

$$|p| = p_T \cosh \eta$$

$$E_T = \frac{E}{\cosh \eta}$$

# Missing transverse energy and transverse mass

- If invisible particles are created, only their transverse momentum can be constrained: **missing transverse energy**

$$E_T^{\text{miss}} = \sum p_T(i)$$

- If a heavy particle is produced and decays into two particles one of which is invisible, the mass of the parent particle can be constrained with the **transverse mass quantity**

$$\begin{aligned} M_T^2 &\equiv [E_T(1) + E_T(2)]^2 - [\mathbf{p}_T(1) + \mathbf{p}_T(2)]^2 \\ &= m_1^2 + m_2^2 + 2[E_T(1)E_T(2) - \mathbf{p}_T(1) \cdot \mathbf{p}_T(2)] \end{aligned}$$

$$\text{if } m_1 = m_2 = 0 \quad M_T^2 = 2|\mathbf{p}_T(1)||\mathbf{p}_T(2)|(1 - \cos \phi_{12})$$

# $W \rightarrow e \nu$ discovery


Volume 122B, number 1

PHYSICS LETTERS

24 February 1983

**a**


EVENT 2958. 1279.









# ATLAS EXPERIMENT



Run Number: 152409, Event Number: 5966801


Date: 2010-04-05 06:54:50 CEST



$W \rightarrow e \nu$


CDF II preliminary

$$\int L dt \approx 2.2 \text{ fb}^{-1}$$



# Interaction cross section

Flux  $\Phi = \frac{1}{S} \frac{dN_i}{dt}$   $[L^{-2} t^{-1}]$



Reactions per unit of time  $\frac{dN_{\text{reac}}}{dt} = \Phi \overbrace{\sigma N_{\text{target}} dx}^{\text{area obscured by target particle}}$   $[t^{-1}]$

$[L^{-2} t^{-1}]$   $[?]$   $[L^{-1}]$   $[L]$

Reaction rate per target particle  $W_{if} = \Phi \sigma$   $[t^{-1}]$

Cross section per target particle  $\sigma = \frac{W_{if}}{\Phi}$   $[L^2]$  = reaction rate per unit of flux

$1b = 10^{-28} \text{ m}^2$  (roughly the area of a nucleus with  $A = 100$ )

# Fermi Golden Rule

From non-relativistic perturbation theory...


transition probability      matrix element      energy density of final states

$$W_{if} = \frac{2\pi}{\hbar} |M_{if}|^2 \frac{dN}{dE_f}$$

$[\tau^{-1}]$

$[E]$

$[E^{-1}]$



$$M_{if} = -i \int j_{\mu}^{(1)} \left( \frac{1}{q^2} \right) j_{\mu}^{(2)} d^4x$$

$$\sigma \sim |M_{if}|^2 \sim g^4 \left( \frac{1}{q^4} \right)$$

# Cross section: magnitude and units

Standard

cross section unit:

$$[\sigma] = \text{mb}$$

with  $1 \text{ mb} = 10^{-27} \text{ cm}^2$

or in


natural units:

$$[\sigma] = \text{GeV}^{-2}$$

with  $1 \text{ GeV}^{-2} = 0.389 \text{ mb}$

$1 \text{ mb} = 2.57 \text{ GeV}^{-2}$

Estimating the  
proton-proton cross section:



---


using:  $\hbar c = 0.1973 \text{ GeV fm}$   
 $(\hbar c)^2 = 0.389 \text{ GeV}^2 \text{ mb}$

Proton radius:  $R = 0.8 \text{ fm}$


Strong interactions happens up to  $b = 2R$

$$\begin{aligned}\sigma &= \pi(2R)^2 = \pi \cdot 1.6^2 \text{ fm}^2 \\ &= \pi \cdot 1.6^2 \cdot 10^{-26} \text{ cm}^2 \\ &= \pi \cdot 1.6^2 \cdot 10 \text{ mb} \\ &= 80 \text{ mb}\end{aligned}$$

# Proton-proton scattering cross-section



# Cross-sections at LHC



$10^8$  events/s

$\sim 10^{10}$

$10^{-2}$  events/s  $\sim$

10 events/min

[ $m_H \sim 120 \text{ GeV}$ ]

0.2%  $H \rightarrow \gamma\gamma$

1.5%  $H \rightarrow ZZ$

