

Loopfest VII

May 16, 2008

Integral Coefficients for One-Loop Amplitudes

Ruth Britto




University of Amsterdam

One-Loop Amplitudes

Passarino-Veltman reduction brings the one-loop amplitude to the form

$$A_{n;1} = \sum_i d_i (\text{box}) + \sum_i c_i (\text{triangle}) + \sum_i b_i (\text{bubble})$$


where expressions for scalar bubble, scalar triangle and scalar box integrals are known explicitly. (in dim. reg.: Bern, Dixon, Kosower)



Amplitudes from unitarity cuts

$$C = \Delta A_n^{1\text{-loop}} = \sum c \Delta I$$

Tree level data.



Matching 4-dimensional cuts can suffice to determine reduction coefficients!

(Bern, Dixon, Dunbar, Kosower 1994)


But: we get several coefficients together in the same equation.

Unitarity Cuts

$$\Delta A^{1\text{-loop}} = \int d\mu A_{\text{Left}}^{\text{tree}} \times A_{\text{Right}}^{\text{tree}}$$

where


$$d\mu = d^4\ell_1 d^4\ell_2 \delta^{(4)}(\ell_1 + \ell_2 - K) \delta(\ell_1^2) \delta(\ell_2^2)$$



By unitarity, this is the **discontinuity** of the amplitude across a **branch cut**, in a kinematic region selecting the cut momentum K . (Cutkosky 1960)

Box Coefficients from Quadruple Cuts

(RB, Cachazo, Feng)



Generalized Unitarity: Try replacing all four propagators by delta functions.

This operation isolates any given box.

In four dimensions, these four delta functions **localize the integral** completely. This computation is very easy!

The loop momentum solution

The box coefficients computed from quadruple cuts are given by

$$c = \frac{1}{2} \sum_{\mathcal{S}} A_1^{\text{tree}} A_2^{\text{tree}} A_3^{\text{tree}} A_4^{\text{tree}}$$

\mathcal{S} is the set of all solutions of the on-shell conditions for the internal lines.


$$\mathcal{S} = \{ \ell \mid \ell^2 = 0, \quad (\ell - K_1)^2 = 0, \quad (\ell - K_1 - K_2)^2 = 0, \quad (\ell + K_4)^2 = 0 \}$$

Can these equations always be solved?

In [complexified momentum space](#), there are exactly 2 solutions.


(Note: nonvanishing 3-point amplitudes.)

Example: Box Coefficient from Quadruple Cut



$$\begin{aligned}
 \text{coeff} &= \frac{1}{2} \frac{\langle l_1 l_4 \rangle^3}{\langle l_1 2 \rangle \langle 2 l_4 \rangle} \frac{\langle 4 l_2 \rangle^3}{\langle l_2 l_1 \rangle \langle l_1 3 \rangle \langle 3 4 \rangle} \frac{\langle 5 6 \rangle^3}{\langle 6 l_3 \rangle \langle l_3 l_2 \rangle \langle l_2 5 \rangle} \frac{\langle l_3 7 \rangle^3}{\langle 7 1 \rangle \langle 1 l_4 \rangle \langle l_4 l_3 \rangle} \\
 &= - \frac{[1 2]^3 [2 3]^3 \langle 5 6 \rangle^3}{[7 1] [3 4] \langle 5 | P_{3,4} | 2 \rangle \langle 6 | P_{7,1} | 2 \rangle [2 | P_{3,4} P_{5,6} | 7 \rangle [2 | P_{7,1} P_{5,6} | 4 \rangle]}
 \end{aligned}$$

Integral Coefficients from Unitarity Cuts



RHS: cuts of master integrals.

Extract coefficients by **matching** cuts.

Cuts of 4-d Master Integrals

$$\Delta I_2 = \int \langle \ell \, d\ell \rangle [\ell \, d\ell] \frac{K^2}{\langle \ell | K | \ell \rangle^2}$$

$$\Delta I_3 = \int \langle \ell \, d\ell \rangle [\ell \, d\ell] \frac{1}{\langle \ell | K | \ell \rangle \langle \ell | Q | \ell \rangle}$$

$$\Delta I_4 = \int \langle \ell \, d\ell \rangle [\ell \, d\ell] \frac{1}{K^2} \frac{1}{\langle \ell | Q_1 | \ell \rangle \langle \ell | Q_2 | \ell \rangle}$$

$$Q_j \equiv -K_j + \frac{K_j^2}{K^2} K$$

Cutting the Amplitude in 4d

$$C = c \int d^4 \ell \frac{\prod_{i=1}^{k+n} (-2\ell \cdot P_i)}{\prod_{j=1}^k (\ell - K_j)^2} \delta(\ell^2) \delta((\ell - K)^2)$$

We define the following vectors:

$$\begin{aligned} Q_j &= -K_j + \frac{K_j^2}{K^2} K, \\ R_i &= -P_i. \end{aligned}$$

Then the cut integral can be written as follows:

$$C = c \int \langle \ell \, d\ell \rangle [\ell \, d\ell] \frac{(K^2)^{n+1}}{\langle \ell | K | \ell \rangle^{n+2}} \frac{\prod_{i=1}^{k+n} \langle \ell | R_i | \ell \rangle}{\prod_{j=1}^k \langle \ell | Q_j | \ell \rangle}$$

Finish by identifying poles and residues.

We have given the results in general form. (RB, Feng)

See also: Forde

Box coefficients

$$C[K_r, K_s, K] = \frac{(K^2)^{2+n}}{2} \left(\frac{\prod_{j=1}^{k+n} \langle P_{sr,1} | R_j | P_{sr,2} \rangle}{\langle P_{sr,1} | K | P_{sr,2} \rangle^{n+2} \prod_{t=1, t \neq i, j}^k \langle P_{sr,1} | Q_t | P_{sr,2} \rangle} + \{P_{sr,1} \leftrightarrow P_{sr,2}\} \right)$$

$$P_{sr,1} = Q_s + \left(\frac{-2Q_s \cdot Q_r + \sqrt{\Delta_{sr}}}{2Q_r^2} \right) Q_r$$

$$P_{sr,2} = Q_s + \left(\frac{-2Q_s \cdot Q_r - \sqrt{\Delta_{sr}}}{2Q_r^2} \right) Q_r$$

$$\Delta_{sr} = (2Q_s \cdot Q_r)^2 - 4Q_s^2 Q_r^2$$

Triangle coefficients

$$C[K_s, K] = \frac{(K^2)^{1+n}}{2} \frac{1}{(\sqrt{\Delta_s})^{n+1}} \frac{1}{(n+1)! \langle P_{s,1} P_{s,2} \rangle^{n+1}} \\ \times \frac{d^{n+1}}{d\tau^{n+1}} \left(\frac{\prod_{j=1}^{k+n} \langle P_{s,1} - \tau P_{s,2} | R_j Q_s | P_{s,1} - \tau P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} - \tau P_{s,2} | Q_t Q_s | P_{s,1} - \tau P_{s,2} \rangle} + \{P_{s,1} \leftrightarrow P_{s,2}\} \right) \Big|_{\tau=0} .$$

$$P_{s,1} = Q_s + \left(\frac{-2Q_s \cdot K + \sqrt{\Delta_s}}{2K^2} \right) K$$

$$P_{s,2} = Q_s + \left(\frac{-2Q_s \cdot K - \sqrt{\Delta_s}}{2K^2} \right) K$$

$$\Delta_s = (2Q_s \cdot K)^2 - 4Q_s^2 K^2$$

Bubble coefficients

$$C[K] = (K^2)^{1+n} \sum_{q=0}^n \frac{(-1)^q}{q!} \frac{d^q}{ds^q} \left(\mathcal{B}_{n,n-q}^{(0)}(s) + \sum_{r=1}^k \sum_{a=q}^n \left(\mathcal{B}_{n,n-a}^{(r;a-q;1)}(s) - \mathcal{B}_{n,n-a}^{(r;a-q;2)}(s) \right) \right) \Big|_{s=0}$$

$$\mathcal{B}_{n,t}^{(0)}(s) \equiv \frac{d^n}{d\tau^n} \left(\frac{1}{n![\eta|\eta'K|\eta]^n} \frac{(2\eta \cdot K)^{t+1}}{(t+1)(K^2)^{t+1}} \frac{\prod_{j=1}^{n+k} \langle \ell | R_j(K+s\eta) | \ell \rangle}{\langle \ell \eta \rangle^{n+1} \prod_{p=1}^k \langle \ell | Q_p(K+s\eta) | \ell \rangle} \right) \Big|_{|\ell\rangle \rightarrow |K-\tau\eta'|\eta\rangle, \tau=0}$$

$$\mathcal{B}_{n,t}^{(r;b;1)}(s) \equiv \frac{(-1)^{b+1}}{b! \sqrt{\Delta_r}^{b+1} \langle P_{r,1} P_{r,2} \rangle^b} \frac{d^b}{d\tau^b} \left(\frac{1}{(t+1)} \frac{\langle P_{r,1} - \tau P_{r,2} | \eta | P_{r,1} \rangle^{t+1}}{\langle P_{r,1} - \tau P_{r,2} | K | P_{r,1} \rangle^{t+1}} \right. \\ \left. \times \frac{\langle P_{r,1} - \tau P_{r,2} | Q_r \eta | P_{r,1} - \tau P_{r,2} \rangle^b \prod_{j=1}^{n+k} \langle P_{r,1} - \tau P_{r,2} | R_j(K+s\eta) | P_{r,1} - \tau P_{r,2} \rangle}{\langle P_{r,1} - \tau P_{r,2} | \eta K | P_{r,1} - \tau P_{r,2} \rangle^{n+1} \prod_{p=1, p \neq r}^k \langle P_{r,1} - \tau P_{r,2} | Q_p(K+s\eta) | P_{r,1} - \tau P_{r,2} \rangle} \right) \Big|_{\tau=0}$$

$$\mathcal{B}_{n,t}^{(r;b;2)}(s) \equiv \frac{(-1)^{b+1}}{b! \sqrt{\Delta_r}^{b+1} \langle P_{r,1} P_{r,2} \rangle^b} \frac{d^b}{d\tau^b} \left(\frac{1}{(t+1)} \frac{\langle P_{r,2} - \tau P_{r,1} | \eta | P_{r,2} \rangle^{t+1}}{\langle P_{r,2} - \tau P_{r,1} | K | P_{r,2} \rangle^{t+1}} \right. \\ \left. \times \frac{\langle P_{r,2} - \tau P_{r,1} | Q_r \eta | P_{r,2} - \tau P_{r,1} \rangle^b \prod_{j=1}^{n+k} \langle P_{r,2} - \tau P_{r,1} | R_j(K+s\eta) | P_{r,2} - \tau P_{r,1} \rangle}{\langle P_{r,2} - \tau P_{r,1} | \eta K | P_{r,2} - \tau P_{r,1} \rangle^{n+1} \prod_{p=1, p \neq r}^k \langle P_{r,2} - \tau P_{r,1} | Q_p(K+s\eta) | P_{r,2} - \tau P_{r,1} \rangle} \right) \Big|_{\tau=0}$$

D-dimensional unitarity

Orthogonal decomposition within Four Dimensional Helicity (FDH) scheme.

(Mahlon; Bern, Chalmers; Bern, Morgan)

$$\begin{aligned}\int d^{4-2\epsilon} \ell_{4-2\epsilon} &= \int d^{-2\epsilon} \ell_{-2\epsilon} \int d^4 \ell_4 \\ &= \frac{(4\pi)^\epsilon}{\Gamma(-\epsilon)} \int_0^1 du u^{-1-\epsilon} \int d^4 \tilde{\ell}.\end{aligned}$$

where $\ell_{-2\epsilon}^2 = \frac{K^2}{4} u$.

The integral over u will remain. The u -dependence is controlled.

(Anastasiou, RB, Feng, Kunszt, Mastrolia)

$$\Delta A = \int_0^1 du u^{-1-\epsilon} \int d^4 \ell \delta(\ell^2) \delta(\sqrt{1-u} K^2 - 2K \cdot \ell)$$

Problem is reduced to a standard 4-d cut integral.

(Cf. methods by Ossola, Papadopoulos, Pittau; Ellis, Giele, Kunszt; Kilgore; Giele, Kunszt, Melnikov)

D-dimensional unitarity algorithm

$$\Delta A = \int_0^1 du u^{-1-\epsilon} \int d^4 \ell \delta(\ell^2) \delta(\sqrt{1-u} K^2 - 2K \cdot \ell)$$

1. 4d cut: get u -dependent coefficients of master integrals.

u -dependence is polynomial. (RB, Feng, Yang; RB, Feng, Mastrolia)

2. Treat polynomial u -dependence of integrand. Two choices:

(a) For each term in the polynomial, use shift identities to get coefficients of 4d master integrals.

(b) Use dimensionally shifted master integrals.

The u -integral is not done explicitly.

Here, u is like the \tilde{q}^2 of Ossola, Papadopoulos, Pittau; or the s_e^2 of Giele, Kunszt, Melnikov.

Coefficients are polynomials in u

The (maximum) degrees are the following:

Pentagon: 0

Box: $[(n + 2)/2]$

Triangle: $[(n + 1)/2]$

Bubble: $[n/2]$

$$C = c \int d^{4-2\epsilon} p \frac{\prod_{i=1}^{k+n} (-2p \cdot P_i)}{\prod_{j=1}^k (p - K_j)^2} \delta(p^2) \delta((p - K)^2)$$

The terms in the polynomials

Analytically,

$$C(u) = \sum_{s=0}^d \left(\frac{1}{s!} \frac{d^s C(u)}{du^s} \Big|_{u \rightarrow 0} \right) u^s.$$

Or, numerically,

1. Define

$$C_k \equiv C(u_k),$$

for $(k = 0, \dots, d - 1)$, where

$$u_k = e^{-2\pi i k/d}.$$

2. Then,

$$C(u) = \sum_{s=0}^d \left(\frac{1}{d} \sum_{k=0}^{d-1} C_k e^{2\pi i s k/d} \right) u^s.$$

Dimensional shift identities

(Anastasiou, RB, Feng, Kunstz, Mastrolia)

$$\text{Bub}^{(n)} = F_{2 \rightarrow 2}^{(n)} \text{Bub}^{(0)}$$

$$\text{Tri}^{(n)} = F_{3 \rightarrow 3}^{(n)} \text{Tri}^{(0)} + F_{3 \rightarrow 2}^{(n)} \text{Bub}^{(0)}$$

$$\text{Box}^{(n)} = F_{4 \rightarrow 4}^{(n)} \text{Box}^{(0)} + F_{4 \rightarrow 3}^{(n)} \text{Tri}^{(0)} + F_{4 \rightarrow 2}^{(n)} \text{Bub}^{(0)}$$

$$F_{2 \rightarrow 2}^{(n)} = \frac{(-\epsilon)_{\frac{3}{2}}}{(n-\epsilon)_{\frac{3}{2}}}, \quad F_{3 \rightarrow 3}^{(n)} = \frac{-\epsilon}{n-\epsilon} (1-Z^2)^n,$$

$$F_{4 \rightarrow 4}^{(n)} = \frac{(-\epsilon)_{\frac{1}{2}}}{(n-\epsilon)_{\frac{1}{2}}} \left(\frac{B}{A} \right)^n,$$

$$F_{3 \rightarrow 2}^{(n)} = \frac{(-\epsilon)_{\frac{3}{2}}}{n-\epsilon} \sum_{k=1}^n \frac{2Z(1-Z^2)^{n-k}}{(k-\epsilon)_{\frac{1}{2}}}$$

$$F_{4 \rightarrow j}^{(n)} = \frac{D + (Z^2 - 1)C}{(n-\epsilon)_{\frac{1}{2}} Z A} \sum_{k=1}^n \left(\frac{B}{A} \right)^{n-k} (k-1-\epsilon)_{\frac{1}{2}} F_{3 \rightarrow j}^{(k-1)}$$

Cuts of D-dimensional Master Integrals

$$\Delta I_2 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] K^2 \sqrt{1-u} \frac{1}{\langle \ell | K | \ell \rangle^2}$$

$$\Delta I_3 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \sqrt{1-u} \frac{1}{\langle \ell | K | \ell \rangle \langle \ell | Q | \ell \rangle}$$

$$\Delta I_4 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \frac{\sqrt{1-u}}{K^2} \frac{1}{\langle \ell | Q_1 | \ell \rangle \langle \ell | Q_2 | \ell \rangle}$$

$$\Delta I_4 = \frac{1}{2K^2} \frac{\sqrt{1-u}}{\sqrt{(Q_1 \cdot Q_2)^2 - Q_1^2 Q_2^2}} \ln \frac{Q_1 \cdot Q_2 + \sqrt{(Q_1 \cdot Q_2)^2 - Q_1^2 Q_2^2}}{Q_1 \cdot Q_2 - \sqrt{(Q_1 \cdot Q_2)^2 - Q_1^2 Q_2^2}}$$

$$\begin{aligned} \Delta I_5 = & \frac{\sqrt{1-u}}{(K^2)^2} \left(\frac{S[Q_3, Q_2, Q_1, K]}{4\sqrt{(Q_3 \cdot Q_2)^2 - Q_3^2 Q_2^2}} \ln \frac{Q_3 \cdot Q_2 + \sqrt{(Q_3 \cdot Q_2)^2 - Q_3^2 Q_2^2}}{Q_3 \cdot Q_2 - \sqrt{(Q_3 \cdot Q_2)^2 - Q_3^2 Q_2^2}} \right. \\ & + \frac{S[Q_3, Q_1, Q_2, K]}{4\sqrt{(Q_3 \cdot Q_1)^2 - Q_3^2 Q_1^2}} \ln \frac{Q_3 \cdot Q_1 + \sqrt{(Q_3 \cdot Q_1)^2 - Q_3^2 Q_1^2}}{Q_3 \cdot Q_1 - \sqrt{(Q_3 \cdot Q_1)^2 - Q_3^2 Q_1^2}} \\ & \left. + \frac{S[Q_2, Q_1, Q_3, K]}{4\sqrt{(Q_2 \cdot Q_1)^2 - Q_2^2 Q_1^2}} \ln \frac{Q_2 \cdot Q_1 + \sqrt{(Q_2 \cdot Q_1)^2 - Q_2^2 Q_1^2}}{Q_2 \cdot Q_1 - \sqrt{(Q_2 \cdot Q_1)^2 - Q_2^2 Q_1^2}} \right). \end{aligned}$$

$$S[Q_i, Q_j, Q_k, K] = \frac{T_1}{T_2}$$

$$T_1 = -8 \det \begin{pmatrix} K \cdot Q_k & Q_i \cdot K & Q_j \cdot K \\ Q_i \cdot Q_k & Q_i^2 & Q_i \cdot Q_j \\ Q_j \cdot Q_k & Q_i \cdot Q_j & Q_j^2 \end{pmatrix}; \quad T_2 = -4 \det \begin{pmatrix} Q_k^2 & Q_i \cdot Q_k & Q_j \cdot Q_k \\ Q_i \cdot Q_k & Q_i^2 & Q_i \cdot Q_j \\ Q_j \cdot Q_k & Q_i \cdot Q_j & Q_j^2 \end{pmatrix}.$$

Cutting the Amplitude in D dimensions

$$C = c \int d^{4-2\epsilon} p \frac{\prod_{i=1}^{k+n} (-2p \cdot P_i)}{\prod_{j=1}^k (p - K_j)^2} \delta(p^2) \delta((p - K)^2)$$

Let us define the following four-vectors:

$$Q_j = -(\sqrt{1-u})K_j + \frac{K_j^2 - (1 - \sqrt{1-u})(K_j \cdot K)}{K^2} K,$$

$$R_i = -(\sqrt{1-u})P_i - \frac{(1 - \sqrt{1-u})(P_i \cdot K)}{K^2} K.$$

Then the cut integral can be written as follows:

$$C = c \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] (\sqrt{1-u}) \frac{(K^2)^{n+1}}{\langle \ell | K | \ell \rangle^{n+2}} \frac{\prod_{i=1}^{k+n} \langle \ell | R_i | \ell \rangle}{\prod_{j=1}^k \langle \ell | Q_j | \ell \rangle}$$

Coefficient formulas look the same, now in terms of u -dependent Q_j, R_i .

We have checked 4- and 5-gluon examples to all orders in ϵ . (Anastasiou, RB, Feng, Kunszt, Mastrolia; RB, Feng, Yang. Original results from Bern, Dixon, Dunbar, Kosower.)

Box coefficients

$$C[K_r, K_s, K] = \frac{(K^2)^{2+n}}{2} \left(\frac{\prod_{j=1}^{k+n} \langle P_{sr,1} | R_j | P_{sr,2} \rangle}{\langle P_{sr,1} | K | P_{sr,2} \rangle^{n+2} \prod_{t=1, t \neq i, j}^k \langle P_{sr,1} | Q_t | P_{sr,2} \rangle} + \{P_{sr,1} \leftrightarrow P_{sr,2}\} \right)$$

$$P_{sr,1} = Q_s + \left(\frac{-2Q_s \cdot Q_r + \sqrt{\Delta_{sr}}}{2Q_r^2} \right) Q_r$$

$$P_{sr,2} = Q_s + \left(\frac{-2Q_s \cdot Q_r - \sqrt{\Delta_{sr}}}{2Q_r^2} \right) Q_r$$

$$\Delta_{sr} = (2Q_s \cdot Q_r)^2 - 4Q_s^2 Q_r^2$$

Pentagon coefficients

(RB, Feng, Yang)

If $k = 3$:

$$C[K_i, K_j, K_t, K] = (K^2)^{3+n} \prod_{s=1}^{n+3} \beta_s^{(q_i, q_j, q_t; p_s)}$$

If $k \geq 4$:

$$C[K_i, K_j, K_t, K] = (K^2)^{3+n} \frac{\prod_{s=1}^{n+3} \beta_s^{(q_i, q_j, q_t; p_s)}}{\prod_{w=1, w \neq i, j, t}^k \gamma_w^{(K_i, K_j; K_w, K_t)}}.$$

$$\alpha_j \equiv \frac{K_j^2 - K_j \cdot K}{K^2}$$

$$\beta_s^{(q_i, q_j, q_t; p_s)} \equiv \left(\beta_s - \sum_{h=i, j, k} \alpha_h^{(q_i, q_j, q_t; p_s)} \right)$$

$$\gamma_s^{(K_i, K_j; K_s, K_t)} \equiv \frac{K_i^2 \epsilon(K, K_j, K_s, K_t) + K_j^2 \epsilon(K_i, K, K_s, K_t) + K_s^2 \epsilon(K_i, K_j, K, K_t) + K_t^2 \epsilon(K_i, K_j, K_s, K)}{K^2 \epsilon(K_i, K_j, K, K_t)}.$$

Triangle coefficients

$$C[K_s, K] = \frac{(K^2)^{1+n}}{2} \frac{1}{(\sqrt{\Delta_s})^{n+1}} \frac{1}{(n+1)! \langle P_{s,1} P_{s,2} \rangle^{n+1}} \\ \times \frac{d^{n+1}}{d\tau^{n+1}} \left(\frac{\prod_{j=1}^{k+n} \langle P_{s,1} - \tau P_{s,2} | R_j Q_s | P_{s,1} - \tau P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} - \tau P_{s,2} | Q_t Q_s | P_{s,1} - \tau P_{s,2} \rangle} + \{P_{s,1} \leftrightarrow P_{s,2}\} \right) \Big|_{\tau=0} .$$

$$P_{s,1} = Q_s + \left(\frac{-2Q_s \cdot K + \sqrt{\Delta_s}}{2K^2} \right) K$$

$$P_{s,2} = Q_s + \left(\frac{-2Q_s \cdot K - \sqrt{\Delta_s}}{2K^2} \right) K$$

$$\Delta_s = (2Q_s \cdot K)^2 - 4Q_s^2 K^2$$

Bubble coefficients

$$C[K] = (K^2)^{1+n} \sum_{q=0}^n \frac{(-1)^q}{q!} \frac{d^q}{ds^q} \left(\mathcal{B}_{n,n-q}^{(0)}(s) + \sum_{r=1}^k \sum_{a=q}^n \left(\mathcal{B}_{n,n-a}^{(r;a-q;1)}(s) - \mathcal{B}_{n,n-a}^{(r;a-q;2)}(s) \right) \right) \Big|_{s=0}$$

$$\mathcal{B}_{n,t}^{(0)}(s) \equiv \frac{d^n}{d\tau^n} \left(\frac{1}{n![\eta|\eta'K|\eta]^n} \frac{(2\eta \cdot K)^{t+1}}{(t+1)(K^2)^{t+1}} \frac{\prod_{j=1}^{n+k} \langle \ell | R_j(K + s\eta) | \ell \rangle}{\langle \ell \eta \rangle^{n+1} \prod_{p=1}^k \langle \ell | Q_p(K + s\eta) | \ell \rangle} \right) \Big|_{|\ell\rangle \rightarrow |K - \tau\eta'|\eta\rangle, \tau=0}$$

$$\mathcal{B}_{n,t}^{(r;b;1)}(s) \equiv \frac{(-1)^{b+1}}{b! \sqrt{\Delta_r}^{b+1} \langle P_{r,1} P_{r,2} \rangle^b} \frac{d^b}{d\tau^b} \left(\frac{1}{(t+1)} \frac{\langle P_{r,1} - \tau P_{r,2} | \eta | P_{r,1} \rangle^{t+1}}{\langle P_{r,1} - \tau P_{r,2} | K | P_{r,1} \rangle^{t+1}} \right. \\ \left. \times \frac{\langle P_{r,1} - \tau P_{r,2} | Q_r \eta | P_{r,1} - \tau P_{r,2} \rangle^b \prod_{j=1}^{n+k} \langle P_{r,1} - \tau P_{r,2} | R_j(K + s\eta) | P_{r,1} - \tau P_{r,2} \rangle}{\langle P_{r,1} - \tau P_{r,2} | \eta K | P_{r,1} - \tau P_{r,2} \rangle^{n+1} \prod_{p=1, p \neq r}^k \langle P_{r,1} - \tau P_{r,2} | Q_p(K + s\eta) | P_{r,1} - \tau P_{r,2} \rangle} \right) \Big|_{\tau=0}$$

$$\mathcal{B}_{n,t}^{(r;b;2)}(s) \equiv \frac{(-1)^{b+1}}{b! \sqrt{\Delta_r}^{b+1} \langle P_{r,1} P_{r,2} \rangle^b} \frac{d^b}{d\tau^b} \left(\frac{1}{(t+1)} \frac{\langle P_{r,2} - \tau P_{r,1} | \eta | P_{r,2} \rangle^{t+1}}{\langle P_{r,2} - \tau P_{r,1} | K | P_{r,2} \rangle^{t+1}} \right. \\ \left. \times \frac{\langle P_{r,2} - \tau P_{r,1} | Q_r \eta | P_{r,2} - \tau P_{r,1} \rangle^b \prod_{j=1}^{n+k} \langle P_{r,2} - \tau P_{r,1} | R_j(K + s\eta) | P_{r,2} - \tau P_{r,1} \rangle}{\langle P_{r,2} - \tau P_{r,1} | \eta K | P_{r,2} - \tau P_{r,1} \rangle^{n+1} \prod_{p=1, p \neq r}^k \langle P_{r,2} - \tau P_{r,1} | Q_p(K + s\eta) | P_{r,2} - \tau P_{r,1} \rangle} \right) \Big|_{\tau=0}$$

These coefficients are polynomials in u .

Proof is constructive. For alternate formulas with more transparent u -dependence, see [arXiv:0803.3147](https://arxiv.org/abs/0803.3147). (RB, Feng, Yang)

Incorporating Masses

(RB, Feng; RB, Feng, Mastrolia)

$$C = c \int d^{4-2\epsilon} p \frac{\prod_{i=1}^{k+n} (-2p \cdot P_i)}{\prod_{j=1}^k ((p - K_j)^2 - m_j^2)} \delta(p^2 - M_1^2) \delta((p - K)^2 - M_2^2)$$

We define the following four-vectors:

$$Q_j = - \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) K_j + \frac{K_j^2 + M_1^2 - m_j^2 - 2zK \cdot K_j}{K^2} K$$

$$R_i = - \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) P_i - \frac{z(2P_i \cdot K)}{K^2} K,$$

where

$$z = \frac{\alpha - \beta \sqrt{1 - u}}{2}$$

$$\alpha = \frac{K^2 + M_1^2 - M_2^2}{K^2},$$

$$\beta = \frac{\sqrt{(K^2)^2 + (M_1^2)^2 + (M_2^2)^2 - 2K^2 M_1^2 - 2K^2 M_2^2 - 2M_1^2 M_2^2}}{K^2}$$

Cut amplitude:

$$\int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) \frac{(K^2)^{n+1}}{\langle \ell | K | \ell \rangle^{n+2}} \frac{\prod_{j=1}^{n+k} \langle \ell | R_j | \ell \rangle}{\prod_{i=1}^k \langle \ell | Q_i | \ell \rangle}$$

Cut masters:

$$\Delta I_2 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) \frac{(K^2)}{\langle \ell | K | \ell \rangle^2}$$

$$\Delta I_3 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) \frac{1}{\langle \ell | K | \ell \rangle \langle \ell | Q | \ell \rangle}$$

$$\Delta I_4 = \int_0^1 du u^{-1-\epsilon} \int \langle \ell d\ell \rangle [\ell d\ell] \left((1 - 2z) + \frac{M_1^2 - M_2^2}{K^2} \right) \frac{(K^2)^{-1}}{\langle \ell | Q_1 | \ell \rangle \langle \ell | Q_2 | \ell \rangle}$$

Again, the formulas for integral coefficients will look the same!

See also: Kilgore

We have reproduced cuts for $gg \rightarrow gg$ and $gg \rightarrow gH$ with massive fermion loops. (RB, Feng, Mastrolia. Original results from Bern, Morgan; Rozowsky.)

Coefficients of cut-free integrals are to be fixed by universal divergent behavior, or other constraints. (Bern, Morgan; Kilgore)

How were these formulas derived?

$$C = c \int d^4 \ell \frac{\prod_{i=1}^{k+n} (-2\ell \cdot P_i)}{\prod_{j=1}^k (\ell - K_j)^2} \delta(\ell^2) \delta((\ell - K)^2)$$

Change to spinor variables: $\ell_{a\dot{a}} = t \lambda_a \tilde{\lambda}_{\dot{a}}$. (Cachazo, Svrček, Witten)

$$\int d^4 \ell \delta(\ell^2) \delta((\ell - K)^2) (\bullet) = \int_0^\infty dt t \int \langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}] \delta((t\lambda\tilde{\lambda} - K)^2) (\bullet)$$

Use 2nd delta function to perform t -integral.

$$C = \int \langle \lambda d\lambda \rangle [\tilde{\lambda} d\tilde{\lambda}] \frac{(K^2)^{n+1}}{\langle \lambda | K | \tilde{\lambda} \rangle^{n+2}} \frac{\prod_{i=1}^{k+n} \langle \lambda | R_i | \tilde{\lambda} \rangle}{\prod_{j=1}^k \langle \lambda | Q_j | \tilde{\lambda} \rangle}$$

$$C = \int \langle \ell d\ell \rangle [\ell d\ell] \frac{(K^2)^{n+1}}{\langle \ell | K | \ell \rangle^{n+2}} \frac{\prod_{i=1}^{k+n} \langle \ell | R_i | \ell \rangle}{\prod_{j=1}^k \langle \ell | Q_j | \ell \rangle}$$

Split the factors in the denominator with **partial fractions**.

$$\frac{\prod_{j=1}^{k-1} [a_j \ell]}{\prod_{i=1}^k \langle \ell | Q_i | \ell \rangle} = \sum_{i=1}^k \frac{1}{\langle \ell | Q_i | \ell \rangle} \frac{\prod_{j=1}^{k-1} [a_j | Q_i | \ell]}{\prod_{m=1, m \neq i}^k \langle \ell | Q_m Q_i | \ell \rangle} \quad \text{box \& pentagon}$$

$$\begin{aligned} \frac{\prod_{j=1}^{n-1} [a_j \ell]}{\langle \ell | K | \ell \rangle^n \langle \ell | Q | \ell \rangle} &= \frac{\prod_{j=1}^{n-1} [a_j | Q | \ell]}{\langle \ell | K Q | \ell \rangle^{n-1}} \frac{1}{\langle \ell | K | \ell \rangle \langle \ell | Q | \ell \rangle} \quad \text{triangle} \\ &+ \sum_{p=0}^{n-2} (-1)^{n-p} \frac{\prod_{j=1}^{n-p-2} [a_j | Q | \ell] [a_{n-p-1} | K | \ell] \prod_{t=n-p}^{n-1} [a_t \ell]}{\langle \ell | K | \ell \rangle^{p+2} \langle \ell | Q K | \ell \rangle^{n-p-1}} \quad \text{bubble} \end{aligned}$$

Multiple poles lead to derivatives.

Alternate formula: triangle coefficients without derivatives

$$C[K_s, K]_{n=-2} = 0$$

$$C[K_s, K]_{n=-1} = \frac{1}{2} \left(\frac{\prod_{j=1}^{k-1} \langle P_{s,1} | R_j | P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} | Q_t | P_{s,2} \rangle} \right)$$

$$C[K_s, K]_{n=0} = \frac{K^2}{2\Delta_s} \left[\frac{\prod_{j=1}^k \langle P_{s,1} | R_j | P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} | Q_t | P_{s,2} \rangle} \left(\sum_{j=1}^k \frac{(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)}{\langle P_{s,1} | R_j | P_{s,2} \rangle} \right. \right. \\ \left. \left. - \sum_{t=1, t \neq s}^k \frac{(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)}{\langle P_{s,1} | Q_t | P_{s,2} \rangle} \right) + \{P_{s,1} \leftrightarrow P_{s,2}\} \right]$$

$$\begin{aligned}
C[K_s, K]_{n=1} = & \\
& \frac{(K^2)^2}{4\Delta_s^2} \left[\frac{\prod_{j=1}^{k+1} \langle P_{s,1} | R_j | P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} | Q_t | P_{s,2} \rangle} \left[\left(\sum_{j=1}^{k+1} \frac{(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)}{\langle P_{s,1} | R_j | P_{s,2} \rangle} \right)^2 \right. \right. \\
& - \left. \sum_{t=1, t \neq s}^k \frac{(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)}{\langle P_{s,1} | Q_t | P_{s,2} \rangle} \right]^2 \\
& + \sum_{j=1}^{k+1} \frac{-[(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)]^2 + 2Q_s^2 K^2 \langle P_{s,1} | R_j | P_{s,2} \rangle \langle P_{s,2} | R_j | P_{s,1} \rangle}{\langle P_{s,1} | R_j | P_{s,2} \rangle^2} \\
& - \sum_{t=1, t \neq s}^k \frac{-[(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)]^2 + 2Q_s^2 K^2 \langle P_{s,1} | Q_t | P_{s,2} \rangle \langle P_{s,2} | Q_t | P_{s,1} \rangle}{\langle P_{s,1} | Q_t | P_{s,2} \rangle^2} \\
& + \{P_{s,1} \leftrightarrow P_{s,2}\}
\end{aligned}$$

$$C[K_s, K]_{n=2} = \frac{(K^2)^3}{12\Delta_s^3} \left[\frac{\prod_{j=1}^{k+2} \langle P_{s,1} | R_j | P_{s,2} \rangle}{\prod_{t=1, t \neq s}^k \langle P_{s,1} | Q_t | P_{s,2} \rangle} (\mathcal{A}^3 + 3\mathcal{A}\mathcal{B} + \mathcal{C}) + \{P_{s,1} \leftrightarrow P_{s,2}\} \right]$$

$$\begin{aligned} \mathcal{A} &= \sum_{j=1}^{k+2} \frac{(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)}{\langle P_{s,1} | R_j | P_{s,2} \rangle} - \sum_{t=1, t \neq s}^k \frac{(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)}{\langle P_{s,1} | Q_t | P_{s,2} \rangle} \\ \mathcal{B} &= - \sum_{j=1}^{k+2} \frac{[(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)]^2 + 2Q_s^2 K^2 \langle P_{s,1} | R_j | P_{s,2} \rangle \langle P_{s,2} | R_j | P_{s,1} \rangle}{\langle P_{s,1} | R_j | P_{s,2} \rangle^2} \\ &\quad + \sum_{t=1, t \neq s}^k \frac{[(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)]^2 - 2Q_s^2 K^2 \langle P_{s,1} | Q_t | P_{s,2} \rangle \langle P_{s,2} | Q_t | P_{s,1} \rangle}{\langle P_{s,1} | Q_t | P_{s,2} \rangle^2} \\ \mathcal{C} &= \sum_{j=1}^{k+2} \frac{[(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)]^3 - 3Q_s^2 K^2 \langle P_{s,1} | R_j | P_{s,2} \rangle \langle P_{s,2} | R_j | P_{s,1} \rangle}{\langle P_{s,1} | R_j | P_{s,2} \rangle^2} \\ &\quad - \sum_{t=1, t \neq s}^k \frac{2[(2Q_s \cdot K)(2R_j \cdot Q_s) - 2Q_s^2(2R_j \cdot K)]}{\langle P_{s,1} | R_j | P_{s,2} \rangle} - \sum_{t=1, t \neq s}^k \frac{2[(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)]}{\langle P_{s,1} | Q_t | P_{s,2} \rangle} \\ &\quad - \frac{[(2Q_s \cdot K)(2Q_t \cdot Q_s) - 2Q_s^2(2Q_t \cdot K)]^2 - 3Q_s^2 K^2 \langle P_{s,1} | Q_t | P_{s,2} \rangle \langle P_{s,2} | Q_t | P_{s,1} \rangle}{\langle P_{s,1} | Q_t | P_{s,2} \rangle^2} \end{aligned}$$

Summary

- Coefficients of master integrals can be obtained without explicit reduction in a D -dimensional unitarity method.
- General analytic formulas available for the massless case and in D dimensions.