

Particle Physics

The Standard Model

Antonio Pich

IFIC, CSIC – Univ. Valencia

Antonio.Pich@ific.uv.es

The Standard Model

1. Constituents & Interactions
2. Quarks
3. Gauge Invariance
4. Quantum Chromodynamics
5. Electroweak Unification
6. Symmetry Breaking
7. Electroweak Phenomenology
8. Flavour Dynamics

1. Constituents & Interactions

- Table of Elementary Fermions
- Interactions: Gauge Bosons
- Charged Leptons
- Neutrinos

Periodic Table of the Elements

1	IA																																0																	
1	H	IIA																																																He
2	Li	Be																	B	C	N	O	F	Ne																										
3	Na	Mg	III B IV B V B VI B VII B — VII — IB IB																Al	Si	P	S	Cl	Ar																										
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr																																
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe																																
6	Cs	Ba	* La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn																																
7	Fr	Ra	+ Ac	Rf	Ha	106	107	108	109	110	111	112																																						


Naming conventions of new elements

* Lanthanide Series

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu


+ Actinide Series


90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr






The Sta








p


p


p


n

n










<p>ALL ORDINARY MATTER BELONGS TO THIS GROUP.</p> <p>THESE PARTICLES EXISTED JUST AFTER THE BIG BANG.</p> <p>NOW THEY ARE FOUND ONLY IN COSMIC RAYS AND ACCELERATORS.</p>	LEPTONS		QUARKS	
	<p>electron Electric charge -1. Responsible for electricity and chemical reactions</p>	<p>electron neutrino Electric charge 0. Rarely interacts with other matter.</p>	<p>up Electric charge $+2/3$. Protons have 2 up quarks Neutrons have 1 up quark</p>	<p>down Electric charge $-1/3$. ... and one down quark. ... and two down quarks.</p>
<p>muon A heavier relative of the electron.</p>	<p>muon neutrino Created with muons when some particles decay.</p>	<p>charm A heavier relative of the up.</p>	<p>strange A heavier relative of the down.</p>	
<p>tau Heavier still.</p>	<p>tau neutrino Not yet observed directly.</p>	<p>top Heavier still, recently observed.</p>	<p>bottom Heavier still.</p>	




<p>ALL ORDINARY MATTER BELONGS TO THIS GROUP.</p>	LEPTONS		QUARKS	
	<p>THESE PARTICLES EXISTED JUST AFTER THE BIG BANG.</p> <p>NOW THEY ARE FOUND ONLY IN COSMIC RAYS AND ACCELERATORS.</p>	<p>electron Electric charge -1. Responsible for electricity and chemical reactions</p>	<p>electron neutrino Electric charge 0. Rarely interacts with other matter.</p>	<p>up Electric charge $+2/3$. Protons have 2 up quarks Neutrons have 1 up quark</p>
	<p>muon A heavier relative of the electron.</p>	<p>muon neutrino Created with muons when some particles decay.</p>	<p>charm A heavier relative of the up.</p>	<p>strange A heavier relative of the down.</p>
	<p>tau Heavier still.</p>	<p>tau neutrino Not yet observed directly.</p>	<p>top Heavier still, recently observed.</p>	<p>bottom Heavier still.</p>

ANTIMATTER
Each particle also has an antimatter counterpart ... sort of a mirror image.



The forces in Nature

TYPE	INTENSITY OF FORCES (DECREASING ORDER)	BINDING PARTICLE (FIELD QUANTUM)	OCCURS IN :
STRONG NUCLEAR FORCE	~ 1	GLUONS (NO MASS)	ATOMIC NUCLEUS
ELECTRO -MAGNETIC FORCE	$\sim 10^{-3}$	PHOTONS (NO MASS)	ATOMIC SHELL ELECTROTECHNIQUE
WEAK NUCLEAR FORCE	$\sim 10^{-5}$	BOSONS Z^0, W^+, W^- (HEAVY)	RADIOACTIVE BETA DESINTEGRATION
GRAVITATION	$\sim 10^{-38}$	GRAVITONS (?)	HEAVENLY BODIES




RADIOACTIVITY

(β Decay)



$$n \rightarrow p + e^{-} + \bar{\nu}_e$$

$$"p \rightarrow n + e^{+} + \nu_e"$$



$$Q_{\nu_e} = Q_{\bar{\nu}_e} = 0$$

$$m_{\nu_e} = m_{\bar{\nu}_e} \approx 0$$

$\nu_e \equiv$ Neutrino ; $\bar{\nu}_e \equiv$ Anti-Neutrino

$$p p \rightarrow (p n) + e^{+} + \nu_e$$



Weak Interaction

$$W^{\pm}, Z^0$$

$$M_W \sim M_Z \approx 100 m_p$$

Quarks



up



down



charm



strange



top



beauty

Leptons



electron



neutrino e



muon



neutrino μ



tau




neutrino τ



photon



gluon




Z^0 W^\pm



Higgs

QM + Relativity




Antiparticles


(Dirac)

ANTIMATTER

u	d	ν_e	e^-
\bar{u}	\bar{d}	$\bar{\nu}_e$	e^+



$$E = m c^2$$



Quarks



up



down



charm



strange



top



beauty

Leptons



electron



neutrino e



muon



neutrino μ



tau



neutrino τ


Bosons



photon



gluon



Z^0 W^\pm



Higgs

standard model



THEORETICAL FRAMEWORK

Quantum Mechanics (\hbar) + Special Relativity (c)



Quantum Field Theory


STANDARD THEORY:

- 1) Electricity + Magnetism + Optics (light): γ
Quantum Electrodynamics (QED)
- 2) QED + Weak Interaction: γ, Z, W^\pm
Electroweak Theory $SU(2)_L \otimes U(1)_Y$
- 3) Strong Interaction: 8 Gluons
Quantum Chromodynamics (QCD)

OPEN QUESTIONS:

- The Higgs Boson (Mass scales)
- Gran Unification (Electroweak + Strong)
- SuperSymmetry
- Gravitation: SuperGravity, Strings, ...

LEPTONS



- Do not have Strong Interactions
- Spin $\frac{1}{2}$
- Seen as Free Particles
- Pointlike ($r < \text{few} \times 10^{-17} \text{ cm}$)


Family Structure:

$$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L, \begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}_L, \begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}_L$$

$m_e = 0.5 \text{ MeV}$	$m_\mu = 106 \text{ MeV}$	$m_\tau = 1777 \text{ MeV}$
$\tau_e > 6 \cdot 10^{24} \text{ y}$	$\tau_\mu = 2 \cdot 10^{-6} \text{ s}$	$\tau_\tau = 3 \cdot 10^{-13} \text{ s}$
$m_{\nu_e} < 2 \text{ eV}$	$m_{\nu_\mu} < 0.2 \text{ MeV}$	$m_{\nu_\tau} < 18 \text{ MeV}$

Why 3 ?

NEUTRAL CURRENTS




- Flavour Conserving $\mu \not\rightarrow e \gamma$; $Z \not\rightarrow e^\mp \mu^\pm$
- $g_\gamma \sim Q_l$ ($Q_e = Q_\mu = Q_\tau$; $Q_\nu = 0$)
- Same γ interaction for both lepton helicities
- NC Universality: $g_{Zee} = g_{Z\mu\mu} = g_{Z\tau\tau} \neq g_{Z\nu\nu}$
- Different Z coupling to l_R and l_L
- Left-handed neutrinos only
- 3 Families with light (nearly massless) neutrinos

HOW MANY NEUTRINOS ?





$\sigma(Z \rightarrow \text{hadrons})$




$$N_{\nu} = \frac{\Gamma(Z \rightarrow \text{invisible})}{\Gamma(Z \rightarrow \nu_i \bar{\nu}_i)_{\text{Th}}}$$
$$= 2.9840 \pm 0.0082$$

$$\Gamma(Z \rightarrow \text{invisible}) \equiv \Gamma(Z \rightarrow \text{all}) - \Gamma(Z \rightarrow \text{visible})$$


The heavier leptons μ and τ are unstable





CHARGED CURRENTS




- Left-handed leptons (Right-handed antileptons)






~~\mathcal{P}~~ and ~~\mathcal{C}~~ in Weak Interactions
 CP still a good symmetry

CHARGED CURRENTS



- Left-handed leptons (Right-handed antileptons)



- Doublet partners:

$$l^- \leftrightarrow \nu_l$$

$$\nu_\mu X \rightarrow \mu^- X' \quad ; \quad \nu_\mu X \not\rightarrow e^- X'$$

- Universal Strength

$$T(l \rightarrow \nu_l l' \bar{\nu}_{l'}) \sim \frac{g_W^2}{M_W^2 - q^2} \xrightarrow{q^2 \ll M_W^2} \frac{g_W^2}{M_W^2} \sim G_F \quad \longrightarrow \quad \Gamma(l \rightarrow \nu_l l' \bar{\nu}_{l'}) \sim G_F^2 m_l^5$$

$$\Gamma(\tau \rightarrow \nu_\tau e \bar{\nu}_e) / \Gamma(\mu \rightarrow \nu_\mu e \bar{\nu}_e) \approx (m_\tau / m_\mu)^5$$

LEPTON FLAVOUR VIOLATION

90% CL Upper Limits on $\text{Br}(l^- \rightarrow X^-)$

[BABAR / BELLE]

Decay	U.L.	Decay	U.L.	Decay	U.L.
$\mu^- \rightarrow e^- \gamma$	$1.2 \cdot 10^{-11}$	$\mu^- \rightarrow e^- e^+ e^-$	$1.0 \cdot 10^{-12}$	$\mu^- \rightarrow e^- \gamma \gamma$	$7.2 \cdot 10^{-11}$
$\tau^- \rightarrow e^- \gamma$	$1.1 \cdot 10^{-7}$	$\tau^- \rightarrow e^- e^+ e^-$	$3.6 \cdot 10^{-8}$	$\tau^- \rightarrow e^- e^+ \mu^-$	$2.7 \cdot 10^{-8}$
$\tau^- \rightarrow \mu^- \gamma$	$4.5 \cdot 10^{-8}$	$\tau^- \rightarrow e^- \mu^+ \mu^-$	$3.7 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- e^+ \mu^-$	$2.3 \cdot 10^{-8}$
$\tau^- \rightarrow e^- e^- \mu^+$	$2.0 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- \mu^+ \mu^-$	$3.2 \cdot 10^{-8}$	$\tau^- \rightarrow e^- \pi^0$	$8.0 \cdot 10^{-8}$
$\tau^- \rightarrow \mu^- \pi^0$	$1.1 \cdot 10^{-7}$	$\tau^- \rightarrow e^- \eta'$	$1.6 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^- \eta'$	$1.3 \cdot 10^{-7}$
$\tau^- \rightarrow e^- \eta$	$9.2 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- \eta$	$6.5 \cdot 10^{-8}$	$\tau^- \rightarrow e^- K^{*0}$	$7.8 \cdot 10^{-8}$
$\tau^- \rightarrow e^- K_S$	$5.6 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- K_S$	$4.9 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- \rho^0$	$6.8 \cdot 10^{-8}$
$\tau^- \rightarrow e^- K^+ K^-$	$1.4 \cdot 10^{-7}$	$\tau^- \rightarrow e^- K^+ \pi^-$	$1.6 \cdot 10^{-7}$	$\tau^- \rightarrow e^- \pi^+ K^-$	$3.2 \cdot 10^{-7}$
$\tau^- \rightarrow \mu^- K^+ K^-$	$2.5 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^- K^+ \pi^-$	$3.2 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^- \pi^+ K^-$	$2.6 \cdot 10^{-7}$
$\tau^- \rightarrow e^- \pi^+ \pi^-$	$1.2 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^- \pi^+ \pi^-$	$2.9 \cdot 10^{-7}$	$\tau^- \rightarrow \Lambda \pi^-$	$7.2 \cdot 10^{-8}$
$\tau^- \rightarrow e^+ K^- K^-$	$1.5 \cdot 10^{-7}$	$\tau^- \rightarrow e^+ K^- \pi^-$	$1.8 \cdot 10^{-7}$	$\tau^- \rightarrow e^+ \pi^- \pi^-$	$2.0 \cdot 10^{-7}$
$\tau^- \rightarrow \mu^- K^{*0}$	$5.9 \cdot 10^{-8}$	$\tau^- \rightarrow e^- \phi$	$7.3 \cdot 10^{-8}$	$\tau^- \rightarrow \mu^- \omega$	$8.9 \cdot 10^{-8}$
$\tau^- \rightarrow \mu^+ K^- K^-$	$4.4 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^+ K^- \pi^-$	$2.2 \cdot 10^{-7}$	$\tau^- \rightarrow \mu^+ \pi^- \pi^-$	$0.7 \cdot 10^{-7}$

NEUTRINOS

- Weakly Interacting Particles
- Among most abundant particles in the Universe
- Each second pass through your body

$\sim 10^{14} \nu_e$ from the SUN




NEUTRINOS

Each second pass through your body

$\sim 10^{14}$ ν_e from the SUN



They also come
from below!




SOLAR NEUTRINO PROBLEM

ν_e Measured $<$ ν_e Predicted

SNO

- **CC:** $\nu_e + d \rightarrow p + p + e^-$
- **ES:** $\nu_x + e^- \rightarrow \nu_x + e^-$
- **NC:** $\nu_x + d \rightarrow p + n + \nu_x$


($x = e, \mu, \tau$)



Neutrino Oscillations

$$\nu_e \rightarrow \nu_{\mu, \tau}$$

Sudbury Neutrino Observatory




ATMOSPHERIC NEUTRINOS

(produced by Cosmic Rays)

$$p X \rightarrow \pi X'$$

$$\pi^- \rightarrow \mu^- \bar{\nu}_\mu \quad ; \quad \mu^- \rightarrow \nu_\mu e^- \bar{\nu}_e$$




$$\frac{N_{\nu_\mu}}{N_{\nu_e}} \approx 2$$

Super-Kamiokande:




$$\frac{N_{\nu_\mu}}{N_{\nu_e}} \sim 1$$

$$\nu_\mu \rightarrow \nu_\tau$$

The Standard Model





Super-Kamiokande




The Standard Model

K2K



CNGS

CERN Neutrinos To Gran Sasso



Quarks



up



down



charm



strange



top



beauty

Leptons



electron



neutrino e



muon



neutrino μ



tau



neutrino τ


Bosons



photon



gluon



Z^0 W^\pm



Higgs