

CERN-FNAL HCP 2009

# Experimental Aspects of Heavy Flavour Physics

.... The saga of the penguin and the polar bear....


Valerie Gibson



# Disclaimer....



The title “Experimental Aspects of Heavy Flavour Physics” covers an enormous range of topics. Therefore I can only present a very selective personal view. Concentrate on mostly experimental aspects of



# Acknowledgements




I have taken my inspiration from many recent results and conference talks....If you would like to know more about the current status of Heavy Flavour Physics, have a look at...

Recontres de Moriond EW/QCD 2009

<http://www.slac.stanford.edu/xorg/hfag/>

Many thanks to those who (un)knowingly help me:  
 T.Browder, L.Esteve, T.Gershon,  
 G.Hamel de Monchenault,  
 U.Kerzel, T.Ruf, Y.Sakai, K.Trabelsi, G.Wilkinson  
 and many more....



*Flavor Physics & CP Violation 2009*  
**FPCP 2009**  
 May 27– June 1, Lake Placid, NY

*To integrate past results with recent developments in flavor physics and CP violation, both in theory and experiment. Among the topics included will be: Bottom, Charm and Kaon mixing and decays, determination of the CKM matrix elements, neutrino masses and mixings, new spectroscopy, and future facilities.*

**International Advisory Committee**

- J. A. Appel (FNAL), G. Bonneaud (Ecole Polytechnique), G. Borissov (Lancaster), R. Briere (CMU), T. Browder (Hawaii), K.T. Chao (Beijing), H.Y. Cheng (Accademia Sincal), U. Eggede (Imperial), H. Evans (Indiana), F. Gilman (CMU), M. Giorgi (Pisa), B. Grinstein (San Diego), M. Gronau (Technion), M. Hazumi (KEK), D. Hillin (Cal Tech), C. Jarlskog (Lund), C. S. Kim (Yonsei), J. Kroll (Penn), H. Lacker (Humbolt), Z. Ligeti (LBNL), K. Pitts (Illinois), J. Rosner (Chicago), T. Sanda (Kanagawa), M.D. Shapiro (LBNL), S. Stone (Syracuse—Chair)

**Local Organizing Committee**  
 (Syracuse University)

- Sheldon Stone, Chair
- Marina Artuso
- Steven Blusk
- Ray Mountain
- Tomasz Skwarnicki
- Jianchun Wang

<http://fpcp2009.syr.edu>  
 Questions: [fpcp2009@physics.syr.edu](mailto:fpcp2009@physics.syr.edu)



- Introduction
- The Standard Model
- B Physics
- Celebrating the B factories

---

- What have we learnt from the Tevatron ?
- The LHC era
- and beyond.....
- Summary

Lecture 1

Lecture 2

# Role of Heavy Flavour Physics



Heavy flavour physics has led the way to

- The 3 generation Standard Model
- The CKM picture of flavour
- CP Violation



2008



SM cannot be ultimate theory

- low-energy effective theory of a more fundamental theory at a higher energy scale (TeV range)
- **Hierarchy problem:** New Physics required to cancel radiative corrections to the Higgs mass but leave the SM EW predictions unaffected

NP needs to have a special flavour structure

- provide the suppression mechanism for FCNC processes already observed.
- we need to measure the flavour structure to distinguish between the NP models.

Flavour physics goes hand-in-hand with direct searches

# Historical Note




## New Physics in box diagrams

1970's

Glashow, Iliopoulos and Maini : existence of  $c$  quark (GIM mechanism)

Gaillard, Lee and Rosner :  $m_c \sim 1.5$  GeV from  $K^0 - \bar{K}^0$  mixing




1974  $c$  quark discovered

## New Physics in interference

1980's

$$e^+ e^- \rightarrow \mu^+ \mu^-$$





Existence of  $Z^0$  at high  $\sqrt{s}$

## New Physics in loops

1990's

Virtual effects at the  $Z^0$   
LEP

$$m_t = 170 \pm 10_{-19}^{+17} \text{ GeV}$$



# The Standard Model



Physical quark states in the Standard Model

$$\begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L, \dots, u_R, d_R, c_R, s_R, t_R, b_R$$

Lagrangian for charged current weak decays

$$L_{cc} = -\frac{g}{\sqrt{2}} J_{cc}^\mu W_\mu^* + h.c.$$

where

$$J_{cc}^\mu = (\bar{u}, \bar{c}, \bar{t})_L \gamma^\mu V_{CKM} \begin{pmatrix} d \\ s \\ b \end{pmatrix}_L$$

# CKM Matrix



$V_{\text{CKM}}$  describes rotation between the weak eigenstates ( $d', s', b'$ ) and mass eigenstates ( $d, s, b$ )

weak states


CKM matrix

mass states

Quarks


$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

$V_{ij}$  proportional to transition amplitude from quark  $j$  to quark  $i$



Antiquarks

$$\begin{pmatrix} \bar{d}' \\ \bar{s}' \\ \bar{b}' \end{pmatrix} = \begin{pmatrix} V_{ud}^* & V_{us}^* & V_{ub}^* \\ V_{cd}^* & V_{cs}^* & V_{cb}^* \\ V_{td}^* & V_{ts}^* & V_{tb}^* \end{pmatrix} \begin{pmatrix} \bar{d} \\ \bar{s} \\ \bar{b} \end{pmatrix}$$



CPV due to complex phases of CKM matrix elements



# CKM Matrix



- CKM matrix is complex and unitary
- 4 independent parameters
  - These 4 numbers are **fundamental constants** of nature and **must be determined from experiment**
- Standard parametrization (PDG)
  - 3 angles  $(\theta_{12}, \theta_{23}, \theta_{13})$  1 phase  $\delta$

$$\hat{V}_{CKM}^+ \hat{V}_{CKM} = 1$$

$$V_{CKM} = R_{23} \times R_{13} \times R_{12}$$

$$R_{12} = \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \quad R_{13} = \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix}$$

$$s_{ij} = \sin \theta_{ij} \quad c_{ij} = \cos \theta_{ij}$$

# Wolfenstein Parameterization




Wolfenstein parameterization (perturbative form)

$$\lambda = s_{12} \quad A = \frac{s_{23}}{s_{12}^2} \quad \rho = \frac{s_{13} \cos \delta}{s_{12} s_{23}} \quad \eta = \frac{s_{13} \sin \delta}{s_{12} s_{23}}$$

$$\lambda = \sin \theta_{12} \approx 0.23$$

Reflects hierarchy of strengths of quark transitions



# Wolfenstein Parameterization



Wolfenstein parameterization to  $O(\lambda^3)$ :

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4)$$

Next-to leading order corrections in  $\lambda$  will be important in LHC era:

$$V_{CKM} = \begin{pmatrix} 1 - \frac{\lambda^2}{2} - \frac{\lambda^4}{8} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda + A^2\lambda^5\left(\frac{1}{2} - \rho - i\eta\right) & 1 - \frac{\lambda^2}{2} - \frac{\lambda^4}{8}(1 + 4A^2) & A\lambda^2 \\ A\lambda^3(1 - \bar{\rho} - i\bar{\eta}) & -A\lambda^2 + A\lambda^4(1/2 - \rho - i\eta) & 1 - A^2\lambda^4/2 \end{pmatrix} + O(\lambda^6)$$

$$(\bar{\rho}, \bar{\eta}) \equiv (1 - \lambda^2/2)(\rho, \eta)$$



Requirements for CP violation

$$\left(m_t^2 - m_c^2\right) \left(m_t^2 - m_u^2\right) \left(m_c^2 - m_u^2\right) \\ \times \left(m_b^2 - m_s^2\right) \left(m_b^2 - m_d^2\right) \left(m_s^2 - m_d^2\right) \times J_{CP} \neq 0$$



$J_{CP}$  Jarlskog determinant

where

$$J_{CP} = \left| \text{Im} \left\{ V_{i\alpha} V_{j\beta} V_{i\beta}^* V_{j\alpha}^* \right\} \right| \quad (i \neq j, \alpha \neq \beta)$$

Using parameterizations

$$J_{CP} = s_{12} s_{13} s_{23} c_{12} c_{23} c_{13} \sin \delta = \lambda^6 A^2 \eta = O(10^{-5})$$

**→** CP violation is small in the Standard Model

# Unitarity Triangles



CKM matrix is unitary : 12 conditions (6 normalisation, 6 orthogonality)

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0 \quad (\text{db})$$







$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0 \quad (\text{sb})$$

$$V_{ud}V_{us}^* + V_{cd}V_{cs}^* + V_{td}V_{ts}^* = 0 \quad (\text{ds})$$

$$V_{ud}V_{td}^* + V_{us}V_{ts}^* + V_{ub}V_{tb}^* = 0 \quad (\text{ut})$$

$$V_{cd}V_{td}^* + V_{cs}V_{ts}^* + V_{cb}V_{tb}^* = 0 \quad (\text{ct})$$

$$V_{ud}V_{cd}^* + V_{us}V_{cs}^* + V_{ub}V_{cb}^* = 0 \quad (\text{uc})$$




All 6  $\Delta$ 's have the same area ( $= J_{CP}/2$ ), a measure of CPV in the Standard Model.

# The Unitarity Triangle



Redraw “unsquashed”  $\Delta$ 's and take  $V_{cb}^* V_{cd}$  real divide by  $|V_{cb}^* V_{cd}|$



$$\alpha \equiv \pi - \beta - \gamma$$

$$\gamma \equiv \arg \left[ -\frac{V_{ub}^* V_{ud}}{V_{cb}^* V_{cd}} \right] = \tan^{-1} \frac{\eta}{\rho} \sim 70^\circ$$

$$\beta \equiv \arg \left[ -\frac{V_{cb}^* V_{cd}}{V_{tb}^* V_{td}} \right] = \tan^{-1} \frac{\bar{\eta}}{1 - \rho} \sim 21^\circ$$



# The Ultimate Quest...



# To discover New Physics

# beyond the Standard Model



# The Quest...




NP models introduce new particles which could

- be produced and discovered as real particles
- appear as virtual particles in loop processes → observable deviations from the SM expectations in flavour physics and CPV

## Heavy flavour programme

- Precision measurements of CKM elements
- Compare tree level processes with loop processes sensitive to NP
- Measure all angles and sides in many different ways and look for inconsistencies
- Measure processes very suppressed in SM






# B Physics



# Why the b-quark ?




- Heaviest quark that forms hadronic bound states ( $m \sim 4.7$  GeV)
- Must decay outside 3<sup>rd</sup> family
  - All decays are CKM suppressed
  - Long lifetime ( $\sim 1.6$  ps)
- High mass: many accessible final states (all Br's are small)




- Dominant decay process: “tree”  $b \rightarrow c$  transition
- Very suppressed “tree”  $b \rightarrow u$  transition
- FCNC: “penguin”  $b \rightarrow s, d$  transition
- Flavour oscillations ( $b \rightarrow t$  “box” diagram)
- CP violation – expect large CP asymmetries in some B decays

# The Birth of B Physics




1977

Discovery of b quark  
Fermilab fixed target




1980

Discovery of  $\Upsilon(4s)$   
& B meson  
Cornell CUSB




1983

B meson lifetime  
MAC-Mark II  
 $\tau \sim 490 \mu\text{s}$



1987

Exclusive reconstruction of  
several  $b \rightarrow c$  decay modes  
CLEO



1987

B mixing observed  
ARGUS

# B Physics



PDG 1986

BOTTOM MESONS <sup>a</sup>		$[B^+ = u\bar{b}, B^0 = d\bar{b}, \bar{B}^0 = \bar{d}b, B^- = \bar{u}b]$	
$B^\pm$	$\frac{1}{2}(0^-) \rho$	5271.2	
		$\pm 3.0$	
	$m_{B^0} - m_{B^\pm} = 4.0$		
		$\pm 3.4$	
$B^0, \bar{B}^0$	$\frac{1}{2}(0^-) \rho$	5275.2	
		$\pm 2.8$	
		$B^+ \bar{\phantom{B}}^-$ (or $B^- \rightarrow$ chg. conj.)	
		$\bar{D}^0 \pi^+$	( 1.1 $\pm$ 0.6 )% 2303
		$D^*(2010)^- \pi^+ \pi^+$	( 2.7 $\pm$ 1.7 )% 2243
		$J/\psi(3097)K^+$	( < 2.6 ) $\times 10^{-3}$ 1678
		$\rho^0 \pi^+$	( < 6 ) $\times 10^{-4}$ 2578
		$B^0 \bar{\phantom{B}}^0$ (or $\bar{B}^0 \rightarrow$ chg. conj.)	
		$\bar{D}^0 \pi^+ \pi^-$	( 7 $\pm$ 5 )% 2299
		$D^*(2010)^- \pi^+$	( 1.7 $\pm$ 0.7 )% 2253
		$D^*(2010)^- \rho^+$	( 8 $\pm$ 7 )% 2180
		$J/\psi(3097)K^+ \pi^-$	( < 6.3 ) $\times 10^{-3}$ 1649
		$\pi^+ \pi^-$	( < 5 ) $\times 10^{-4}$ 2634
		$e^+ e^-$	( < 3 ) $\times 10^{-4}$ FC 2638
		$\mu^+ \mu^-$	( < 2 ) $\times 10^{-4}$ FC 2635
		$e^+ \mu^- + c.c.$	( < 3 ) $\times 10^{-4}$ LF 2637
$B^\pm, B^0, \bar{B}^0$		$(14.2 \pm 2.7) \times 10^{-13}$	$e^\pm \nu$ hadrons ( 12.3 $\pm$ 0.8 )%
	(not separated) <sup>q</sup>	$c\tau = 0.043$	$\mu^\pm \nu$ hadrons ( 11.0 $\pm$ 0.9 )%
	$\frac{\Gamma(B \rightarrow \bar{B} \rightarrow \ell^- \text{ any})}{\Gamma(B \rightarrow \ell^\pm \text{ any})} < 0.12$		$D^0$ anything ( 80 $\pm$ 28 )%
			$K$ anything ( seen )
			$p$ anything ( > 3.6 )%
			$\Lambda$ anything ( > 2.2 )%
			$e^+ e^-$ anything ( < 0.6 )%
			$\mu^+ \mu^-$ anything ( < 0.6 )%
			$J/\psi(3097)$ anything ( 1.2 $\pm$ 0.3 )%
			$D^*(2010)^\pm$ anything ( 23 $\pm$ 9 )%

PDG 2009

> 25 pages

- 1992 Evidence for  $\Lambda_b, B_s$
- 1993 First observation of  $B \rightarrow K^* \gamma, B \rightarrow \pi^+ \pi^-,$  time dependent  $B$  mixing
- 1994 Evidence for  $\Xi_b, B^{**},$  measurement of exclusive B lifetime
- 1998 Discovery of  $B_c$
- 2001 Discovery of CPV in B system
- 2004 Direct CPV in B system
- 2006 Measurement of  $B_s$  mixing

# B Mixing



Mixing of neutral B mesons governed by

$$i \frac{\partial}{\partial t} \begin{pmatrix} a \\ b \end{pmatrix} = H \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} M_{11} - \frac{i}{2} \Gamma_{11} & M_{12} - \frac{i}{2} \Gamma_{12} \\ M_{12}^* - \frac{i}{2} \Gamma_{12}^* & M_{22} - \frac{i}{2} \Gamma_{22} \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}$$

Physical mass eigenstates

$$|B_{L,H}\rangle = p |B^0\rangle \pm q |\bar{B}^0\rangle$$



$$\frac{q}{p} = \sqrt{\frac{M_{12}^* - \frac{i}{2} \Gamma_{12}^*}{M_{12} - \frac{i}{2} \Gamma_{12}}} \quad |p|^2 + |q|^2 = 1$$

– p and q represent the amount of state mixing

$$\Delta m = m_H - m_L = 2|M_{12}|$$

$$\Delta \Gamma = \Gamma_L - \Gamma_H = 2|\Gamma_{12}|$$

$$|q/p| = 1$$



# CP Violation in B System



Decay amplitudes of flavour states  $\rightarrow$  final state  $f$

$$A_f = \langle f | H | B^0 \rangle \quad \bar{A}_f = \langle f | H | \bar{B}^0 \rangle$$

Define

$$\lambda_f = \frac{q}{p} \frac{\bar{A}_f}{A_f}$$

General time dependence of decay rate for initially pure flavour states

$$\Gamma_f \equiv \left| \langle f | H | B^0(t) \rangle \right|^2 = \frac{1 + |\lambda_f|^2}{2} |A_f|^2 e^{-t/\tau} \left[ \cosh yt/\tau + \Omega_f \sinh yt/\tau + C_f \cos xt/\tau - S_f \sin xt/\tau \right]$$

$$\bar{\Gamma}_f \equiv \left| \langle f | H | \bar{B}^0(t) \rangle \right|^2 = \frac{1 + |\lambda_f|^2}{2} \left| \frac{p}{q} A_f \right|^2 e^{-t/\tau} \left[ \cosh yt/\tau + \Omega_f \sinh yt/\tau - C_f \cos xt/\tau + S_f \sin xt/\tau \right]$$

$$S_f \equiv \frac{2 \operatorname{Im} \lambda_f}{1 + |\lambda_f|^2} \quad C_f \equiv \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2} \quad \Omega_f^2 + S_f^2 + C_f^2 = 1$$

$$\tau \equiv 1/\Gamma$$

$$x \equiv \Delta m/\Gamma$$

$$y \equiv \Delta\Gamma/2\Gamma$$

# 3 Types of CP Violation



CP violation if  $\Gamma_f \neq \bar{\Gamma}_f$

CPV in Decay  
Direct CP Violation

$$\left| \frac{\bar{A}_f}{A_f} \right| \neq 1$$

CPV in Mixing  
Indirect CP Violation

$$\left| \frac{q}{p} \right| \neq 1$$


$$\text{Im}\{\Gamma_{12}^* M_{12}\} \neq 0$$

CPV in Interference between mixing and decay

Indirect CP Violation

$$|\lambda_f| = 1, \quad \text{Im}\{\lambda_f\} \neq 0$$

$$A_f^{CP}(t) = \frac{\Gamma_f(t) - \bar{\Gamma}_f(t)}{\Gamma_f(t) + \bar{\Gamma}_f(t)} = \frac{-C_f \cos(\Delta mt) + S_f \sin(\Delta mt)}{\cosh(\Delta\Gamma t/2) + \Omega_f \sinh(\Delta\Gamma t/2)}$$



Golden case: CP final state and single dominating amplitude



$$A_{f_{CP}}^{CP}(t) = \text{Im} \lambda_{f_{CP}} \sin(\Delta mt)$$






# B Decays






## Dominant decays



## Rare hadronic decays



## Radiative and leptonic decays




# Where to start ?



$\Upsilon$  (4s)



# $\Upsilon(4s)$ Resonance




Cleanest way to produce B mesons  
 $e^+e^-$  collisions around

$$\sqrt{s} = 10.58 \text{ GeV}$$


$$\sigma_{b\bar{b}} = 1.1 \text{ nb} \quad \text{continuum} \sim 3 \text{ nb}$$

$\sim 1.1$  million  $B\bar{B}$  pairs per  $\text{fb}^{-1}$



50% / 50%  
 $B^0\bar{B}^0 / B^+B^-$

$B\bar{B}$  pair is produced in a coherent L=1 state  
Two B mesons evolve until one decays




# $\Upsilon(4S)$ Resonance



Symmetric-energy collider : B mesons produced  $\sim$  at rest in the CM frame which, combined with a short B lifetime ( $\sim 1.5$  ps), makes flight distance unmeasurably small.

Asymmetric-energy collider : with boost  $\beta\gamma \sim 0.6$



$$\Delta t \approx \frac{\Delta z}{\langle \beta\gamma \rangle c}$$

$$\langle |\Delta z| \rangle \approx 200 \mu m$$

# Asymmetric B Factories





## PEP-II @ SLAC


High Energy Ring : 9.0 GeV  $e^-$   
 Low Energy Ring : 3.1 GeV  $e^+$   
 Design luminosity :  $3 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$   
 Peak luminosity :  $1.207 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

## KEK-B @ KEK

High Energy Ring : 8.0 GeV  $e^-$   
 Low Energy Ring : 3.5 GeV  $e^+$   
 Design luminosity :  $1 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$   
 Peak luminosity :  $1.71 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$   
 Beam crossing angle : 22 mrad



# Luminosity at B Factories





K.Trabelsi, Moriond 2009

# BELLE and BABAR



3/4 layers DSSD silicon detector  
 Threshold Cherenkov + TOF  
 Central drift chamber  
 EM Cal Csl  
 RPCs  
 1.5T solenoid



5 layers DSSD silicon detector  
 Ring-Imaging Cherenkov DIRC  
 Drift chamber  
 EM Cal Csl  
 RPCs  
 1.5T SC solenoid





# Kinematics at the $\Upsilon(4s)$



Precise knowledge of CM energy used to constrain kinematics of B system.



$$\Delta E \equiv E_B^* - E_{beam}^*$$

$$m_{ES} \equiv \sqrt{E_{beam}^{*2} - p_B^{*2}} \quad E_{beam}^* = \sqrt{s}/2$$




Estimate background from sidebands

Continuum background suppressed using event shape





# Time-Dependent Analyses






# What to measure first ?






# Sin (2β) ≡ Sin (2φ<sub>1</sub>)



$$\beta = \tan^{-1} \frac{\bar{\eta}}{1 - \bar{\rho}}$$




# $\beta : b \rightarrow ccs$ modes



Golden decay mode:  $B^0 \rightarrow J/\psi K^0$ , final state CP eigenstate

Method:

Count number of signal events reconstructed with  $B^0$  and  $\bar{B}^0$  tags as a function of  $\Delta t$   
 Observed asymmetry depends on time resolution and tagging purity of sample.






$$A_{CP}(t) = \frac{N(B_{tag} = B^0) - N(B_{tag} = \bar{B}^0)}{N(B_{tag} = B^0) + N(B_{tag} = \bar{B}^0)}$$

$$\approx \pm \{ (1 - 2\omega) \times \sin 2\beta \times \sin(\Delta m \Delta t) \} \otimes R(\Delta t)$$

$$\lambda_f = \left( \frac{q}{p} \right)_{B^0} \left( \frac{q}{p} \right)_{K^0} \left( \frac{\bar{A}}{A} \right)_{J/\psi K^0}$$

$$= - \left( \frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \right) \left( \frac{V_{cs} V_{cd}^*}{V_{cs}^* V_{cd}} \right) \left( \frac{V_{cb} V_{cs}^*}{V_{cb}^* V_{cs}} \right)$$

$$\text{Im}\{\lambda_f\} = \text{Im}\{-e^{-2i\beta}\} = \sin 2\beta$$




# $\beta$ : Then...




Year 2000

## First Physics Results



$$\sin 2\beta = +0.45^{+0.43}_{-0.44}$$




$$\sin 2\beta = 0.12 \pm 0.37 \pm 0.09$$

# And now...




Belle : 535M BB; PRL98 (2007) 031802

BaBar : 465M BB; PRD79 (2009) 072009




$$\sin 2\beta = 0.642 \pm 0.031 \pm 0.017$$




$$\sin 2\beta = 0.666 \pm 0.031 \pm 0.013$$



# $\beta : b \rightarrow ccs$ modes



Other measurements sensitive to  $\cos(2\beta)$  remove ambiguity



$$\beta = (21.1 \pm 0.9)^\circ$$

# $\beta_{\text{eff}}$ in other modes




$b \rightarrow ccs$	$b \rightarrow ccd$	$b \rightarrow sqq$
$J/\psi K_s^0, \psi(2s)K_s^0, \chi_{c1}K_s^0,$ $\eta_c K_s^0, J/\psi K_L^0,$ $J/\psi K^{*0} (K^{*0} \rightarrow K_s^0 \pi^0)$	$D^{*+}D^-, D^+D^-,$ $J/\psi \pi^0, D^{*+}D^{*-}$	$\phi K^0, K^+K^-K_s^0,$ $K_s^0 K_s^0 K_s^0, \eta' K^0, K_s^0 \pi^0,$ $\omega K_s^0, f_0(980)K_s^0$

Increasing tree diagram amplitude



Increasing sensitivity to new physics








# $\beta_{\text{eff}} : b \rightarrow sqq$ modes



Most sensitive to new physics



Moriond 2003



2.6  $\sigma$  : Hint of New Physics ?

Belle : PRD67 (2003) 031102


# $\beta_{\text{eff}} : b \rightarrow s q \bar{q}$ modes




Time-dependent Dalitz plot analyses sensitive to  $\sin(2\beta_{\text{eff}})$  and  $\cos(2\beta_{\text{eff}})$





BaBar : 465M BB;  
ArXiv:0808.0700  
Belle : 657M BB;  
ICHEP 08



BaBar : 383M BB;  
ArXiv:0905.3615  
Belle : 657M BB;  
PRD79 (2009) 072004




# $\beta_{\text{eff}} : b \rightarrow sqq$ modes



Non-zero  $\beta_{\text{eff}}$  is observed in  $b \rightarrow sqq$  penguin transition  
Smaller error is required to see any deviation between  $\sin(2\beta_{\text{eff}})$  and  $\sin(2\beta)$




# Angle $\alpha \equiv \phi_2$



$$\alpha = \pi - \beta - \gamma$$

$$\beta = \tan^{-1} \frac{\bar{\eta}}{1 - \bar{\rho}}$$


$$\gamma = \tan^{-1} \frac{\eta}{\rho}$$



# $\alpha : B \rightarrow \pi^+\pi^-$



Strangeless-charmless two-body decay;  $b \rightarrow uud$  transition, final state CP eigenstate.



If tree amplitude dominates

$$\lambda_{\pi^+\pi^-}^t = +e^{-2i(\beta+\gamma)} = e^{2i\alpha}$$

$$S_{\pi^+\pi^-} = \sin 2\alpha \quad C_{\pi^+\pi^-} = 0$$

$$S_f \equiv \frac{2\text{Im}\lambda_f}{1+|\lambda_f|^2} \quad C_f = \frac{1-|\lambda_f|^2}{1+|\lambda_f|^2}$$

... but penguin contributions cannot be neglected

$$\left| \frac{P}{T} \right| \sim 30\%$$

$$\lambda_{\pi^+\pi^-}^{t+p} \approx e^{2i\alpha} \left[ 1 + 2i \left| \frac{P}{T} \right| \sin \alpha e^{i(\delta_P - \delta_T)} \right]$$

$$S_{\pi^+\pi^-} = \sin 2\alpha - 2 \left| \frac{P}{T} \right| \sin \alpha \cos 2\alpha \cos(\delta_P - \delta_T) + O\left(\left| \frac{P}{T} \right|^2\right)$$

In general let  $\lambda_{\pi^+\pi^-} \equiv \left| \lambda_{\pi^+\pi^-} \right| e^{2i\alpha_{eff}}$  and fit time-dependent CP asymmetry

$$A_{CP}(t) = \sqrt{1 - C_{\pi\pi}^2} \sin 2\alpha_{eff} \sin(\Delta mt) - C_{\pi\pi} \cos(\Delta mt)$$

# $\alpha : B \rightarrow \pi^+\pi^-$



Two types of CP violation are observed




- Direct CP violation  $C_{\pi^+\pi^-} \neq 0$
- Mixing induced CP violation  $S_{\pi^+\pi^-} \neq 0$

Need to estimate  $\Delta\alpha \equiv \alpha_{eff} - \alpha$

$$\sigma(\alpha_{eff}) \sim 4^\circ$$

BaBar : 467M BB; ArXiv:0807.4226

Belle : 535M BB; PRL98 (2007) 211801



# Isospin Analysis $B \rightarrow \pi\pi$



Gronau and London, PRL65 (1990) 3381.

Isospin amplitude relations for


$$B^0 \rightarrow \pi^+ \pi^-$$

$$B^0 \rightarrow \pi^0 \pi^0$$

$$B^+ \rightarrow \pi^+ \pi^0$$

$$A_{+-} + \sqrt{2}A_{00} = \sqrt{2}A_{0+}$$


$$\bar{A}_{+-} + \sqrt{2}\bar{A}_{00} = \sqrt{2}\bar{A}_{0+}$$



Neglecting EWP

$$|A_{0+}| = |\bar{A}_{0+}|$$

Ambiguities: 4 triangle orientations  $\rightarrow$  4-fold ambiguity for  $\Delta\alpha$   
 $\alpha \Leftrightarrow (\pi - \alpha) \Rightarrow$  8-fold ambiguity for  $\alpha$



Need to measure  $Br(B^0 \rightarrow \pi^0 \pi^0)$  and  $Br(\bar{B}^0 \rightarrow \pi^0 \pi^0)$

$\sigma(\Delta\alpha)$  determines  $\sigma(\alpha)$

# Isospin Analysis $B \rightarrow \pi\pi$





6 inputs :  $Br(B^0 \rightarrow \pi^+\pi^-)$ ,  $Br(B^+ \rightarrow \pi^+\pi^0)$ ,  $Br(B^0 \rightarrow \pi^0\pi^0)$ ,  $C_{+-}$ ,  $S_{+-}$ ,  $C_{00}$

$\alpha$  scan : find minimum  $\chi^2$  in fit of isospin triangle to the measurements

convert to C.L. (frequentist)

BaBar : 467M BB; ArXiv:0807.4226

Belle : 535M BB; PRL98 (2007) 211801



Belle:	$11 < \alpha < 79^\circ$	excluded at 95% C.L.
BaBar :	$23 < \alpha < 67^\circ$	excluded at 90% C.L.



# $\alpha : B \rightarrow \rho\rho$



$B \rightarrow \rho\rho$  similar to  $B \rightarrow \pi\pi$

😊  $Br(B^0 \rightarrow \rho^+\rho^-) \sim 5 \times Br(B^0 \rightarrow \pi^+\pi^-)$  and  $Br(B^0 \rightarrow \rho^0\rho^0) \sim 4\% \times Br(B^0 \rightarrow \rho^+\rho^-)$


😊  $|P/T|$  smaller  $\sim 4\%$

😊 Time-dependent measurement for  $B^0 \rightarrow \rho^0\rho^0$  possible  $\rightarrow S_{00}$

lifts 4-fold ambiguity

😞  $\rho$  has spin 1  $\rightarrow$  3 polarization states  
Mixture CP odd and CP even

1 longitudinal  $\propto \cos\theta_1 \cos\theta_2$   
2 transverse  $\propto \sin\theta_1 \sin\theta_2 e^{\pm i\phi}$



Integrate over  $\phi$

$$\frac{1}{\Gamma} \frac{d^2\Gamma}{d\cos\theta_1 d\cos\theta_2} = \frac{9}{16} \left[ 4 \cos^2\theta_1 \cos^2\theta_2 f_L + \sin^2\theta_1 \sin^2\theta_2 (1 - f_L) \right]$$

😊 is almost longitudinally polarized

$$f_L(\rho^+\rho^-) = 0.992 \pm 0.024^{+0.026}_{-0.013}$$




# $\alpha : B \rightarrow \rho\rho$



BaBar : 387M BB; PRD76 (2007) 052007

Belle : 535M BB; PRD76 (2007) 011104





$\sigma(\alpha_{eff}) \sim 6^\circ$

# $\alpha : B \rightarrow \rho\rho$



$B^0 \rightarrow \rho^0 \rho^0$

$B^+ \rightarrow \rho^+ \rho^0$



$Br(B \rightarrow \rho^0 \rho^0) = (0.92 \pm 0.32 \pm 0.14) \times 10^{-6}$   
 $f_L = 0.75^{+0.11}_{-0.14} \pm 0.04$   
 $S_L^{00} = 0.3 \pm 0.7 \pm 0.2$       $C_L^{00} = 0.2 \pm 0.8 \pm 0.3$

$Br(B \rightarrow \rho^0 \rho^0) < 1.0 \times 10^{-6}$  @ 90% c.l.

$Br(B^+ \rightarrow \rho^+ \rho^0) = (23.7 \pm 1.4 \pm 1.4) \times 10^{-6}$   
 $f_L = 0.950 \pm 0.015 \pm 0.006$

New: Moriond 2009


BaBar : 465M BB; PRD78 (2008) 071104  
Belle : 657M BB; PRD78 (2008) 111102

BaBar : 465M BB; PRL102 (2009) 141802

# B → ρρ



Perform isospin analysis:





$$\alpha = \left(92.4^{+6.0}_{-6.5}\right)^{\circ}$$




Why the significant improvement?  
Increase of  $Br(B^+ \rightarrow \rho^+ \rho^0)$  by  $\sim 2\sigma$   
flattens isospin triangle

Two distinct solutions → single solution



# Summary $\alpha$




$$\alpha = \left(89.0^{+4.4}_{-4.2}\right)^{\circ} \text{ 60\% c.l. interval}$$


$\alpha$  is now a precision 4.8% measurement  
 Note:  $\beta$  @ 4.2%



# Angle $\gamma \equiv \phi_3$



$$\gamma = \tan^{-1} \frac{\eta}{\rho}$$




# $\gamma : B \rightarrow DK$



The measurement of  $\gamma$  via tree processes provides a SM benchmark that must be met by any New Physics model

The theoretically cleanest method measures  $\gamma$  via the interference between  $B \rightarrow D^0 K$  and  $B \rightarrow \bar{D}^0 K$



Common parameters:

CKM angle  $\gamma$

Amplitude ratio,  $r_B$

Strong phase difference,  $\delta_B$

$\gamma$  precision very sensitive to value of  $r_B$

$$\frac{\langle B \rightarrow \bar{D}^0 K \rangle}{\langle B \rightarrow D^0 K \rangle} = r_B e^{i(\delta_B - \gamma)}$$

$$r_B \sim \frac{|V_{ub} V_{cs}^*|}{|V_{cb} V_{us}^*|} \times |\text{col. supp}| = 0.1 - 0.2$$

# $\gamma : B \rightarrow DK$



Reconstruct  $D$  in final states accessible to both  $D^0$  and  $\overline{D}^0$

**“ADS” Method:** Atwood, Dunietz, Soni; PRL78 (1997) 3257; PRD63 (2001) 036005.

$D \rightarrow$  Cabibbo favoured and doubly-suppressed decays e.g.  $K^+\pi^-$ ,  $K^+\pi^-\pi^0$

**“GLW” Method:** Gronau, London, Wyler; PLB253 (1991) 483; PLB265 (1991) 172.

$D \rightarrow$  CP Eigenstates e.g.  $K^+K^-$ ,  $\pi^+\pi^-$ ,  $K_S\pi^0$

**“Dalitz” or “GGSZ” Method:** Giri, Grossman, Soffer, Zupan, PRD68 054018 (2003).

$D \rightarrow$  three-body decays e.g.  $K_S\pi^+\pi^-$ ,  $K_S K^+K^-$

Time-integrated analyses, tagging not required

Effects due to charm mixing and CP violation negligible

Different B decays (e.g.  $DK$ ,  $D^*K$ ,  $DK^*$ ) have different hadronic factors  $r_B$ ,  $\delta_B$

Strategy: combine as many channels as possible to improve  $\gamma$  sensitivity




# GGSZ (Dalitz) Method



Currently most powerful method for extraction of  $\gamma$

Exploit interference pattern in Dalitz plot for  $B^\mp \rightarrow D(K_s h^+ h^-) K^\mp$




Sensitivity varies strongly over Dalitz plane (mixture ADS+GLW)





Input knowledge of  $D$  decay amplitude, introduces model uncertainty

Simultaneous fit to Dalitz plot density for  $B^+/B^-$  data


# GGSZ (Dalitz) Method



BaBar : 383M BB; PRD78 (2008) 034023  
Belle : 657M BB; ArXiv:0803.3375



# GGSZ (Dalitz) Method



BaBar : 383M BB; PRD78 (2008) 034023

Belle : 657M BB; ArXiv:0803.3375

$$x_{\pm} = r_B \cos(\delta_B \pm \gamma)$$

$$y_{\pm} = r_B \sin(\delta_B \pm \gamma)$$

$$\gamma = (76_{-13}^{+12} \pm 4 \pm 9)^{\circ}$$

$$B \rightarrow DK, D^*K$$

$$D \rightarrow K_s \pi^+ \pi^-$$



model error

$$\gamma = (76 \pm 22 \pm 5 \pm 5)^{\circ}$$

$$B \rightarrow DK, D^*K, DK^*$$

$$D \rightarrow K_s \pi^+ \pi^-, K_s K^+ K^-$$




Note: difference in Belle & BaBar stat errors due to values of  $r_B$

# Summary $\gamma$



World average combinations vary according to statistics philosophy...


frequentist




$$\gamma = (70^{+27}_{-29})^\circ$$

$$r_B = 0.103^{+0.017}_{-0.023}$$

Bayesian




$$\gamma = (78 \pm 12)^\circ$$



# Summary of the Angles



$$\alpha = (89.0^{+4.4}_{-4.2})^\circ \text{ 60\% c.l. interval}$$




$$\gamma = (70^{+27}_{-29})^\circ$$


$$\beta = (21.1 \pm 0.9)^\circ$$



# $|V_{ub} / V_{cb}|$



$$\left| \frac{V_{ub}}{V_{cb}} \right| = \frac{\lambda}{1 - \lambda^2/2} \sqrt{\bar{\rho}^2 + \bar{\eta}^2}$$



# $V_{ub}$ and $V_{cb}$



$|V_{ub}|$  and  $|V_{cb}|$  determined from semi-leptonic B decays

At tree level everything is clean

QCD corrections must be included


- Inclusive measurements :  
OPE  $\rightarrow$  total s.l. decay rate, moments
- Exclusive measurement :  
Form factors from LQCD

$V_{ub}$  suffers from large  $b \rightarrow c$  background

$$\frac{\Gamma(b \rightarrow ul\nu)}{\Gamma(b \rightarrow cl\nu)} \sim \left| \frac{V_{ub}}{V_{cb}} \right|^2 \sim \frac{1}{50}$$

Theory and experiment analyses are well advanced


- Briefly comment on current status






**Inclusive**

Global fit to moments of  $E_\ell$ ,  $M_X^2$  in  $b \rightarrow c\ell\nu$  and  $E_\gamma$  in  $b \rightarrow s\gamma$  decays



Belle : PRD78 (2008) 032016





HFAG: Inclusion of  $b \rightarrow s\gamma$  lowers  $m_b$  contrary to Belle moment analysis

$|V_{cb}| = (41.67 \pm 0.43 \pm 0.08 \pm 0.58) \times 10^{-3}$



**2.5 $\sigma$**

**Exclusive**

Select  $D^{(*)}\ell\nu$  decays on the recoil of a fully reconstructed  $B$



BaBar : ArXiv:0807.4978



$|V_{cb}| = (39.1 \pm 1.4 \pm 0.9) \times 10^{-3} \quad (D)$

$|V_{cb}| = (38.3 \pm 0.5 \pm 1.0) \times 10^{-3} \quad (D^*)$



# $V_{ub}$




## Inclusive

Experimental measurements of partial rates in regions of  $M_X$ ,  $q^2$ ,  $P_+$ ...



### Recoil analysis

PRL100 (2008) 171802



### Multivariate analysis measures full $b \rightarrow ul\nu$ rate

FPCP09






$$\frac{\Delta|V_{ub}|}{|V_{ub}|} \sim 4\% (exp)$$

$$\sim 6-7\% (th)$$

## Exclusive

$B \rightarrow \pi l \nu$ : reconstruct  $\nu$  from whole event and extract signal yield in  $q^2$  bins from fit to  $\Delta E$  and  $M_{ES}$  distributions



$$\frac{\Delta|V_{ub}|}{|V_{ub}|} \sim 5\% (exp)$$

$$\sim +17\% (th)$$


$$\sim -11\% (th)$$

Lower than inclusive  $V_{ub}$

# B → τν



The helicity suppressed B → τν annihilation decay sensitive to  $f_B |V_{ub}|$   
 Also sensitive to tree-level charged Higgs



$$Br(B^+ \rightarrow \tau^+ \nu) = \frac{G_F^2}{8\pi} |V_{ub}|^2 f_B^2 m_B m_\tau^2 \tau_B \left(1 - \frac{m_\tau^2}{m_B^2}\right)^2$$

An excess of events is clearly visible in signal region of “Extra Energy in Calorimeter”




$$Br(B^+ \rightarrow \tau^+ \nu) = \left(1.65^{+0.38+0.35}_{-0.37-0.37}\right) \times 10^{-4}$$

Semi-leptonic tag



$$Br(B^+ \rightarrow \tau^+ \nu) = (1.2 \pm 0.4 \pm 0.3 \pm 0.2) \times 10^{-4}$$

hadronic tag



BaBar : 383M BB; PRD77 (2008) 011107

Belle : 657M BB; ArXiv:0809.3834



# B → τν



Within SM,  $\text{Br}(B \rightarrow \tau\nu)$  determines  $f_B |V_{ub}|$

- Input  $f_B$  from lattice →  $|V_{ub}|$

~2.5σ discrepancy between  $\text{Br}(B \rightarrow \tau\nu)$  and CKM from other measurements



Combining with B mixing results removes dependence on  $f_B$

$$\frac{\text{Br}(B^+ \rightarrow \tau^+ \nu)}{\Delta m_d} = \frac{3\pi}{4} \frac{m_\tau^2 \tau_B}{m_W^2 S(x_t)} \left(1 - \frac{m_\tau^2}{m_B^2}\right)^2 \frac{\sin^2 \beta}{\sin^2 \gamma} \frac{1}{|V_{ud}|^2 B_{B_d}}$$


Tension with  $\sin 2\beta$  persists.

Theory free prediction for  $B_{B_d} \sim 2.7\sigma$  from LQCD value

# B → τν




Tree-level charged Higgs contribution interferes destructively with SM W diagram



e.g. MSSM (G.Isidori; ArXiv:0710.5377)

$$Br(B^+ \rightarrow \tau^+ \nu) \approx Br^{SM} \times \left( 1 - \frac{m_B^2}{m_H^2} \tan^2 \beta \right)^2$$

$$\frac{Br(B^+ \rightarrow \tau^+ \nu)}{Br^{SM}} = 1.77 \pm 0.65$$




Large  $Br/Br^{SM}$  ruled out by Br upper limits


# $B \rightarrow \tau \nu$



## B factories versus LHC (ATLAS) for the charged Higgs




Current flavour constraints are already competitive with LHC direct search sensitivity for charged Higgs




U.Haisch, hep-ph/0805.2141; ATLAS curve added by Steve Robertson (LLWI 2009)  
D.Eriksson, F.Mahmoudi and O.Stal, JHEP 0811:0.35 (2008) for MSSM interpretation



$$\left| V_{td} / V_{ts} \right|$$




$$\left| \frac{V_{td}}{V_{ts}} \right| \approx \sqrt{(1 - \bar{\rho})^2 + \bar{\eta}^2}$$



# B Mixing



Traditional method for measuring  $V_{td}$  and  $V_{ts}$  is through box diagrams



SM

$$\Delta m_q = 2|M_{12}| = \frac{G_F^2 m_W^2 \eta_{B_q} m_{B_q} B_{B_q} f_{B_q}^2}{6\pi^2} S(m_t^2/m_W^2) |V_{tq}^* V_{tb}|^2 \quad |q/p| = 1$$

Measure oscillation rates for flavour eigenstates


Input theory from LQCD for  $B_{B_q}$  and  $f_{B_q}$  to extract  $|V_{td}|$  or  $|V_{td}/V_{ts}|$

HPQCD; ArXiv:0902.1815

$$f_{B_d} \sqrt{B_{B_d}} = 216(15) \text{ MeV}$$

$$f_{B_s} \sqrt{B_{B_s}} = 266(18) \text{ MeV}$$



$$\xi = \frac{f_{B_s} \sqrt{B_{B_s}}}{f_{B_d} \sqrt{B_{B_d}}} = 1.258(33)$$




$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{B_s}}{m_{B_d}} \xi^2 \left| \frac{V_{ts}}{V_{td}} \right|^2$$

3%

# B Mixing



$$A_{mix}(\Delta t) \equiv \left\{ (1 - 2\omega) \times \cos \Delta m \Delta t \right\} \otimes R(\Delta t)$$



$$\left| \frac{V_{ts}}{V_{td}} \right| = 0.214 (1)_{\text{exp.}} (5)_{\text{lattice.}}$$



FPCP 09:  
R.Van de Water




# FCNC : $B \rightarrow (\rho, \omega)\gamma, K^*\gamma$




Access to  $|V_{td}/V_{ts}|$  through ratios of exclusive Br's.  
 Provides comparison between penguins vs box  
 Many recent and accurate exclusive measurements



$$\frac{Br(B \rightarrow (\rho/\omega)\gamma)}{Br(B \rightarrow K^*\gamma)} = \left| \frac{V_{td}}{V_{ts}} \right|^2 \left( \frac{1 - m_\rho^2/m_B^2}{1 - m_{K^*}^2/m_B^2} \right)^3 \xi^2 [1 + \Delta R]$$




$\xi = 1.17 \pm 0.09$   
 Flavour SU(3) breaking  
 $\Delta R = 0.1 \pm 0.1$   
 Weak annihilation correction



# Summary of Sides



CP Conserving Measurements  $|V_{ub}/V_{cb}|$ ,  $\Delta m_d$ ,  $|\Delta m_d/\Delta m_s|$ ,  $B \rightarrow \tau\nu$




# Global CKM Fit




Tree processes only:

$$|V_{ub}/V_{cb}|, B \rightarrow \tau\nu, \gamma, \pi - \alpha - \beta$$




Loop processes only:

$$|\epsilon_K|, \sin 2\beta, \Delta m_d, \Delta m_s$$




Tension between  $B \rightarrow \tau\nu$  and  $\sin 2\beta$  evident ( $\sim 2.4\sigma$ )

# Global CKM Fit



PDG 1997



Moriond 2009

$$\begin{aligned}
 A &= 0.8116^{+0.0097}_{-0.0241} \\
 \lambda &= 0.22521 \pm 0.00082 \\
 \bar{\rho} &= 0.139^{+0.025}_{-0.027} \\
 \bar{\eta} &= 0.341^{+0.016}_{-0.015}
 \end{aligned}$$

$$J_{CP} = (2.92 \pm 0.15) \times 10^{-5}$$


# The “Kπ Puzzle”



“Kπ Puzzle” published by Belle in Nature 2008....

Belle : 535M BB; Nature 452 (2008) 332

Direct CPV asymmetry in  $B \rightarrow K^+ \pi^-$  decays different to  $B \rightarrow K^+ \pi^0$  decays ??



HFAG 2009

$$A_{CP}(K^+ \pi^-) = -0.098^{+0.012}_{-0.011} \quad 8.1\sigma$$

$$A_{CP}(K^+ \pi^0) = +0.050 \pm 0.025$$

$$\Delta A_{K\pi} = A_{CP}(K^+ \pi^-) - A_{CP}(K^+ \pi^0) = -0.147 \pm 0.028 \quad 5.3\sigma$$


“Kπ Puzzle”

# Solutions to the “ $K\pi$ Puzzle”




See Nature commentary by Michael Peskin

M.Peskin : Nature 452 (2008) 293



If T and P dominant then  $\Delta A_{K\pi} \cong 0$   
(recent expectation)

Gronau & Rosner : PRD59 (1999) 113002



Enhancement of C with large strong phase to T  $\Rightarrow$  strong Interactions ??

Enhancement of  $P_{EW}$  from  $\Rightarrow$  New Physics

Also explains pattern of  $B \rightarrow \pi\pi$  and  $B \rightarrow \rho\rho$  Br's

Fleischer et al (& ref therein) : ArXiv 0806.2900

Li & Mishima : ArXiv 0901.1272


# Solutions to the “Kπ Puzzle”




Model independent method to detect NP

M.Gronau : PLB82 (2005) 627

$$\mathcal{A}_{CP}(K^+\pi^-) + \mathcal{A}_{CP}(K^0\pi^+) \frac{\mathcal{B}(K^0\pi^+) \tau_0}{\mathcal{B}(K^+\pi^-) \tau_+} = \mathcal{A}_{CP}(K^+\pi^0) \frac{2\mathcal{B}(K^+\pi^0) \tau_0}{\mathcal{B}(K^+\pi^-) \tau_+} + \mathcal{A}_{CP}(K^0\pi^0) \frac{2\mathcal{B}(K^0\pi^0)}{\mathcal{B}(K^+\pi^-)}$$




BaBar : 467M BB; PRD79 (2009) 052003  
 Belle : 657M BB; ArXiv:0809.4366



# FCNC Rare Decays: $B \rightarrow K^* \mu \mu$



FCNC  $b \rightarrow s$  transition, very sensitive to NP




The forward-backward asymmetry arises from the interference between  $\gamma$  and  $Z^0$  contributions

$$A_{FB}(s = m_{\mu\mu}^2) = -C_{10} \xi(s) \left[ \text{Re}(C_9) F_1 + \frac{1}{s} C_7 F_2 \right]$$

The zero crossing point is most theoretically clean

$$s_0^{SM} = 4.36_{-0.31}^{+0.33} \text{ GeV}^2$$

Beneke et al; EPJC41 (2005) 173



Ali et al; PLB273 (1991) 505






# FCNC Rare Decays: $B \rightarrow K^* \mu \mu$



Recent results of  $A_{FB}$  from the B factories show interesting behaviour

$$A_{FB}(s = m_{\mu\mu}^2) = \frac{N_F - N_B}{N_F + N_B}$$



BaBar : 384M BB; ArXiv:0804.4412

Belle : 657M BB; ArXiv:0904.0770



Note: opposite sign convention

Data points tend to be above the SM curve  
Is the sign of C7 wrong ? Need more statistics

# Some hints/puzzles exist...?



$$\beta_{\text{eff}}$$



“ $K\pi$  Puzzle”

# Summary



B factories have dramatically improved our understanding of flavour physics, far beyond expectations.

Clear demonstration of the SM CKM mechanism as the dominant source of CP violation

Overall good agreement in the global SM CKM fit


- A huge step forward on precision of  $\alpha$

A few puzzles remaining.....

- Tension between  $\sin 2\beta$  and  $|V_{ub}|$  with  $B \rightarrow \tau \nu$
- $K\pi$  puzzle
- $A_{FB}$  in  $B \rightarrow K^* \mu \mu$

Next lecture:

- What have we learnt from the Tevatron ?
- Is there still room for NP ?
- If so, can it be discovered at the LHC and beyond ?



New Physics is  
still hiding...





# CP Violation in Interference






Golden case: CP final state and single dominating amplitude

$$A_{f_{CP}}^{CP}(t) = \text{Im} \lambda_{f_{CP}} \sin(\Delta m t)$$



# $\beta$ : $b \rightarrow ccs$ modes





Mode	$\sin 2\beta$
$J/\psi K_S$	$0.657 \pm 0.036 \pm 0.012$
$J/\psi K_L$	$0.694 \pm 0.061 \pm 0.031$
<b><math>J/\psi K^0</math></b>	<b><math>0.666 \pm 0.031 \pm 0.013</math></b>
$\psi(2S) K_S$	$0.897 \pm 0.100 \pm 0.036$
$\chi_{c1} K_S$	$0.614 \pm 0.160 \pm 0.040$
$\eta_c K_S$	$0.925 \pm 0.160 \pm 0.057$
$J/\psi K^{*0}$	$0.601 \pm 0.239 \pm 0.087$
<b><math>c\bar{c}K^{(*)0}</math></b>	<b><math>0.687 \pm 0.028 \pm 0.012</math></b>

# Resolve 2-fold $\sin 2\beta$ ambiguity





- Amplitude analysis in  $B^0 \rightarrow [K_s^0 \pi^+ \pi^-]_D \pi^0$

Belle : 386M BB; PRL97 (2006) 081801  
BaBar : 383M BB; PRL99 (2007) 231802



- Time-dependent analysis in  $B^0 \rightarrow J/\psi K^{*0}$



- Time-dependent analysis of  $B^0 \rightarrow D^{*+} D^{*-} K_s^0$

BaBar : 230M BB; PRD74 (2006) 091101

Belle : 449M BB; PRD76 (2007) 072004




# $\beta_{\text{eff}} : b \rightarrow ccd$ modes



$B^0 \rightarrow J/\psi \pi^0$

$B^0 \rightarrow D^+ D^-$

$B^0 \rightarrow D^{*+} D^{*-}$





Good agreement with  $b \rightarrow ccs$  modes

$$S = -\sin 2\beta \quad C = 0$$



More info needed for C in  $D^+ D^-$  mode




$$\alpha : B \rightarrow \pi^0 \pi^0$$



# Isospin Analysis $B \rightarrow \pi\pi$



BaBar : 467M BB; ArXiv:0807.4226  
 Belle : 535M BB; PRL98 (2007) 211801




	<b>Br(10<sup>-6</sup>)</b>	<b>C</b>	<b>S</b>
$\pi^+\pi^-$	$5.5 \pm 0.4 \pm 0.3$	$-0.25 \pm 0.08 \pm 0.02$	$-0.68 \pm 0.10 \pm 0.03$
$\pi^+\pi^0$	$5.02 \pm 0.46 \pm 0.29$		
$\pi^0\pi^0$	$1.83 \pm 0.21 \pm 0.13$	$-0.43 \pm 0.26 \pm 0.05$	<b>No vertex Not possible</b>

# Isospin Analysis $B \rightarrow \pi\pi$



- Input  $Br(B^0 \rightarrow \pi^+\pi^-)$ ,  $Br(B^+ \rightarrow \pi^+\pi^0)$ ,  $Br(B^0 \rightarrow \pi^0\pi^0)$ ,  $C_{+-}$ ,  $S_{+-}$ ,  $C_{00}$



# Isospin Analysis $B \rightarrow \pi\pi$




- Grossmann-Quinn limit**

Grossmann, Quinn; arXiv:hep-ph/9712306

$$\sin^2 \Delta\alpha \leq \frac{Br(B^0 \rightarrow \pi^0\pi^0)}{Br(B^+ \rightarrow \pi^+\pi^0)}$$

- $\pi^0\pi^0$  rate too large to obtain useful limit
- $\pi^0\pi^0$  rate too small for a precision CP measurement

$$|\Delta\alpha| < 43^\circ \text{ @ } 90\% \text{ c.l.}$$




$$\alpha : [86^\circ, 108^\circ] \text{ @ } 68\% \text{ c.l.}$$

Standard peak

BaBar : 467M BB; ArXiv:0807.4226

Belle : 535M BB; PRL98 (2007) 211801



$$\alpha : [71^\circ, 109^\circ] \text{ @ } 68\% \text{ c.l.}$$

# $B \rightarrow \pi^+\pi^-\pi^0$




Three-pion final state: dominated by transitions through  $\rho$  mesons

- Interfering contributions from  $\rho^+\pi^-$ ,  $\pi^+\rho^-$  (and  $\rho^0\pi^0$ )


Snyder-Quinn method

Snyder&Quinn, PRD48 (1993) 2139; Quinn&Silva, PRD62 (2000) 054002

- Time-dependent Dalitz analysis of 6 decay amplitudes
- BW phase variations break degeneracy in solutions



$$\alpha = (87^{+45}_{-13})^\circ @ 68\% \text{ c.l.}$$



$$68^\circ < \alpha < 95^\circ @ 68\% \text{ c.l.}$$

BaBar : 375M BB; PRD76 (2007) 012004


Belle : 449M BB; PRL98 (2007) 221602

# $B \rightarrow \pi^+ \pi^- \pi^0$



Dalitz plot

$m^2(\pi^- \pi^0) \text{ GeV}^2$




$m^2(\pi^+ \pi^0) \text{ GeV}^2$



# ADS Method



Favoured and doubly-suppressed D decays e.g.  $K^+\pi^-$ ,  $K^+\pi^-\pi^0$



Same final state  
Large interference  
 $\sim O(1)$


Measure  $B^+$  and  $B^-$  yields to determine ADS observables:

$$R_{ADS} \equiv \frac{\Gamma(B^- \rightarrow D[\rightarrow f]K^-) + \Gamma(B^+ \rightarrow D[\rightarrow \bar{f}]K^+)}{\Gamma(B^- \rightarrow D[\rightarrow \bar{f}]K^-) + \Gamma(B^+ \rightarrow D[\rightarrow f]K^+)} = r_B^2 + r_D^2 + 2r_B r_D \cos \gamma \cos(\delta_B + \delta_D)$$


$$A_{ADS} \equiv \frac{\Gamma(B^- \rightarrow D[\rightarrow f]K^-) - \Gamma(B^+ \rightarrow D[\rightarrow \bar{f}]K^+)}{\Gamma(B^- \rightarrow D[\rightarrow f]K^-) + \Gamma(B^+ \rightarrow D[\rightarrow \bar{f}]K^+)} = 2r_B r_D \sin \gamma \sin(\delta_B + \delta_D) / R_{ADS}$$

ADS method useful at present to constrain  $r_B$   $r_D = \left| \frac{A(\bar{D}^0 \rightarrow f)}{A(D^0 \rightarrow f)} \right|$ ,  $\delta = \arg \left[ \frac{A(\bar{D}^0 \rightarrow f)}{A(D^0 \rightarrow f)} \right]$

# ADS Method



Belle : 657M BB; PRD78 (2008) 071901




No significant signal observed in any modes yet  
Upper limits on  $R_{ADS}$  translated to constraint  $r_B$



# ADS Method



BaBar : 379M BB; CKM008 prelim.




# GLW Method



D → CP eigenstates e.g.  $K^+K^-$ ,  $\pi^+\pi^-$  (CP+) and  $K_S\pi^0$  (CP-)

Measure  $B^+$  and  $B^-$  yields to determine GLW observables,  $A_{CP}$  and  $R_{CP}$



First result from CDF  
ArXiv:0809.4809

No direct measurement of  $\gamma$ , but helps in combination with other methods.  
Sensitivity to  $\gamma$  depends on strong phase.



- Observables

$$R_{CP\pm} \equiv \frac{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) + \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)}{2\Gamma(B^- \rightarrow D^0 K^-)} = 1 \pm 2r_B \cos\gamma \cos\delta_B + r_B^2$$

$$A_{CP\pm} \equiv \frac{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) - \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)}{\Gamma(B^- \rightarrow D_{CP\pm}^0 K^-) + \Gamma(B^+ \rightarrow D_{CP\pm}^0 K^+)} = \pm 2r_B \sin\gamma \sin\delta_B / R_{CP\pm}$$

- Alternate set

$$x_{\pm} = r_B \cos(\delta_B + \gamma) = \frac{R_{CP+}(1 \mp A_{CP+}) - R_{CP-}(1 \mp A_{CP+})}{4}$$

$$r_B^2 = x_{\pm}^2 + y_{\pm}^2 = \frac{R_{CP+} + R_{CP-} - 2}{2}$$


# GGSZ (Dalitz) Method






- Dalitz model

$$K_s \pi^+ \pi^-$$


$$K_s K^+ K^-$$



# Inclusive Vcb




# Exclusive Vcb



$$F(1) = 0.921 \pm 0.013 \pm 0.020$$

$$|V_{cb}| = (38.1 \pm 0.6 \pm 0.9) \times 10^{-3} \quad (D^*)$$

C. Bernard et al, ArXiv:0808.2519




$$G(1) = 1.074 \pm 0.018 \pm 0.016$$

$$|V_{cb}| = (39.7 \pm 1.4 \pm 0.9) \times 10^{-3} \quad (D)$$

M. Okamoto et al, NPPS140 (2005) 461

# $B \rightarrow \tau \nu$




# FCNC Rare Decays: $B \rightarrow K^* \mu \mu$





Expect Isospin asymmetry to be small  
( $< 15\%$ ) in the SM

$$A_I(s = m_{\mu\mu}^2) = \frac{(\tau_{B^+}/\tau_{B^0})N(B^0 \rightarrow K^{(*)0} \ell^+ \ell^-) - N(B^\pm \rightarrow K^{(*)\pm} \ell^+ \ell^-)}{(\tau_{B^+}/\tau_{B^0})N(B^0 \rightarrow K^{(*)0} \ell^+ \ell^-) + N(B^\pm \rightarrow K^{(*)\pm} \ell^+ \ell^-)}$$



Feldmann & Matias JHEP 0301, 074 (2003)



Intriguing large and negative  $A_I$

BaBar : 384M BB; ArXiv:0807.4119  
Belle : 657M BB; ArXiv:0904.0770