

The (d,p) reaction on ^{206}Hg

B. P. Kay¹, C. R. Hoffman¹, M. Avila¹, S. Bottoni¹, P. A. Butler², S. J. Freeman³,
L. P. Gaffney⁴, R. V. F. Janssens¹, M. Labiche⁵, R. D. Page², Zs. Podolyák⁶, R. Raabe⁷,
P. H. Regan⁶, M. Rudigier⁶, D. Santiago-Gonzalez^{8,1}, G. Savard¹, J. P. Schiffer¹, D. K. Sharp³,
J. F. Smith⁴, R. Talwar¹, and S. Zhu¹.

¹Argonne National Laboratory, ²University of Liverpool, ³University of Manchester,
⁴University of the West of Scotland, ⁵STFC Daresbury Laboratory, ⁶University of Surrey,
⁷KU Leuven, ⁸Louisiana State University

Requested shifts: 18

Beam: (ideally) 10 MeV/u ^{206}Hg , 1×10^6 Hz, >99% purity

Target: deuterated polyethylene $(\text{CD}_2)_n$



Installation: ISOL solenoidal spectrometer

INTC meeting, June 29, 2016

Motivation — general comments


$N = 127$ isotones below Pb

- *Terra incognita*. Below Pb, around $N = 126$, **very little known** (limited knowledge on masses, decays).
- **Evolution of single-particle states** has **not been explored** in nuclei around ^{208}Pb as these require **radioactive ion beams**.
- Data on 2^+ and 3^- in even nuclei allows us to make some assumptions.
- **Few / no theoretical studies** on single-particle excitations.




Motivation — loosely bound systems

s-states in loosely bound systems tend to linger below threshold—this feature seems to **dominate the structural changes in light nuclei**, and that results in **halo structures**. Does this characteristic of s-states play a role in loosely bound heavier systems?



Motivation — r-process physics



The proposed measurement

The $^{206}\text{Hg}(d,p)$ reaction at 10 MeV/u using the ISOL Solenoidal Spectrometer (ISS)

Why 10 MeV/u?

- Cross sections
- Angular momentum matching
- Angular distributions

Why ISS?

Resolution



- Charged-particle spectroscopy with **<100-keV Q-value resolution** using thin targets

Efficiency

- Limited only by geometrical acceptance, not intrinsic efficiency of the detectors.


Direct probe of excited states

- **Does not** require coincident γ -rays de-exciting the states (\therefore no concerns with isomers*, ground state, states not connected by γ -ray decay, etc).




*Isomers prevalent in the region around Pb
Cross sections estimated using DWBA code Ptolemy using standard parameterizations.

The ISOL Solenoidal Spectrometer (ISS)





Kinematics: $^{206}\text{Hg}(d,p)$, 10 MeV/u



No kinematic compression ($A = 0.31$), only **modest** kinematic shift (~ 17 keV/mm) *cf.* other techniques.

For this measurement the Si array used in the comparable HELIOS spectrometer at Argonne National Laboratory will be used in place of the one shown in this schematic.

The solenoidal-spectrometer technique



Simulation:

Marc Labiche, STFC Daresbury, using NPTool, assuming 40-keV intrinsic Si resolution¹ and the geometry of the ANL array, beam properties of the linac². Comparable to actual performance of the HELIOS spectrometer at ANL. Location of states states in ^{207}Hg estimated from Woods-Saxon calculations³.

¹Mean value for ANL Si array, J. C. Lighthall *et al.*, Nucl. Instrum. Methods Phys. Res. A 622, 97 (2010).

²Beam spot: 2.3 mm FWHM, Beam divergence: 1.8 mrad, Beam energy spread: 0.26%

³<http://www.volya.net>

Beam time request — 18 shifts

Assume:

1×10^6 Hz of ^{206}Hg , **>99%** purity desired, **10 MeV/u** desired, **$75 \mu\text{g}/\text{cm}^2$** CD_2 target, cross sections from DWBA calculations using standard parameterizations, **40%** solid angle for Si array over angular range $10^\circ \lesssim \theta_{\text{cm}} \lesssim 30^\circ$.

5 days (**18 shifts**) of beam on target yields **3000, 11300, 8700, 900, and 150** counts in single-particle states populated in $\ell = 0, 2, 4, 6, \text{ and } 7$ transfer.

1 additional day is requested for the optimization and calibration of the set up (1 shift), target changes (1 shift), and to record background events (1 shift).

Answer to TAC question: What degree of contamination could be tolerated?

This is an exploration of an unknown system. Though likely contaminants are known (isobars, ^{206}Pb) they could still obscure new states. We require the use of the VADLIS to achieve >99% purity.


Summary

- A study of the $^{206}\text{Hg}(d,p)$ reaction will be a flagship measurement—not possible at any other facility in the foreseeable future, particularly at this ideal energy for transfer.
- **First ever exploration** of single-particle structure of this region of the chart—terra incognita.
- Impact on nuclear structure — evolution of single-neutron states along $N=126$ — and on nuclear astrophysics, offering a first look at ***s-states below Pb*** on approach to the ***3rd r-process peak*** (*poorly understood in astrophysical models due to lack of data constraining them*).
- Solenoidal spectrometer technique ***well proven***, removing many complications plaguing other techniques. Ideal for ***extracting reliable spectroscopic factors*** from the data.
- Collaboration with the Argonne group—use of Si array, etc.

Supplemental material — beam purity

Use of VADLIS source


*No evidence of ^{206}Tl or ^{206}Pb
in the time of flight
spectrum*



Supplemental material — beam purity

Use of VADLIS source

From ^{208}Hg measurements, some small amount of Pb expected, though predicted to be about <600 ions/s cf. $>10^6$ ions/s of Hg.



Supplemental material — time lines


Ordering of events prior to experimental campaign

- **Cool down** the solenoid
- **Energize** and **verify the field**
- **Locate** in ISOLDE hall
- **Shield**
- Install various **mechanical components**
- Install **ANL Si array, electronics, DAQ**
- **Sources tests & take data with test beams for the beam line commissioning**

2016


2017

Supplemental material — ISS in situ




Supplemental material — ISS in situ


Schematic courtesy of Ian Burrows, STFC Daresbury




Supplemental material — ANL Si array



- 4 sides, 6 elements long
- Detector size, 9×50 mm
- 700- μm thick (e.g. ~ 10 MeV protons)
- Φ coverage, **0.48 of 2π**
- $\Omega_{\text{element}} = \sim 21 \text{ msr}$ (depending on kinematics, field, etc)
- $\Omega_{\text{array}} = \sim 500 \text{ msr}$




Supplemental material — r-process path





r-process model example (Image from: <http://www.ph.sophia.ac.jp/~shinya/research/research.html>)
From study described in S. Wanajo, S. Goriely, M. Samyn, and N. Itoh, ApJ 606, 1057 (2004)


Supplemental material — ^{208}Pb (d,p) at 10 MeV/u




Supplemental material — $^{208}\text{Pb} (d,p)$ at 10 MeV/u



Supplemental material — level structure



Supplemental material — $N = 127$ isotones



Supplemental material — fragmentation

Fragmentation of the $s_{1/2}$ strength

Fragmentation of the neutron s -state strength would be valuable data for **estimations of neutron-capture cross sections**.

In ^{207}Pb , below $N = 126$, the s -state strength appears at relatively high excitation energy, around 4.5-5 MeV in *at least 3 fragments*.

In ^{211}Po , one neutron outside 126, but above $Z = 82$, *two strong fragments* of the s -state strength are seen.

In ^{207}Hg , the $3s_{1/2}$ state could lie around 1.7 MeV in excitation energy (1.9 MeV below threshold like in ^{209}Pb), but could mix with the nearby core 2^+ (1.1 MeV) resulting in fragments lying closer to threshold.

A measurement of the (d,p) reaction on ^{206}Hg would provide a clear assessment of the fragmentation.