


# ZIMÁNYI SCHOOL

## 2012

December 3-7.  
Budapest, Hungary



# INITIAL ENERGY DENSITY IN LHC P+P COLLISIONS

04 Dec 2012

M. Csanad & T. Cs org 

# The Bjorken-estimate


2

- The original idea: energy density based on  $dE/dy$
- QGP critical  $\varepsilon: \leq 1 \text{ GeV/fm}^3$  (from  $\varepsilon_c = 6-8 \times T_c^4$ )
- Result (1900x cited)

$$E = N \frac{dE}{dy} \Delta y = N \frac{dE}{dy} \frac{1}{2} \frac{2d}{t} = \varepsilon \cdot Ad$$

$$\varepsilon_{Bj} = \frac{1}{R^2 \pi t_0} \frac{dE}{d\eta} = \frac{\langle E \rangle}{R^2 \pi t_0} \frac{dn}{d\eta}$$

- Needs correction!
- Ref.: Phys.Rev. D27 (1983)



# An advanced estimate

3

- Fact:  $d\eta/dy$  not flat
- Finiteness & acceleration
- Need accelerating solution  
of relativistic hydro

- Two modifications:


- $y \neq \eta$  &  $\eta_{\text{final}} \neq \eta_{\text{initial}}$

- Corrected initial energy density:

$$\varepsilon = \varepsilon_{Bj} \frac{dy}{d\eta_f} \frac{d\eta_f}{d\eta_i}$$

- Work needed to accelerate!



- Correction w.r.t. EoS:  $\varepsilon = \varepsilon_{Bj} (2\lambda - 1) \left( \frac{\tau_f}{\tau_i} \right)^{(\lambda-1)(2-c_{\text{sound}}^2)}$



# A solution of relativistic hydro

4

- Velocity:  $\tanh(\lambda\eta)$
- Acceleration:  $\lambda \neq 1$
- Density:  $(\tau/\tau_0)^\lambda$
- $dN/dy$  calculable!




- Compare this to RHIC data!
- $dN/dy$  measurement yields advanced initial  $\varepsilon$  estimate
- Significant correction at RHIC!

- Reference: Phys.Lett. B663 (2008) 306-311 (nucl-th/0605070)


# Initial energy density at RHIC

5

- Bjorken estimate from BRAHMS:  $5 \text{ GeV/fm}^3$
- Advanced estimate gives:  $\varepsilon = \varepsilon_{Bj} (2\lambda - 1) \left( \frac{\tau_f}{\tau_i} \right)^{(\lambda-1)(2-c_s^2)}$
- Correction:  $2-3x$ , result  $15 \text{ GeV/fm}^3$ , QCD agreement!
- Corresponds to  $T_{\text{ini}} \approx 2T_c \approx 340 \text{ MeV}$
- Confirmed by photon spectra at PHENIX, published 2010
- Reference e.g.: J.Phys.G35 (2008) 104128 (arXiv:0805.1562)



Normalized pseudorapidity distributions from BRAHMS



# Initial energy density at LHC

6

- Rough estimate possible via the Bjorken formula
  - Number of particles at midrapidity: 5.89 (CMS, ALICE)
  - Average energy:  $\langle m_t \rangle = 0.562 \text{ GeV}$  (CMS)
  - Initial radius of the system R:  $\sim 1.081 \text{ fm}$  (TOTEM,  $\sigma_{\text{inel}}$ )
  - Formation time  $\tau_0$ : 1 fm/c (conservative estimate)
- Energy density from this:

$$\varepsilon_{Bj} = \frac{1}{R^2 \pi \tau_0} \frac{dE}{d\eta} = \frac{\langle E \rangle}{R^2 \pi \tau_0} \frac{dn}{d\eta} = \frac{0.562 \times 5.89 \text{ GeV}/c}{3.67 \text{ fm}^3/c} \approx 0.90 \text{ GeV/fm}^3$$


- Just below critical? Important question!

# Correction from initial acceleration

7

- Initial acceleration pushes outer volume elements
- This modifies the  $d\eta/d\eta$  distribution
- Estimate acceleration from it!
- $\lambda = 1$ : no acceleration
- TOTEM fit:  $\lambda = 1.073 \pm 0.001 \pm 0.004$
- Without EoS: 20%


$$\varepsilon = \varepsilon_{Bj} (2\lambda - 1) \left( \frac{\tau_f}{\tau_i} \right)^{(\lambda-1)(1-c_s^2)}$$



# Advanced estimate

8


- Fit result:  $\lambda = 1.073$
- Conservatively:  $c_s^2 = 0.1$
- $\sim 25\%$  correction
- Input parameters:
  - $dN/dy$  at midrap.: 5.89
  - Average  $m_t$ : 0.562 GeV
  - Area: 3.67 fm $^2$ 
    - From cross sections
  - Freeze-out time / form. time: at least 2
- Bjorken result: 0.90 GeV/fm $^3$
- Corrected result: 1.14 GeV/fm $^3$



# Dependence on particle number

9


- Several multiplicity classes, 6-20, even 30 seen!



# Initial temperature estimate

10

- Temperature from  $\varepsilon \sim T^4$



# Sources of uncertainties

11

- For the correction factor  $\varepsilon/\varepsilon_{\text{Bj}}$ :
  - Fit parameter  $\lambda$
  - Statistical error (from the data)
  - Speed of sound  $c_s^2$
  - Duration  $\tau_f/\tau_i$
- For the original Bjorken-estimate:
  - Main uncertainty source: multiplicity at midrapidity  $dN/dy$
  - Area (if taken from cross-section): very precise
  - Formation time
  - Average transverse mass

# Systematic uncertainties

12

## □ All sources of uncertainties:

parameter	value	statistical uncertainty	systematic uncertainty
$\lambda$	1.073	0.1%	0.4% (from data)
$c_s^2$	0.1	-	-2%+0.2% (if $0.05 < c_s^2 < 0.5$ )
$\tau_f/\tau_0$	2	-	-4%+10% (if ratio 1.5...4)
$\tau_0$	1 fm/c	-	understimation on $\varepsilon$
R (from $\sigma_{inel}$ )	1.081 fm	0.5%	1.5% (from data)
$\langle m_t \rangle$	0.562 GeV/c <sup>2</sup>	0.5%	3% (from data)
dN/d $\eta$	5.985	0.2%	3% (from data)

## □ Conclusion:

$$\square \varepsilon_{ini} = (1.14 \pm 0.01(\text{syst}))^{+0.21}_{-0.16} (\text{syst}) \text{ GeV/fm}^3$$

# Is it unprecedented? Consequences?

13

- Bjorken and Landau worked out hydro for pp and pA
- Success of hydro to describe h+p, with  $\langle n \rangle = 7-8\dots$   
Phys.Lett. B422 (1998) 359-368
- Bjorken: it is not hadrons that play billiard balling
- If p+p is a complex system:
  - Radial flow
  - V2 scaling
  - HBT radii scaling
  - Low mass dilepton enhancement?
  - Direct photon enhancement?
- $R_{AA}$  might not be the best measure: divide by length scale?

# Summary

- Experimentally widely used Bjorken est.
- Advanced estimate: acceleration work, from  $d\eta/dn$
- Results on the initial  $\varepsilon$  for  $c_s^2=0.1$ , at  $\tau_f/\tau_{ini}=2$
- From TOTEM data:  $\varepsilon_{ini} = (1.14 \pm 0.01 \pm 0.2) \text{ GeV/fm}^3$ 
  - This at  $dN/dy=6$  & linearly rises with multiplicity!
- Critical energy density:  $1 \text{ GeV/fm}^3$
- Results not incompatible with supercrit. sQGP phase