



Contribution ID: 92

Type: **not specified**

The Chirality-Flow Formalism for the Standard Model

Tuesday 20 April 2021 15:50 (20 minutes)

Scattering amplitudes are often split up into their gauge ($\text{su}(N)$) and kinematic (two copies of complexified $\text{su}(2)$) components. Since the $\text{su}(N)$ gauge part is often calculated using flows of colour, it should similarly be possible to describe the $\text{su}(2) \oplus \text{su}(2)$ kinematics of an amplitude in terms of flows of chirality. In two recent papers (hep-ph:2003.05877 & hep-ph:2011.10075) we showed that this is indeed the case, introducing the chirality-flow formalism for Standard Model calculations. In the chirality-flow method (which simplifies the spinor-helicity method) Feynman diagrams can be directly written down in terms of Lorentz-invariant spinor inner products, allowing the simplest and most direct possible path from Feynman diagram to complex number. In this talk, I will introduce this method and show some examples

Primary author: LIFSON, Andrew (Lund University)

Co-authors: SJODAHL, Malin (Lund University); Dr REUSCHLE, Christian (Lund University); Mr ALNEFJORD, Joakim (Lund University)

Presenter: LIFSON, Andrew (Lund University)

Session Classification: Student Talks or Discussion Session