



LUND
UNIVERSITY

Soft Modelling and Heavy Ions (1)

Leif Lönnblad

Department of Astronomy
and Theoretical Physics
Lund University

CTEQ-MCnet School
Dresden 2021-09-13

Outline

- ▶ The (semi-) inclusive pp cross section(s)
- ▶ Minimum bias and Regge theory
- ▶ Multiple interactions
- ▶ Underlying events
- ▶ Summary: General Purpose Event Generators
- ▶ Introduction to Heavy Ions part.



What happens at LHC? (13 TeV)

Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Jets $p_{\perp} > 2 \text{ GeV}$	900 mb
Jets $p_{\perp} > 4 \text{ GeV}$	120 mb
Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb



What happens at LHC? (13 TeV)

Jets $p_{\perp} > 2 \text{ GeV}$	900 mb
Jets $p_{\perp} > 4 \text{ GeV}$	120 mb
Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb
BSM	$\sim 0? \text{ fb}$




What happens at LHC? (13 TeV)

Jets $p_{\perp} > 2 \text{ GeV}$	900 mb
Jets $p_{\perp} > 4 \text{ GeV}$	120 mb
Total	100 mb
Non-diffractive	56 mb
Elastic	22 mb
Diffractive	22 mb
Jets $p_{\perp} > 150 \text{ GeV}$	220 nb
W+Z	200 nb
Top	600 pb
Higgs	30 pb
BSM	$\sim 0? \text{ fb}$


Almost everything at LHC is pure QCD



Inclusive cross sections.




Minimum Bias: The typical pp collision




soft $gg \rightarrow gg$



Minimum Bias: The typical pp collision




soft $gg \rightarrow gg$




+ISR




Minimum Bias: The typical pp collision



soft $gg \rightarrow gg$




+ISR




Related to elastic scattering
via optical theorem




Minimum Bias: The typical pp collision




soft $gg \rightarrow gg$



+ISR




Related to elastic scattering
via optical theorem




Pomeron exchange




Minimum Bias: The typical pp collision




soft $gg \rightarrow gg$



+ISR



Related to elastic scattering
via optical theorem




Pomeron exchange

A **cut** pomeron in an elastic amplitude gives the non-diffractive cross section.

(From **Regge** theory [Regge. T, *Nuovo Cim.* **14** (1959) 951])




Multi-pomeron diagrams




Each cut pomeron contributes with evenly distributed particle production in the corresponding rapidity interval.
Like two flat strings.




Diffraction and triple-pomeron vertices



single diffractive excitation




central diffraction




double diffraction




Diffraction and triple-pomeron vertices




single diffractive excitation



central diffraction



double diffraction



?



Soft multiple interactions

- ▶ PHOJET [Engel et al.]
- ▶ Shrimps (SHERPA) [Zapp et al.]
- ▶ EPOS-LHC (also Heavy ions) [Werner et al.]

Where are the (mini-) jets?



(Semi-) Hard Multiple Interactions

Starting Point in PYTHIA:

$$\frac{d\sigma^H}{dk_{\perp}^2} = \sum_{ij} \int dx_1 dx_2 f_i(x_1, \mu_F^2) f_j(x_2, \mu_F^2) \frac{d\hat{\sigma}_{ij}^H}{dk_{\perp}^2}$$

The QCD $2 \rightarrow 2$ cross section is divergent $\propto \alpha_S^2(k_{\perp}^2)/k_{\perp}^4$

$\int_{k_{\perp c}^2} d\sigma^H$ will exceed the total (non-diffractive) pp cross section at the LHC for $k_{\perp c} \lesssim 5$ GeV.

There are more than one partonic interaction per pp -collision

$$\langle N_H \rangle(k_{\perp c}) = \frac{\int_{k_{\perp c}^2} d\sigma^H}{\sigma^{\text{ND}}}$$



The trick in PYTHIA is to treat everything as if it is perturbative.

$$\frac{d\hat{\sigma}_{ij}^H}{dk_{\perp}^2} \rightarrow \frac{d\hat{\sigma}_{ij}^H}{dk_{\perp}^2} \times \left(\frac{\alpha_S(k_{\perp}^2 + k_{\perp 0}^2)}{\alpha_S(k_{\perp}^2)} \cdot \frac{k_{\perp}^2}{k_{\perp}^2 + k_{\perp 0}^2} \right)^2$$

Where $k_{\perp 0}^2$ is motivated by colour screening (saturation) and is dependent on collision energy.

$$k_{\perp 0}(E_{\text{CM}}) = k_{\perp 0}(E_{\text{CM}}^{\text{ref}}) \times \left(\frac{E_{\text{CM}}}{E_{\text{CM}}^{\text{ref}}} \right)^{\epsilon \sim 0.16}$$

(using handwaving about the the rise of the total cross section)



The total and non-diffractive cross section is put in by hand (or with a Donnachie—Landshoff parameterization).

- Pick a hardest scattering according to

$$\frac{1}{\sigma^{\text{ND}}} \frac{d\sigma^H}{dk_{\perp}^2} \times \exp \left(- \int_{k_{\perp}^2} dq_{\perp}^2 \frac{1}{\sigma^{\text{ND}}} \frac{d\sigma^H}{dq_{\perp}^2} \right)$$

- Pick an impact parameter, b , from the overlap function (high k_{\perp} gives bias for small b).
- Generate additional scatterings with decreasing k_{\perp} using $d\sigma^H(b)/\sigma^{\text{ND}}$



Hadronic matter distributions

We assume that we have factorization

$$\mathcal{L}_{ij}(x_1, x_2, b, \mu_F^2) = \mathcal{O}(b) f_i(x_1, \mu_F^2) f_j(x_2, \mu_F^2)$$

$$\mathcal{O}(b) = \int dt \int dx dy dz \rho(x, y, z) \rho(x + b, y, z + t)$$

Where ρ is the matter distribution in the proton
(note: general width determined by σ^{ND})

- ▶ A simple Gaussian
- ▶ Double Gaussian
- ▶ x-dependent Gaussian



x-dependent overlap

Small-x partons are more spread out

$$\rho(r, x) \propto \exp\left(-\frac{r^2}{a^2(x)}\right)$$

with $a(x) = a_0(1 + a_1 \log 1/x)$

Note that high k_{\perp} generally means higher x and more narrow overlap distribution.



A note on resummation

There are many scales in an event: $S, \hat{s}, -\hat{t} \sim k_{\perp}^2, \Lambda_{\text{QCD}}^2, m_W^2, \dots$

Every time we have two widely separated scales there may be large logarithms that need to be resummed:

- ▶ $k_{\perp}^2 \gg \Lambda_{\text{QCD}}^2$: DGLAP
- ▶ $S \gg k_{\perp}^2$: BFKL (CCFM)
- ▶ $\hat{s} \gg k_{\perp}^2$: FKL



A note on resummation

There are many scales in an event: $S, \hat{s}, -\hat{t} \sim k_{\perp}^2, \Lambda_{\text{QCD}}^2, m_W^2, \dots$

Every time we have two widely separated scales there may be large logarithms that need to be resummed:

- ▶ $k_{\perp}^2 \gg \Lambda_{\text{QCD}}^2$: DGLAP
- ▶ $S \gg k_{\perp}^2$: BFKL (CCFM)
- ▶ $\hat{s} \gg k_{\perp}^2$: FKL



A note on resummation

There are many scales in an event: $S, \hat{s}, -\hat{t} \sim k_{\perp}^2, \Lambda_{\text{QCD}}^2, m_W^2, \dots$

Every time we have two widely separated scales there may be large logarithms that need to be resummed:

- ▶ $k_{\perp}^2 \gg \Lambda_{\text{QCD}}^2$: DGLAP
- ▶ $S \gg k_{\perp}^2$: BFKL (CCFM)
- ▶ $\hat{s} \gg k_{\perp}^2$: FKL



A note on resummation

There are many scales in an event: $S, \hat{s}, -\hat{t} \sim k_{\perp}^2, \Lambda_{\text{QCD}}^2, m_W^2, \dots$


Every time we have two widely separated scales there may be large logarithms that need to be resummed:

- ▶ $k_{\perp}^2 \gg \Lambda_{\text{QCD}}^2$: DGLAP
- ▶ $S \gg k_{\perp}^2$: BFKL (CCFM)
- ▶ $\hat{s} \gg k_{\perp}^2$: FKL



Is it reasonable to use collinear factorization even for very small k_{\perp} ?

Soft interactions means very small x , should we not be using k_{\perp} -factorization and BFKL?



For very small x and small k_{\perp} we also have *saturation*



Energy–momentum conservation

Each scattering consumes momentum from the proton, and eventually we will run out of energy.

- ▶ Continue generating MI's with decreasing k_{\perp} , until we run out of energy.
- ▶ Or rescale the PDF's after each additional MI.
(Taking into account flavour conservation).

Note that also initial-state showers take away momentum from the proton.



Interleaved showers

When do we shower?

- ▶ First generate all MI's, then shower each?
- ▶ Generate shower after each MI?

Is it reasonable that a low- k_{\perp} MI prevents a high- k_{\perp} shower emission? Or vice versa?

- ▶ Include MI's in the shower evolution



Interleaved showers

When do we shower?

- ▶ First generate all MI's, then shower each?
- ▶ Generate shower after each MI?

Is it reasonable that a low- k_{\perp} MI prevents a high- k_{\perp} shower emission? Or vice versa?

- ▶ Include MI's in the shower evolution



After the primary scattering we can have

- ▶ Initial-state shower splitting, P_{ISR}
- ▶ Final-state shower splitting, P_{FSR}
- ▶ Additional scattering, P_{MI}
- ▶ Rescattering of final-state partons, P_{RS}

Let them compete


$$\frac{d\mathcal{P}_a}{dk_{\perp}^2} = \frac{dP_a}{dk_{\perp}^2} \times \exp - \left(\int_{k_{\perp}^2} (dP_{\text{ISR}} + dP_{\text{FSR}} + dP_{\text{MI}} + dP_{\text{RS}}) \right)$$



Colour Connections




Every MI will stretch out new colour-strings.


Evidently not all of them can stretch all the way back to the proton remnants.



To be able to describe observables such as $\langle p_{\perp} \rangle (n_{\text{ch}})$ we need (a lot of) colour (re-)connections.







Beyond simple strings

What if we kick out two valens quarks from the same proton?


Normally it is assumed that the proton remnant has a di-quark, giving rise to a leading baryon in the target fragmentation.

PYTHIA8 has can hadronize **string junctions**
(also used for baryon-number violating BSM models)


Non-trivial baryon number distribution in rapidity.



What is the Underlying Event?




What is the Underlying Event?



Everything except the **hard sub-process**?



What is the Underlying Event?



Everything except the **hard sub-process** and **initial-** and **final-**state showers?



Subtracting underlying events from jets.

- ▶ ISR adds energy
- ▶ FSR removes energy
- ▶ UE adds energy
- ▶ Hadronization removes energy

Some of these can be made to cancel each other by adjusting the size of the jet cone.

But we still need to understand the underlying event.



Subtracting underlying events from jets.

- ▶ ISR adds energy
- ▶ FSR removes energy
- ▶ UE adds energy
- ▶ Hadronization removes energy

Some of these can be made to cancel each other by adjusting the size of the jet cone.

But we still need to understand the underlying event.



UE is not MB

- ▶ Harder processes gives a bias towards larger overlap (smaller b) giving more UE.
- ▶ The UE fluctuates — we can't just subtract a number
- ▶ Beware of jet cuts in a steeply falling spectrum





UE is not MB

- ▶ Harder processes gives a bias towards larger overlap (smaller b) giving more UE.
- ▶ The UE fluctuates — we can't just subtract a number
- ▶ Beware of jet cuts in a steeply falling spectrum

Also relevant for pile-up





A note on Tuning

The Min-bias and UE machineries contains a fair number of parameters that need to be tuned to data. In PYTHIA we have:

- ▶ Soft regularisation parameters
- ▶ Overlap function parameters
- ▶ Cross section parameterisations
- ▶ Colour reconnection parameters
- ▶ Intrinsic transverse momenta
- ▶ PDF choices
- ▶ ...



Global Tuning

General purpose event generators should describe everything.
They should not be tuned to a single observable.

- ▶ Hadronization parameters and final-state showers can be tuned to e^+e^- data (LEP).
- ▶ Initial-state showers and UE/MPI can be tuned to MB data.
- ▶ Anythings else should be fixed by measured Standard Model parameters.
- ▶ ... in principle



Global Tuning

General purpose event generators should describe everything.
They should not be tuned to a single observable.

- ▶ Hadronization parameters and final-state showers can be tuned to e^+e^- data (LEP).
- ▶ Initial-state showers and UE/MPI can be tuned to MB data.
- ▶ Anything else should be fixed by measured Standard Model parameters.
- ▶ ... in principle



Jet universality



There may be problems with flavour and meson/baryon issues.

Also at LEP there were mainly quark jets, gluon jets are softer and not very well measured.

At LHC there will be very hard gluon jets.

We need to check that jet universality works.





General Purpose Event Generators

There are only a few programs which deals with the whole picture of the event generation

- ▶ Hard sub-processes
- ▶ Parton showers
- ▶ Multiple interactions
- ▶ Hadronization
- ▶ Decays



Many more programs deal with a specific part of the event generation

- ▶ Hard subprocess: AlpGen, MadEvent, ... can be used with other generators using the Les Houches interface (but be sure to do proper merging)
- ▶ Parton Shower: ARIADNE, CASCADE, Vincia, DIRE, ... need to be integrated with a specific general purpose generator
- ▶ Multiple interactions: JIMMY (HERWIG) Shrimps (SHERPA)
- ▶ Hadronization (?)
- ▶ Decays: Tauola, EvtGen, typically called from within other generators.



PYTHIA8

- ▶ A few simple MEs, the rest from Les Houches
- ▶ k_{\perp} -ordered initial-/final-state DGLAP-based shower
- ▶ (N)LO multi-leg matching (not automatic)
- ▶ Multiple interactions interleaved with shower
- ▶ Lund String Fragmentation
- ▶ Particle decays

<https://pythia.org>



HERWIG++

- ▶ Construction of arbitrary MEs using helicity amplitudes
- ▶ Angular ordered and dipole shower
- ▶ Different matching schemes via MatchBox
- ▶ Soft+hard multiple interactions
- ▶ Cluster hadronization
- ▶ Particle decays with correlations

<http://projects.hepforge.org/herwig>



SHERPA

- ▶ Built-in automated ME generator
- ▶ Dipole-based shower
- ▶ Semi-automatic (N)LO multi-leg matching
- ▶ Multiple interactions (\sim old PYTHIA) with some CKKW features (also Shrimps)
- ▶ Cluster hadronization (string fragmentation via old PYTHIA).
- ▶ Standard particle decays.

<https://sherpa-team.gitlab.io>



Related Tools

Matrix Element Generators

- ▶ **MadGraph5(aMC@NLO)**
- ▶ POWHEG
- ▶ ALPGEN
- ▶ HELAC
- ▶ CompHEP
- ▶ ...

PDF parametrizations

- ▶ LHAPDF



Rivet.hepforge.org

(Buckley et al.)

Analyze Event Generator output and compare with published experimental data, using exactly the same cuts, triggers, etc.

400+ analyses are already in there.


If you want to make your analyses useful for others —
[Publish them in Rivet!](#)

Connected to *Professor* for tuning of parameters



MCplots.cern.ch

(Skands et al.)




pp vs. AA (from the pp point of view)

My immediate reactions when encountering Heavy Ion physics:


- ▶ That's just smashing bunches of nucleons together!
- ▶ Who is this Glauber guy anyway?
- ▶ You do you mean with centrality?
- ▶ When is many particles too many?
- ▶ I'm from Lund, I want to use string fragmentation!
- ▶ You measured what?





$$T_C \sim 170 \text{ MeV} \sim 2 \cdot 10^{12} \text{ K}$$






Pb+Pb @ $\sqrt{s} = 2.76$ ATeV

2010-11-08 11:30:46


Fill : 1482

Run : 137124


Event : 0x00000000D3BBE693





Flow




Fourier decomposition:
 $dN/d\Delta\phi = 1 + 2 v_2 \cos(2 \Delta\phi)$



Jet quenching



The R_{AA} factor



$$R_{AA} = \frac{d^2 N^{AA} / dp_T d\eta}{\langle T_{AA} \rangle d^2 \sigma^{pp} / dp_T d\eta}$$

$$\langle T_{AA} \rangle \sigma^{pp} = \langle N_{coll} \rangle$$

N_{coll} is the # of binary collisions

For perturbative QCD processes:

$R_{AA} < 1$: suppression


$R_{AA} = 1$: no nuclear effects

$R_{AA} > 1$: enhancement




The ridge

(a) pp $\sqrt{s} = 7$ TeV, $N_{\text{ch}}^{\text{offline}} \geq 110$



(b) pPb $\sqrt{s_{NN}} = 5.02$ TeV, $220 < N_{\text{ch}}^{\text{offline}} \leq 260$



(c) PbPb $\sqrt{s_{NN}} = 2.76$ TeV, $220 < N_{\text{ch}}^{\text{offline}} \leq 260$

