

Form factors for semileptonic $B_s \rightarrow K$ and $B_s \rightarrow D_s$ decays

Jonathan Flynn
Physics & Astronomy
University of Southampton

RBC/UKQCD Collaboration

UC Berkeley/LBNL

Aaron Meyer

BNL and BNL/RBRC

Yasumichi Aoki (KEK)
Peter Boyle (Edinburgh)

Taku Izubuchi

Yong-Chull Jang

Chulwoo Jung

Christopher Kelly

Meifeng Lin

Hiroshi Ohki

Shigemi Ohta (KEK)

Amarjit Soni

CERN

Andreas Jüttner (Southampton)

Columbia University

Norman Christ

Duo Guo

Yikai Huo

Yong-Chull Jang

Joseph Karpie

Bob Mawhinney

Ahmed Sheta

Bigeng Wang

Tianle Wang

Yidi Zhao

University of Connecticut

Tom Blum
Luchang Jin (RBRC)
Michael Riberdy
Masaaki Tomii

Edinburgh University

Matteo di Carlo
Luigi Del Debbio
Felix Erben
Vera Gülpers
Tim Harris
Nelson Lachini
Raoul Hodgson
Michael Marshall
Fionn Ó hÓgáin
Antonin Portelli
James Richings
Azusa Yamaguchi
Andrew ZN Yong

KEK

Julien Frison

University of Liverpool

Nicolas Garron

Michigan State U

Dan Hoying

Milano Bicocca

Mattia Bruno

Peking University

Xu Feng

University of Regensburg

Davide Giusti
Christoph Lehner (BNL)

University of Siegen

Matthew Black
Oliver Witzel

University of Southampton

Nils Asmussen
Alessandro Barone
Jonathan Flynn
Ryan Hill
Rajnandini Mukherjee
Chris Sachrajda

U of Southern Denmark

Tobias Tsang

Stony Brook University

Jun-Sik Yoo
Sergey Syritsyn (RBRC)

Motivation: Marcella Bona UT fit update EPS HEP 26 July 2021²

V_{cb} and V_{ub}

from FLAG 2019 arXiv:1902.08191

$$|V_{cb}| \text{ (excl)} = (39.09 \pm 0.68) 10^{-3}$$

$$|V_{cb}| \text{ (incl)} = (42.16 \pm 0.50) 10^{-3}$$

from Bordone et al.

arXiv:2107.00604

$\sim 2.8\sigma$ discrepancy

from FLAG 2019 arXiv:1902.08191

$$|V_{ub}| \text{ (excl)} = (3.73 \pm 0.14) 10^{-3}$$


$$|V_{ub}| \text{ (incl)} = (4.19 \pm 0.17 \pm 0.18 \text{ [flat]}) 10^{-3}$$

from GGOU HFLAV 2021

adding a flat uncertainty

covering the spread
of central values

$\sim 1.5\sigma$ discrepancy



$$|V_{ub} / V_{cb}| \text{ (LHCb)} = (9.46 \pm 0.79) 10^{-2}$$

From B_s to K at high q^2

$|V_{cb}|$


$$|V_{ub} / V_{cb}| \text{ (LHCb)} = (7.9 \pm 0.6) 10^{-2}$$

From Λ_b , excluded following FLAG guidelines

[LHCb first $B_s \rightarrow K\mu\nu$ result¹]

Pseudoscalar to pseudoscalar semileptonic decay

Semileptonic decay of pseudoscalar meson B_s of mass M and momentum p to pseudoscalar meson P of mass m and momentum k , with $q = p - k$



$$\frac{d\Gamma(B_s \rightarrow P \ell \nu)}{dq^2} = \eta \frac{G_F^2 |V_{xb}|^2}{24\pi^3} \frac{(q^2 - m_\ell^2)^2 |\vec{k}|}{(q^2)^2} \left[\left(1 + \frac{m_\ell^2}{2q^2}\right) \vec{k}^2 |f_+(q^2)|^2 + \frac{3m_\ell^2}{8q^2} \frac{(M^2 - m)^2}{M^2} |f_0(q^2)|^2 \right]$$

Form factors f_+ and f_0 from decomposition

$$\langle P(k) | \mathcal{V}^\mu(0) | B_s(p) \rangle = 2f_+(q^2) \left(p^\mu - \frac{p \cdot q}{q^2} q^\mu \right) + f_0(q^2) \frac{M^2 - m^2}{q^2} q^\mu$$

where $\mathcal{V}^\mu = \bar{x} \gamma^\mu b$, with $x = u$ or c

Lattice setup

- Subset of 6 RBC/UKQCD 2+1-flavour DWF and Iwasaki gauge field ensembles 
- Three lattice spacings $a \sim 0.11, 0.08, 0.07$ fm, with $267 \text{ MeV} < M_\pi < 433 \text{ MeV}$
- Light and strange quarks: Shamir DWF, $M_5 = 1.8$
- Lattice spacings from combined RBC/UKQCD analysis³⁻⁵
- Bottom and charm quarks
 - Bottom quarks: RHQ [Christ, Li, Lin^{6,7}; Columbia variant of Fermilab action⁸] with three nonpt-tuned parameters $(m_0 a, c_P, \zeta)$ ⁹
 - Charm quarks: Möbius DWF, $M_5 = 1.6$ ^{4,5,10}
 - 3 masses below m_c^{phys} on C ensembles
 - 2 masses which bracket m_c^{phys} on M and F
- Relate continuum and lattice currents: \mathcal{V}_μ and V_μ ^{11,12}
$$\langle P | \mathcal{V}_\mu | B_s \rangle = Z_{V_\mu}^{bx} \langle P | V_\mu | B_s \rangle$$
$$Z_{V_\mu}^{bx} = \rho_{V_\mu}^{bx} \sqrt{Z_V^{xx} Z_V^{bb}}$$
$$V_0 = V_0^0 + c_t^3 V_0^3 + c_t^4 V_0^4$$
$$V_i = V_i^0 + c_s^1 V_i^1 + c_s^2 V_i^2 + c_s^3 V_i^3 + c_s^4 V_i^4$$
 - $\rho_{V_\mu}^{bx}$ and $c_{t,s}^n$ computed perturbatively [Lehner]
 - Z_V^{bb} from $Z_V^{bb} \langle B_s | V_0 | B_s \rangle = 2M_{B_s}$
 - Z_V^{xx} from $Z_V^{xx} = Z_A^{xx} + O(am_{\text{res}})$

Extracting form factors

Calculate


$$f_{\parallel}(E) = \frac{\langle P | \mathcal{V}^0(0) | B_s \rangle}{\sqrt{2M_{B_s}}}, \quad f_{\perp}(E) = \frac{\langle P | \mathcal{V}^i(0) | B_s \rangle}{k^i \sqrt{2M_{B_s}}}$$

from which

$$f_0(q^2) = \frac{\sqrt{2M_{B_s}}}{M_{B_s}^2 - M_P^2} [(M_{B_s} - E)f_{\parallel}(E) + (E^2 - M_P^2)f_{\perp}]$$

$$f_+(q^2) = \frac{1}{\sqrt{2M_{B_s}}} [f_{\parallel}(E) + (M_{B_s} - E)f_{\perp}]$$

Extract from a correlator ratio (written here for $t_{\text{src}} = 0$)




$$R_{3,\mu}(t, t_{\text{snk}}, \vec{k}) = \frac{C_{3,\mu}(t, t_{\text{snk}}, \vec{k})}{\sqrt{C_2^P(t, \vec{k}) C_2^{B_s}(t_{\text{snk}} - t, \vec{0})}} \sqrt{\frac{2E_P}{e^{-E_P t - M_{B_s}(t_{\text{snk}} - t)}}}$$

$$f_{\parallel}^{\text{bare}}(\vec{k}) = \lim_{0 \ll t \ll t_{\text{snk}}} R_{3,0}(t, t_{\text{snk}}, \vec{k})$$



$$f_{\perp}^{\text{bare}}(\vec{k}) = \lim_{0 \ll t \ll t_{\text{snk}}} \frac{1}{p_P^i} R_{3,i}(t, t_{\text{snk}}, \vec{k})$$

$B_s \rightarrow K$ form factor extraction





$f_{||}$ on coarse (C1)
ensemble

$B_s \rightarrow K$ chiral-continuum extrapolation





$$f^{B_s \rightarrow K}(M_\pi, E_K, a^2) = \frac{\Lambda}{E_K + \Delta} \left[c_0 \left(1 + \frac{\delta f(M_\pi^S) - \delta f(M_\pi^P)}{(4\pi f_\pi)^2} \right) + c_1 \frac{\Delta M_\pi^2}{\Lambda^2} + c_2 \frac{E_K}{\Lambda} + c_3 \frac{E_K^2}{\Lambda^2} + c_4 (a\Lambda)^2 \right]$$

$B_s \rightarrow K$ cumulative error budget



$B_s \rightarrow D_s$ charm inter-/extrapolation and chiral-continuum extrapolation




$$f(q^2, a, M_\pi, M_{D_s}) = \left[c_0 + \sum_{j=1}^{n_{D_s}} c_{1j} h\left(\frac{M_{D_s}}{\Lambda}\right)^j + c_2 (a\Lambda)^2 \right] P_{a,b}(q^2/M_{B_s}^2)$$


$$h\left(\frac{M_{D_s}}{\Lambda}\right) = \frac{\Lambda}{M_{D_s}} - \frac{\Lambda}{M_{D_s}^{\text{phys}}}$$

$$P_{a,b}(x) = \frac{1 + \sum_{i=1}^a a_i x^i}{1 + \sum_{i=1}^b b_i x^i}$$


$B_s \rightarrow D_s$ cumulative error budget



▶ fit sys



z-fits



- Do z-fits after χ -ctm extrapolation
- Use BGL¹³⁻¹⁹ and BCL²⁰ for $B_s \rightarrow K$ and $B_s \rightarrow D_s$
- Example shown is BCL fit for $B_s \rightarrow K$, with $f^+(q^2)(1 - q^2/m_{B^*}^2)$ (lower) and $f^0(q^2)$ (upper) plotted

R ratios for LFU tests

$$R(P) = \frac{\int_{m_\tau^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\tau\bar{\nu}_\tau)}{dq^2}}{\int_{m_\ell^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\ell\bar{\nu}_\ell)}{dq^2}}$$

$$R^{\text{new}}(P) = \frac{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\tau\bar{\nu}_\tau)}{dq^2}}{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \frac{\omega_\tau(q^2)}{\omega_\ell(q^2)} \frac{d\Gamma(B_{(s)} \rightarrow P\ell\bar{\nu}_\ell)}{dq^2}}$$

- Adopt idea proposed for $B_{(s)} \rightarrow V$ decays [Isidori-Sumensari²¹]
 - Common integration range; $q_{\min}^2 \geq m_\tau^2$ [Freytsis et al²², Bernlochner et al²³, Soni²⁴]
 - Same weights for vector parts in integrands for τ and ℓ
- Write

$$\frac{d\Gamma(B_{(s)} \rightarrow P\ell\nu)}{dq^2} = \Phi(q^2) \omega_\ell(q^2) [F_V^2 + (F_S^\ell)^2]$$

$$\Phi(q^2) = \eta \frac{G_F^2 |V_{xb}|^2}{24\pi^3} |\vec{k}|$$

$$\omega_\ell(q^2) = \left(1 - \frac{m_\ell^2}{q^2}\right)^2 \left(1 + \frac{m_\ell^2}{2q^2}\right)$$

$$F_V^2 = \vec{k}^2 |f_+(q^2)|^2$$

$$(F_S^\ell)^2 = \frac{3}{4} \frac{m_\ell^2}{m_\ell^2 + 2q^2} \frac{(M^2 - m^2)^2}{M^2} |f_0(q^2)|^2$$

R ratios for LFU tests

$$R(P) = \frac{\int_{m_\tau^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\tau\bar{\nu}_\tau)}{dq^2}}{\int_{m_\ell^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\ell\bar{\nu}_\ell)}{dq^2}}$$

$$R^{\text{new}}(P) = \frac{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \frac{d\Gamma(B_{(s)} \rightarrow P\tau\bar{\nu}_\tau)}{dq^2}}{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \frac{\omega_\tau(q^2)}{\omega_\ell(q^2)} \frac{d\Gamma(B_{(s)} \rightarrow P\ell\bar{\nu}_\ell)}{dq^2}}$$

- Adopt idea proposed for $B_{(s)} \rightarrow V$ decays [Isidori-Sumensari²¹]
 - Common integration range; $q_{\min}^2 \geq m_\tau^2$ [Freytsis et al²², Bernlochner et al²³, Soni²⁴]
 - Same weights for vector parts in integrands for τ and ℓ
- Write

$$\frac{d\Gamma(B_{(s)} \rightarrow P\ell\nu)}{dq^2} = \Phi(q^2) \omega_\ell(q^2) [F_V^2 + (F_S^\ell)^2]$$

- If drop scalar contribution, $(F_S^\ell)^2$, in denominator ($m_\ell^2/2q^2 \leq m_\mu^2/2m_\tau^2 = 0.002$) expect

$$R^{\text{new,SM}}(P) = 1 + \frac{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \Phi(q^2) \omega_\tau(q^2) (F_S^\tau)^2}{\int_{q_{\min}^2}^{q_{\max}^2} dq^2 \Phi(q^2) \omega_\tau(q^2) F_V^2}$$

Summary

- Finalising results
- Additional lattice spacing since 2015 $B_s \rightarrow K$ and added $B_s \rightarrow D_s$
- Subsequently $B \rightarrow D\ell\nu$, vector final states, rare decays, B_c decays, ...

Backup

Related talks by RBC/UKQCD


- Ryan Hill: Semileptonic form factors for $B \rightarrow \pi/\nu$ decays [Thu 13:45]
- Michael Marshall: Semileptonic $D \rightarrow \pi/\nu$, $D \rightarrow K/\nu$ and $D_s \rightarrow K/\nu$ decays with 2 + 1 f domain wall fermions [Thu 14:30]
- Felix Erben: BSM $B_{(s)} - \bar{B}_{(s)}$ mixing on domain-wall lattices [Wed 6:15]

See also


- Davide Giusti: A new framework to tune an improved relativistic heavy-quark action [Poster]

Motivation

[CKMfitter²⁵]



[HFLAV²⁶]





- Determine fundamental parameters of the Standard Model, eg $|V_{ub}|$, $|V_{cb}|$
- Test/challenge SM: eg lepton flavour universality via $R(D^{(*)})$

Ensembles

	L	T	L_s	a^{-1}/GeV	am_I	am_s^{sea}	M_π/MeV	# cfgs	# sources
C1	24	64	16	1.785	0.005	0.040	340	1636	1
C2	24	64	16	1.785	0.010	0.040	433	1419	1
M1	32	64	16	2.383	0.004	0.030	302	628	2
M2	32	64	16	2.383	0.006	0.030	362	889	2
M3	32	64	16	2.383	0.008	0.030	411	544	2
F1S	48	96	12	2.785	0.002144	0.02144	267	98	24

◀ setup



Chiral-continuum fit systematics: $B_s \rightarrow K$



$\Delta f_i = |f_i^{\text{pref}} - f_i^{\text{alt}}|/f_i^{\text{pref}}$ for $i = 0, +$, for form-factor central values under fit variations.
Shaded band shows statistical uncertainty of preferred fit.



Chiral-continuum fit systematics: $B_s \rightarrow D_s$



$\Delta f_i = |f_i^{\text{pref}} - f_i^{\text{alt}}|/f_i^{\text{pref}}$ for $i = 0, +$, for form-factor central values under fit variations.
Shaded band shows statistical uncertainty of preferred fit.



References I

- [1] R Aaij et al. (LHCb), First observation of the decay $B_s^0 \rightarrow K^- \mu^+ \nu_\mu$ and Measurement of $|V_{ub}| / |V_{cb}|$, *Phys. Rev. Lett.* **126**, 081804 (2021), arXiv:2012.05143 [hep-ex].
- [2] M Bona et al. (UTfit), <http://www.utfit.org/>.
- [3] T Blum et al. (RBC/UKQCD), Domain wall QCD with physical quark masses, *Phys. Rev.* **D93**, 074505 (2016), arXiv:1411.7017 [hep-lat].
- [4] PA Boyle et al. (RBC/UKQCD), The decay constants f_D and f_{D_s} in the continuum limit of $N_f = 2 + 1$ domain wall lattice QCD, *JHEP* **12**, 008 (2017), arXiv:1701.02644 [hep-lat].
- [5] PA Boyle et al. (RBC/UKQCD), SU(3)-breaking ratios for $D_{(s)}$ and $B_{(s)}$ mesons, (2018), arXiv:1812.08791 [hep-lat].
- [6] NH Christ et al., Relativistic heavy quark effective action, *Phys. Rev.* **D76**, 074505 (2007), arXiv:hep-lat/0608006.
- [7] HW Lin and N Christ, Non-perturbatively determined relativistic heavy quark action, *Phys. Rev.* **D76**, 074506 (2007), arXiv:hep-lat/0608005.
- [8] AX El-Khadra et al., Massive fermions in lattice gauge theory, *Phys. Rev.* **D55**, 3933–3957 (1997), arXiv:hep-lat/9604004.
- [9] Y Aoki et al. (RBC/UKQCD), Nonperturbative tuning of an improved relativistic heavy-quark action with application to bottom spectroscopy, *Phys. Rev.* **D86**, 116003 (2012), arXiv:1206.2554 [hep-lat].
- [10] PA Boyle et al., Heavy domain wall fermions: the RBC and UKQCD charm physics program, *EPJ Web Conf.* **175**, edited by M Della Morte et al., 13013 (2018), arXiv:1712.00862 [hep-lat].

References II

- [11] S Hashimoto et al., Lattice QCD calculation of $\bar{B} \rightarrow D/\bar{\nu}$ decay form-factors at zero recoil, *Phys. Rev.* **D61**, 014502 (1999), arXiv:hep-ph/9906376 [hep-ph].
- [12] AX El-Khadra et al., The Semileptonic decays $B \rightarrow \pi/\nu$ and $D \rightarrow \pi/\nu$ from lattice QCD, *Phys.Rev.* **D64**, 014502 (2001), arXiv:hep-ph/0101023.
- [13] C Bourrely et al., Semileptonic decays of pseudoscalar particles ($M \rightarrow M' + \ell + \nu_\ell$) and short distance behavior of quantum chromodynamics, *Nucl.Phys.* **B189**, 157 (1981).
- [14] CG Boyd et al., Constraints on form-factors for exclusive semileptonic heavy to light meson decays, *Phys.Rev.Lett.* **74**, 4603–4606 (1995), arXiv:hep-ph/9412324 [hep-ph].
- [15] CG Boyd et al., Model independent determinations of $\bar{B} \rightarrow D$ (lepton), D^* (lepton) anti-neutrino form-factors, *Nucl. Phys.* **B461**, 493–511 (1996), arXiv:hep-ph/9508211 [hep-ph].
- [16] L Lellouch, Lattice constrained unitarity bounds for $\bar{B}^0 \rightarrow \pi^+/\bar{\nu}_l$ decays, *Nucl.Phys.* **B479**, 353–391 (1996), arXiv:hep-ph/9509358 [hep-ph].
- [17] CG Boyd and MJ Savage, Analyticity, shapes of semileptonic form-factors, and $\bar{B} \rightarrow \pi/\bar{\nu}$, *Phys.Rev.* **D56**, 303–311 (1997), arXiv:hep-ph/9702300 [hep-ph].
- [18] I Caprini et al., Dispersive bounds on the shape of $\bar{B} \rightarrow D^{(*)}\ell\bar{\nu}$ form-factors, *Nucl. Phys.* **B530**, 153–181 (1998), arXiv:hep-ph/9712417 [hep-ph].

References III

- [19] MC Arnesen et al., A precision model independent determination of $|V_{ub}|$ from $B \rightarrow \pi e\nu$, *Phys. Rev. Lett.* **95**, 071802 (2005), arXiv:hep-ph/0504209.
- [20] C Bourrely et al., Model-independent description of $B \rightarrow \pi/\nu$ decays and a determination of $|V_{ub}|$, *Phys. Rev.* **D79**, 013008 (2009), arXiv:0807.2722 [hep-ph].
- [21] G Isidori and O Sumensari, Optimized lepton universality tests in $B \rightarrow V\ell\bar{\nu}$ decays, *Eur. Phys. J. C* **80**, 1078 (2020), arXiv:2007.08481 [hep-ph].
- [22] M Freytsis et al., Flavor models for $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$, *Phys. Rev. D* **92**, 054018 (2015), arXiv:1506.08896 [hep-ph].
- [23] FU Bernlochner and Z Ligeti, Semileptonic $B_{(s)}$ decays to excited charmed mesons with e, μ, τ and searching for new physics with $R(D^{**})$, *Phys. Rev. D* **95**, 014022 (2017), arXiv:1606.09300 [hep-ph].
- [24] JM Flynn et al. (RBC/UKQCD), Nonperturbative calculations of form factors for exclusive semileptonic $B_{(s)}$ decays, PoS **ICHEP2020**, 436 (2021), arXiv:2012.04323 [hep-ph].
- [25] J Charles et al. (CKMfitter Group), <http://ckmfitter.in2p3.fr/>.
- [26] HFALAV: Heavy Flavor Averaging Group, Average of $R(D)$ and $R(D^*)$ for spring 2019, <https://hflav-eos.web.cern.ch/hflav-eos/semi/spring19/html/RDsDsstar/RDRDs.html>, 2019.