

# Investigating the conformal behaviour of $SU(2)$ with one adjoint Dirac flavor

Ed Bennett, Andreas Athenodorou, Georg Bergner, Biagio Lucini

Lattice 2021, 2021-07-30

Paper: arXiv:2103.10485 • Data: doi:10.5281/zenodo.5139617

Slides: <https://edbennett.github.io/lattice2021-talk>



Swansea University  
Prifysgol Abertawe



SUPERCOMPUTING WALES  
UWCHGYFRIFIADURA CYMRU



UNDEB EWROPEAIDD  
EUROPEAN UNION



Llywodraeth Cymru  
Welsh Government

**Cronfa Datblygu  
Rhanbarthol Ewrop  
European Regional  
Development Fund**

# Outline

- Introduction
  - Motivation
  - Previous work
- Results
  - Spectrum
  - Mass anomalous dimension
  - Chiral perturbation theory
- Conclusions

# Introduction

# Why $SU(2) + 1$ adjoint Dirac?

- Conformal window
- BSM strong dynamics
- Topological phase transitions

# What do we know?

- 1412.5994: Single value  $\beta = 2.05$ 
  - Scalar lightest state
  - Roughly constant mass ratios
  - Anomalous dimension  $\gamma_* \approx 0.95$
- 1507.08892: Adds  $\beta = 2.2$ 
  - Constant mass ratios persist
- 1912.11723 (Bi et al.):  $\beta = 2.05, 2.2$ 
  - Consistent masses
  - Composite fermion  $\approx$  hybrid fermion state
- Overall: hints of near-conformality

# What don't we know?

- Chiral limit
- Continuum limit
  - Hadron masses
  - Mass anomalous dimension



# Results

# What have we simulated?


- Unimproved Wilson gauge action
- Unimproved Wilson fermion action
  - as 1412.5994
  - Real action
- $\beta = 2.05, 2.1, 2.15, 2.2$
- $am_{2_s^+} \in (0.42, 1.08)$

# The spectrum


$\beta = 2.05$     $\beta = 2.1$     $\beta = 2.15$     $\beta = 2.2$



# Scalar as the lightest state





# Topological susceptibility




# Mass anomalous dimension


$\beta = 2.05$     $\beta = 2.1$     $\beta = 2.15$     $\beta = 2.2$   
 $\gamma_* = 0.713$     $\gamma_* = 0.773$



$\gamma_* = 0.888$




$\gamma_* = 0.933$




$L(am_{PCAC})^{1/(1+\gamma_*)}$

$L(am_{PCAC})^{1/(1+\gamma_*)}$

# Chiral perturbation theory



# Tensor/scalar ratio



(see also 1605.04258)

# Conclusions

# Summary

- Most ratios remain flat
- Scalar glueball remains lightest state
- Tensor-scalar glueball ratio falls short of gauge-gravity model predictions
- Topological susceptibility very close to quenched,  $N_f = 2$  data
- $\gamma_*$  decreases as  $\beta$  increases
- Results poorly described by chiral perturbation theory

# Possible interpretations

- QCD-like, but probed with too large  $am$ 
  - Inconsistent with observed mass hierarchy
- Conformal, with scaling deviations
- Conformal, but strongly influenced by bulk phase






# Ongoing work

- Smaller  $am$ :
  - To verify observed behaviour continues
- Larger  $\beta$ :
  - To stabilise approach to continuum
  - To probe scaling deviations
- Domain Wall Fermion action:
  - To reduce lattice artefacts towards chiral limit

Thank you


# Backup

# Mode number fits




# $2_s^+$ decay constant

$\beta = 2.05$     $\beta = 2.1$     $\beta = 2.15$     $\beta = 2.2$



# Near-marginal all-beta FSHS fits

$$\gamma_* = 1.016(13), \omega = -0.690(23)$$



(Fit form as in 1401.0195)