Heavy decaying dark matter at future neutrino radio telescopes

Rasmi Hajjar

Scuola Superiore Meridionale (IT) Università degli Studi di Napoli Federico II

rasmienrique.hajjarmunoz@unina.it

12-23 July 2021 Neutrinos en Colombia







Main goal of this work

- radio telescopes.



Rasmi Hajjar

Heavy Decaying Dark Matter at future V radio telescopes

 The main goal of this work is to forecast the limits that we can place on the lifetime of Heavy Dark Matter (HDM) particles using future neutrino

 We assume that the DM particles decay to a pair of SM particles and the minimal decaying DM scenario with only two parameters: $(m_{\rm DM}, \tau_{\rm DM})$





Introduction

- DM is one of the pillars of the standard cosmological model.
- We have only seen DM interacting gravitationally, no other direct observations.
- Indirect DM detection via CRs, gamma rays and neutrinos emitted by annihilation or decay of the DM particles (multi-messenger astronomy)



• We focus on the neutrino detection, where $E_{\nu} > 10$ PeV is observationally unexplored.





Neutrino fluxes from DM

- Indirect DM detection via neutrinos
- Minimal Decaying Dark Matter model:
- Assume DM decay into a pair of SM pa
- Decaying DM expected fluxes:

$$\frac{\mathrm{d}\Phi_{\nu_{\alpha}+\bar{\nu}_{\alpha}}^{\mathrm{gal.}}}{\mathrm{d}E_{\nu}\mathrm{d}\Omega} = \frac{1}{4\pi m_{\mathrm{DM}}\tau_{\mathrm{DM}}} \frac{\mathrm{d}N_{\alpha}}{\mathrm{d}E_{\nu}} \int_{0}^{\infty} \mathrm{d}s \,\rho_{\mathrm{DM}}[r(s,\ell,b)] \\ \frac{\mathrm{d}\Phi_{\nu_{\alpha}+\bar{\nu}_{\alpha}}^{\mathrm{ext.gal.}}}{\mathrm{d}E_{\nu}\mathrm{d}\Omega} = \frac{\Omega_{\mathrm{DM}}\rho_{c}}{4\pi m_{\mathrm{DM}}\tau_{\mathrm{DM}}} \int_{0}^{\infty} \frac{\mathrm{d}z}{H(z)} \frac{\mathrm{d}N_{\alpha}}{\mathrm{d}E_{\nu}} \bigg|_{E_{\nu}'=E_{\nu}(1+z)} \right\} \quad \frac{\mathrm{d}\Phi_{3\nu}^{\mathrm{DM}}}{\mathrm{d}E_{\nu}} = \sum_{\alpha} \int \mathrm{d}\Omega \left[\frac{\mathrm{d}\Phi_{\nu_{\alpha}+\bar{\nu}_{\alpha}}^{\mathrm{gal.}}}{\mathrm{d}E_{\nu}\mathrm{d}\Omega} + \frac{\mathrm{d}\Phi_{\nu_{\alpha}+\bar{\nu}_{\alpha}}^{\mathrm{ext.gal.}}}{\mathrm{d}E_{\nu}\mathrm{d}\Omega} \right]_{E_{\nu}'=E_{\nu}(1+z)}$$

Rasmi Hajjar

Heavy Decaying Dark Matter at future V radio telescopes

$$(m_{\rm DM}, \tau_{\rm DM})$$

articles: $(\rm DM \rightarrow f\bar{f})$

$$b\bar{b}$$

 $\tau^+ \tau^-$
 $\nu \bar{\nu}$

NuCo 2021



3/10

Neutrino fluxes from DM

• High energy \mathcal{V} flux is unknown. Other possible contributions:



colored regions from J. Álvarez-Muniz et. al. [1810.09994]

Heavy Decaying Dark Matter at future V radio telescopes

Rasmi Hajjar

Cosmogenic

guaranteed but uncertain magnitude, come from CRs interacting with CMB

Newborn Pulsars

higher expected astrophysical contribution in literature for neutrino radio telescopes

- Active Galactic Nuclei
- Gamma-ray bursts
- Flat-spectrum radio Quasars
- Black Hole jets embedded in large structures



Future radio v telescopes



<		GRAND 200k						
	1	IceCube-Gen2						en2
2029	2030	2031	2032	2033	2034	2035	2036	2037
future	future V radio telescopes NuCo 2021							





Methodology

- HDMSpectra to generate DM fluxes: C. W. Bauer et. al. [2007.15001]
- Astrophysical neutrinos act as a background.
- Conservative choice: highest theoretical astro fluxes.



Heavy Decaying Dark Matter at future *V* **radio telescopes**

Rasmi Hajjar



Methodology

• For each astrophysical scenario the probability to observe $N_{\rm obs}$ events is

$$p\left(N_{\text{obs}} \mid N_{\text{astro}}\right) = \frac{\left(N_{\text{astro}}\right)^{N_{\text{obs}}} e^{-N_{\text{astro}}}}{N_{\text{obs}}!}$$

- Conservative choice: constrain signals N_{events} of DM > N_{events} observed. \bullet
- Test statistic: (\mathscr{L} assumed Poisson)

$$TS(m_{DM}, \tau_{DM}) = \begin{cases} 0 & \text{for } n_{DM} < N_{obs} \\ -2\ln\left(\frac{\mathscr{L}(N_{obs}|n_{DM})}{\mathscr{L}(N_{obs}|N_{obs})}\right) & \text{for } n_{DM} \ge N_{obs} \end{cases}$$

For $m_{\rm DM}$ and $N_{\rm astro}$ we can determine the lifetime limits.

Rasmi Hajjar

Heavy Decaying Dark Matter at future V radio telescopes

N_{obs} stochastic random variable *N_{astro}* expected astrophysical events



New limits on HDDM



Rasmi Hajjar

Heavy Decaying Dark Matter at future V radio telescopes



NuCo 2021





 10^{15}

New limits on HDDM



- New limits with upcoming neutrino radio telescopes.
 - Neutrino channel: higher constraints.
 - Tau channel: new constraints.
 - b channel: complementary constraints to gamma rays.

Rasmi Hajjar

Heavy Decaying Dark Matter at future V radio telescopes



Conclusions

- Radio neutrino telescopes will have potential to detect a contribution coming from DM.
- HDM particles with $m_{DM} = 10^7 10^{15}$ GeV.
- 3 channels, 4 experiments and 2 different astrophysical signals.
- Future work: obtain limits using the current gamma-ray measurements for all channels.

Forecast analysis in order to set conservative bounds on the lifetime of





Thank you for your attention

http://www.astroparticle-th.unina.it

INTERVIEW NETWORK IN THEORY



rasmienrique.hajjarmunoz@unina.it

