

# Cosmic-ray acceleration and escape from post-adiabatic Supernova remnants

*Wednesday, 14 April 2021 19:15 (15 minutes)*

Supernova remnants are known to accelerate cosmic rays on account of their non-thermal emission of radio waves, X-rays, and gamma rays. Although there are many models for the acceleration of cosmic rays in Supernova remnants, the escape of cosmic rays from these sources is yet understudied.

We use our time-dependent acceleration code RATPaC to study the acceleration of cosmic rays and their escape in post-adiabatic Supernova remnants and calculate the subsequent gamma-ray emission.

We performed spherically symmetric 1-D simulations in which we simultaneously solve the transport equations for cosmic rays, magnetic turbulence, and the hydrodynamical flow of the thermal plasma in the test-particle limit. Our simulations span 100,000 years, thus covering the free-expansion, the Sedov-Taylor, and the beginning of the post-adiabatic phase of the remnant's evolution.

At later stages of the evolution cosmic rays over a wide range of energy can reside outside of the remnant, creating spectra that are softer than predicted by standard diffusive shock acceleration and feature breaks in the 10-100 GeV-range. The total spectrum of cosmic rays released into the interstellar medium has a spectral index of  $s \sim 2.4$  above roughly 10 GeV which is close to that required by Galactic propagation models. We further find the gamma-ray luminosity to peak around an age of 4,000 years for inverse-Compton-dominated high-energy emission. Remnants expanding in low-density media emit generally more inverse-Compton radiation matching the fact that the brightest known supernova remnants - RCW86, Vela Jr, HESSJ1721-347 and RXJ1713.7-3946 - are all expanding in low density environments.

**Primary author:** BROSE, Robert

**Presenter:** BROSE, Robert

**Session Classification:** Exploring the Galaxy: Supernova Remnants & Pulsar Wind Nebulae

**Track Classification:** SNR/PWN