

A SEMI-MICROSCOPIC DESCRIPTION OF ISOSCALAR GIANT MULTIPOLE RESONANCES IN MEDIUM-MASS CLOSED SHELL NUCLEI

Friday, 24 September 2021 14:55 (25 minutes)

Being a microscopically based extension of the standard [1] and nonstandard [2] versions of the continuum-random-phase approximation (cRPA), by taking the spreading effect into account, the semi-microscopic particle-hole (p-h) dispersive optical model (PHDOM) has been proposed [3] and successfully implemented for describing properties of various giant resonances (GRs) in medium-heavy mass closed-shell nuclei (See Ref. [4] and references therein). Within the model, the main relaxation modes of (p-h)-type states, associated with GRs, are together taken into account. These modes are: (i) Landau damping; (ii) coupling mentioned states to the single-particle continuum, and; (iii) coupling to many-quasiparticle configurations (the spreading effect). Landau damping and coupling to the continuum are described microscopically (in terms of a phenomenological mean field and Landau-Migdal p-h interaction), while the spreading effect is treated phenomenologically (in terms of the energy-averaged p-h self-energy term). That allows one to describe within PHDOM the main GR characteristics for a wide excitation-energy interval: (i) double transition density; (ii) strength distribution, and one-body "projected" transition density both related to an appropriate probing operator, and (iii) probabilities of direct one-nucleon decay.

The PHDOM version proposed in Ref. [3] in a rather general form has been adopted in Ref. [4] for describing main characteristics of isoscalar Multipole GRs in medium-heavy mass closed-shell nuclei. The $L = 0 - 3$ multipole resonances together with $L = 0, 2$ multipole overtones have been considered and a rather reasonable description of available experimental data has been obtained for the ^{208}Pb nucleus, taken as an appropriate example. Some of results obtained within cRPA (i.e., in neglecting the spreading effect) were found in agreement with the results obtained within the microscopic RPA-based approach of self-consistent Hartree-Fock, using Skyrme-type forces [5].

The present work is a direct continuation of the above-described study of Ref. [4]. The study is extended to medium-mass closed-shell nuclei $^{40,48}\text{Ca}$, ^{90}Zr , and ^{132}Sn . Calculation results are compared with available experimental data. Some of cRPA results are compared with the results of Ref. [5].

This work was supported in part by the Russian Foundation for Basic Research, under grant no. 19-02-00660 (M.L.G, B.A.T, M.H.U), and by the US Department of Energy, under grant no. DE-FG03-93ER40773 (S.S.).

1. S. Shlomo and G. Bertsch, Nucl. Phys. A 243, 507 (1975).
2. M. H. Urin, Nucl. Phys. A 811, 107 (2008).
3. M. H. Urin, Phys. Rev. C 87, 044330 (2013).
4. M. L. Gorelik, S. Shlomo, B. A. Tulupov, and M. H. Urin, Phys. Rev. C 103, 034302 (2021).
5. G. Bonasera, M. R. Anders, and S. Shlomo, Phys. Rev. C 98, 054316 (2018).

Primary author: Dr GORELIK, Mikhail (Moscow Economic School, Moscow 123022, Russia)

Co-authors: Dr SHLOMO, Shalom (Cyclotron Institute, Texas A&M University, College Station, TX 77843, USA); Dr TULUPOV, Boris (Institute for Nuclear Research, RAS, Moscow 117312, Russia); Prof. URIN, Mikhail (National Research Nuclear University "MEPhI", Moscow 115409, Russia)

Presenter: Dr GORELIK, Mikhail (Moscow Economic School, Moscow 123022, Russia)

Session Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei

Track Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei.