

ON THE TRUE AND FICTITIOUS ENHANCEMENTS OF THE FUNDAMENTAL SYMMETRY BREAKING EFFECTS

Wednesday 22 September 2021 14:45 (25 minutes)

All the enhancements of the P-violation effects in γ -transitions between the compound-nucleus states were analyzed in the classical paper [1] by I.S. Shapiro. The source of these effects is the weak interaction V_W leading to the fact that the wave function ψ of this state contains, besides the wave function of a definite parity ψ_1 , the small admixture of the opposite parity state ψ_2 :

$$\psi = \psi_1 + c\psi_2$$

$$\frac{c(A_a \cdot A_f)}{(A_a + cA_f)^2} \approx \frac{cA_f}{A_a} \equiv \frac{n}{d}$$

(1) The effect is defined by the ratio of the P-forbidden transition normalized by the total transition

(2) Here A_a and A_f are the amplitudes of the P-allowed and P-forbidden transitions. The review [1] indicates 3 types of enhancement: 1) kinematical enhancement, 2) structural enhancement and 3) dynamical enhancement. The kinematical enhancement appears when the allowed transition is the magnetic one which is smaller than the forbidden electric of the same multipolarity by the factor $(v/c) \approx 10$. The structural enhancement appears when the allowed transition amplitude comes to be unusually small due to some suppression caused by the structure of the initial and final states. One should point that both the kinematical and structural enhancements arise because of the decrease of the denominator d in Eq. (2). Only the dynamical enhancement is caused by the increase of the admixture coefficient:

$c = \langle \psi_1 | V_W | \psi_2 \rangle / (E_1 - E_2) = \frac{vP}{D}$ in the numerator of (2). Here vP is the weak interaction matrix element, while the enhancement of the admixture for the high-lying excited states is caused by their strongly decreased level spacing. It is assumed that the largest magnitude of the symmetry-breaking effect allows to measure it with the largest accuracy (i. e. with the smallest relative error). This assumption is shown to be often misleading. Indeed, the experimentally measured value (2) is the ratio of the normally distributed numbers of numerators n to denominators d . Taking their absolute errors to be σ and neglecting the correlation between them, one obtains for the relative error of the measured effect: $\sigma_R = \frac{\sigma}{\sqrt{n^2 + d^2}}$ We see that the dynamical enhancement of n decreases the relative error

References:

1. I.S.Shapiro, Sov. Phys.Uspekhi. **95**, 647 (1968).

Author: Prof. BUNAKOV, Vadim (Petersburg Nuclear Physics Institute, National Research Center Kurchatov Institute, Gatchina)

Presenter: Prof. BUNAKOV, Vadim (Petersburg Nuclear Physics Institute, National Research Center Kurchatov Institute, Gatchina)

Session Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei

Track Classification: Section 1. Experimental and theoretical studies of the properties of atomic nuclei.