

QUASI-ELASTIC KNOCKOUT OF NUCLEON FROM THE SHORT-RANGE NN CORRELATION AND RESCATTERINGS IN THE REACTION $^{12}\text{C}+p\rightarrow^{10}\text{A}+pp+N$

Monday 20 September 2021 14:50 (25 minutes)

Short-range correlated (SRC) NN pairs play an important role in structure of atomic nuclei and are studied using mainly electron beams [1]. A new step was done at BM@N in JINR [2] where the reaction $^{12}\text{C}+p\rightarrow^{10}\text{A}+pp+N$ is studied using the ^{12}C beam at energy of 4 GeV/nucleon in inverse kinematics providing interaction with the hydrogen target to probe the SRC pairs in the ^{12}C . In theoretical analysis [3] of the SRC effects in this reaction is used a properly modified approach developed earlier (see Ref. [4] and references therein) to describe the quasi-elastic knock-out of fast deuterons from the light nuclei ^{12}C and ^7Li by protons in the reactions (p,pd) and (p,nd) [5]. Elementary sub-processes in the (p,Nd) were the backward quasi-elastic scattering of the proton on the two-nucleon clusters $p\{pn\}\rightarrow pd$ and $p\{nn\}\rightarrow nd$ at the proton beam energy 670 MeV. As in Ref. [4], the spectroscopic amplitudes for NN-pairs in the ground state of the ^{12}C nucleus are calculated here within the translation-invariant shell model (TISM) with mixing configurations. Factorization of the two-nucleon momentum distribution over the internal n_{rel} (q_{rel}) and the c.m.s. n_{cm} ($k_{\text{c.m.}}$) momenta is assumed and for n_{rel} (q_{rel}) the squared deuteron (or singlet deuteron) wave function the CD Bonn NN-interaction potential is used. Relativistic effects in the sub-process $p+\{NN\}\rightarrow p+N+N$ of quasi-elastic knockout of nucleon from the SRC pair are taken into account in the light-front dynamics [6]. We found that the c.m. distribution of the deuteron clusters obtained within the TISM and used in [4], [5] to describe the (p,Nd) data [4] has to be modified considerably [6] to describe the kc.m. distribution of the SCR NN pairs measured in the electron data [7]. The ratio of the spin-singlet to spin-triplet pairs $\{pp\}_s/\{pn\}_t$ is calculated and found to be in agreement with existing data. Here the initial and final state interaction effects are estimated within the eikonal approximation using the Glauber model for the $N^{-10}\text{A}$ scattering. The one-loop approximation with elastic $N^{-10}\text{A}$ rescatterings is applied and the effect is found to be moderate.

This work is supported in part by the RFBR grant № 18-02-40046.

References:

1. O.Hen et al., Rev. Mod. Phys. **89**, 045002 (2017) .
2. M. Patsyuk et al., Nature Phys. **17**, 693 (2021); arXiv:2102.02626 [nucl-ex].
3. Yu.N. Uzikov, Izv. RAN, Ser. Fiz. **84** , 580 (2020).
4. M.A. Zhusupov, Yu.N. Uzikov, Fiz. El. Chast. At. Yadr. **18**, 323 (1987).
5. J. Ero et al., Nucl. Phys. A **372**, 317 (1981); D. Albrecht et al., Nucl. Phys. A **322**, 512(1979).
6. Yu.N. Uzikov, EPJ Web Conf. **222**, 03027 (2019).
7. E.O. Cohen et al., Phys. Rev. Lett. **121**, 092501 (2018).

Primary author: UZIKOV, Yuriy (Joint Institute for Nuclear Researches, M.V. Lomonosov State University, Dubna State University)

Presenter: UZIKOV, Yuriy (Joint Institute for Nuclear Researches, M.V. Lomonosov State University, Dubna State University)

Session Classification: Section 4. Relativistic nuclear physics, elementary particle physics and high-energy physics

Track Classification: Section 4. Relativistic nuclear physics, elementary particle physics and high-energy physics.