



Contribution ID: 4

Type: **Talk**

## **【415】 Electrically Pumped Difference Frequency Generation in Bragg-reflection Waveguides**

*Tuesday, 31 August 2021 17:30 (15 minutes)*

Nonlinear optical processes are widely used in quantum communication and computation, as well as laser engineering. In particular, difference frequency generation can be utilized to erase spectral distinguishability between single photons or generate laser light in otherwise hard to reach wavelength regimes. In this work, we present an AlGaAs Bragg-reflection waveguide with an embedded AlInGaAs quantum dot laser. By setting the temperature of this microscopic device, we allow for phase-matching between the internal laser emission and the interacting light in the telecom wavelength range. With an external, tunable telecom laser we investigate the generated photons from 1540 nm to 1630 nm and determine a nonlinear conversion efficiency of  $0.64(21)\%/W/cm^2$ .

**Primary author:** SCHLAGER, Alexander (University of Innsbruck)

**Co-authors:** Mr GÖTSCH, Maximilian (University of Innsbruck); Dr CHAPMAN, Robert (RMIT University); Dr FRICK, Stefan (University of Innsbruck); Ms THIEL, Hannah (University of Innsbruck); Dr SUCHOMEL, Holger (University of Würzburg); Dr KAMP, Martin (University of Würzburg); Prof. HÖFLING, Sven (University of Würzburg); Prof. SCHNEIDER, Christian (University of Oldenburg); Prof. WEIHS, Gregor (University of Innsbruck)

**Presenter:** SCHLAGER, Alexander (University of Innsbruck)

**Session Classification:** Atomic Physics and Quantum Optics

**Track Classification:** Atomic Physics and Quantum Optics