



Contribution ID: 228

Type: Talk

【133】 Does ARPES truly represent high-Tc superconductivity in cuprates?

Wednesday, 1 September 2021 17:30 (15 minutes)

An imminent doubt has always been around whether angle-resolved photoelectron spectroscopy (ARPES) of high-Tc superconductors, visualizing the superconducting gap in \mathbf{k} -space, can truly represent the intrinsic bulk spectral function whose response is distorted by energy- and \mathbf{k} -dependent matrix elements and small photoelectron escape depth. We address this fundamental question with soft-X-ray ARPES of the paradigm high-Tc cuprate $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$. The matrix elements are varied by spanning a dense \mathbf{k} -space grid formed by the lattice superstructure, and probing depth by changing the emission angle. Invariant magnitude of the measured superconducting gap proves the relevance of ARPES for the bulk superconductivity in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$ and calls for similar verification experiments on other high-Tc compounds.

Primary authors: YU, Tianlun (Swiss Light Source, Paul Scherrer Institut); MATT, Christian E. (Swiss Light Source, Paul Scherrer Institut); BISTI, Federico (Swiss Light Source, Paul Scherrer Institut); WANG, Xiaoliang (Swiss Light Source, Paul Scherrer Institut); SCHMITT, Thorsten (Swiss Light Source, Paul Scherrer Institut); CHANG, Johan (Physik-Institut, Universität Zürich); EISAKI, Hiroshi (National Institute of Advanced Industrial Science and Technology (AIST)); FENG, Donglai (State Key Laboratory of Surface Physics and Department of Physics, Fudan University); STROCOV, Vladimir N. (Swiss Light Source, Paul Scherrer Institut)

Presenter: STROCOV, Vladimir N. (Swiss Light Source, Paul Scherrer Institut)

Session Classification: Condensed Matter Physics

Track Classification: Condensed Matter Physics (KOND)