

Leptophilic New Physics

Fiona Kirk



University of
Zurich^{UZH}

PAUL SCHERRER INSTITUT



SPS-ÖPG Meeting 2021, Innsbruck

August 31, 2021

Outline

Leptophilic New Physics

- Vectorlike Leptons

- Singly Charged Scalar Singlet

- Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

- The Cabibbo Angle Anomaly

- Hints for LFUV in leptonic τ decays

Results

- Vectorlike Leptons

- Singly Charged Scalar Singlet

- Leptophilic Z' Bosons

Conclusions

Outline

Leptophilic New Physics

- Vectorlike Leptons

- Singly Charged Scalar Singlet

- Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

- The Cabibbo Angle Anomaly

- Hints for LFUV in leptonic τ decays

Results

- Vectorlike Leptons

- Singly Charged Scalar Singlet

- Leptophilic Z' Bosons

Conclusions

Vectorlike Leptons

Left- & right-handed parts couple in same way to electroweak gauge bosons: “L=R”

⇒ Vectorlike leptons can have “bare” masses (before EWSB):

$$\mathcal{L}_{VLL} = i\bar{\psi}\gamma_{\mu}D^{\mu}\psi - M_{\psi}\bar{\psi}\psi$$

⇒ Anomaly-free

Neutral under $SU(3)_c$

⇒ Only modify the electroweak sector

⇒ Couple to SM-leptons & -Higgs

⇒ Mix with SM-leptons

⇒ modify $W_{\ell\nu}$, $Z_{\nu\nu}$ & $Z_{\ell\ell}$

Vectorlike leptons respecting the SM gauge group

		$SU(3)$	$SU(2)_L$	$U(1)_Y$
SM	ℓ	1	2	-1/2
	e	1	1	-1
	ϕ	1	2	1/2
VLLs	N	1	1	0
	E	1	1	-1
	$\Delta_1 = (\Delta_1^0, \Delta_1^-)$	1	2	-1/2
	$\Delta_3 = (\Delta_3^-, \Delta_3^{--})$	1	2	-3/2
	$\Sigma_0 = (\Sigma_0^+, \Sigma_0^0, \Sigma_0^-)$	1	3	0
$\Sigma_1 = (\Sigma_1^0, \Sigma_1^-, \Sigma_1^{--})$	1	3	-1	

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Conclusions

The Singly Charged Scalar Singlet ϕ^+

	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$	
ϕ^+	1	1	1	new singly charged scalar

The Singly Charged Scalar Singlet ϕ^+

	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$	
ϕ^+	1	1	1	new singly charged scalar

⇒ no couplings to quarks, right-handed leptons

$$\mathcal{L} = \mathcal{L}_{\text{SM}} - \left(\frac{\lambda_{ij}}{2} \bar{L}_{a,i}^c \varepsilon_{ab} L_{b,j} \phi^+ + \text{h.c.} \right)$$

Here

$$\text{a,b: } SU(2)_L \text{ indices} \quad \varepsilon_{ab} = i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

i,j: flavour indices

The Singly Charged Scalar Singlet ϕ^+

	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$	
ϕ^+	1	1	1	new singly charged scalar

\Rightarrow no couplings to quarks, right-handed leptons

$$\mathcal{L} = \mathcal{L}_{\text{SM}} - \left(\frac{\lambda_{ij}}{2} \bar{L}_{a,i}^c \varepsilon_{ab} L_{b,j} \phi^+ + \text{h.c.} \right)$$

Here

$$a,b: SU(2)_L \text{ indices} \quad \varepsilon_{ab} = i\sigma_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

i,j : flavour indices

$\Rightarrow \lambda_{ij}$ antisymmetric in flavour

\Rightarrow automatically lepton flavour (universality) violating

Only 4 new parameters: λ_{12} , λ_{13} , λ_{23} and m_ϕ

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results

Vectorlike Leptons


Singly Charged Scalar Singlet

Leptophilic Z' Bosons


Conclusions

“Standard” Z' Bosons vs. Leptophilic Z' Bosons

- Simple Extension of the SM
- Predicted by many models
- Strongly constrained by di-jet, di-lepton searches



- Z' bosons that couple to leptons
 - \Rightarrow less constrained by LHC data (\Rightarrow LEP)
 - \Rightarrow can be lighter ($<$ TeV-scale)
- Gluons like quarks, Z' likes leptons?



Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results


Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Conclusions

The Cabibbo Angle Anomaly: Determination of V_{us}




“ V_{us} from β decays”:

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1 \Rightarrow V_{us} \sim \sqrt{1 - V_{ud}^2} \quad (V_{ub} \approx 0.0037 \sim 0)$$

$\Rightarrow 3\sigma$ tension (PDG)

Leptophilic Resolutions of the Cabibbo Angle Anomaly



- Singly charged scalar singlet
 \Rightarrow Modified Fermi constant, G_F
- Leptophilic Z' bosons
 \Rightarrow Modified Fermi constant, G_F
- Vectorlike Leptons
 \Rightarrow Modified $W\ell\nu$, $Z\ell\ell$, $Z\nu\nu$

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results


Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Conclusions

Lepton Flavour Universality Violation in τ decays



Plot for $\delta(\mu \rightarrow e \bar{\nu} \nu) = 0$

$$\left. \frac{\mathcal{A}(\tau \rightarrow \mu \bar{\nu} \nu)}{\mathcal{A}(\mu \rightarrow e \bar{\nu} \nu)} \right|_{\text{EXP}} = 1.0029(14)$$



$$\left. \frac{\mathcal{A}(\tau \rightarrow \mu \bar{\nu} \nu)}{\mathcal{A}(\tau \rightarrow e \bar{\nu} \nu)} \right|_{\text{EXP}} = 1.0018(14)$$

$$\left. \frac{\mathcal{A}(\tau \rightarrow e \bar{\nu} \nu)}{\mathcal{A}(\mu \rightarrow e \bar{\nu} \nu)} \right|_{\text{EXP}} = 1.0010(14)$$


$$\delta(\tau \rightarrow \mu \bar{\nu} \nu) = \frac{\mathcal{A}_{NP}(\tau \rightarrow \mu \bar{\nu} \nu)}{\mathcal{A}_{SM}(\tau \rightarrow \mu \bar{\nu} \nu)}$$

$$\delta(\tau \rightarrow e \bar{\nu} \nu) = \frac{\mathcal{A}_{NP}(\tau \rightarrow e \bar{\nu} \nu)}{\mathcal{A}_{SM}(\tau \rightarrow e \bar{\nu} \nu)}$$


Leptophilic New Physics in $\tau \rightarrow \mu \bar{\nu} \nu / \tau(\mu) \rightarrow e \bar{\nu} \nu$



Vectorlike Leptons



Singly charged scalar singlet, ϕ^+



Leptophilic Z' boson

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results


Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons


Conclusions

Vectorlike Leptons: Constraints from $\mu \rightarrow e$



Vectorlike Leptons: Best Fit

- One generation of singlet N coupling **only** to electrons
- One generation of triplet Σ_1 coupling **only** to muons



λ_N : $N\phi e$ -coupling, λ_{Σ_1} : $\Sigma_1\phi\mu$ -coupling; M_{N,Σ_1} : masses of N , Σ_1

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results


Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Conclusions

Singly Charged Scalar: Flavour bounds & EW data



Scenario with $\lambda_{13} = 0$

(constraints from $\mu \rightarrow e\gamma$, $\mu \rightarrow e$ conv.)

Best fit region for

- EW data & the Cabibbo angle anomaly
- $\tau \rightarrow \mu\bar{\nu}\nu / \tau(\mu) \rightarrow e\bar{\nu}\nu$

suggests

- $\text{Br}(\tau \rightarrow e\mu\mu) \sim 10^{-10} \frac{m_\phi^4}{(5 \text{ TeV})^4}$
- $10^{-11} \lesssim \text{Br}(\tau \rightarrow e\gamma) \lesssim 5 \times 10^{-11}$
- $|\lambda_{12}|^2 \sim \frac{0.05}{(1 \text{ TeV})^2}$ (\rightarrow mono photon)

Outline

Leptophilic New Physics

Vectorlike Leptons

Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Lepton Flavour (Universality) Violation

The Cabibbo Angle Anomaly

Hints for LFUV in leptonic τ decays

Results

Vectorlike Leptons


Singly Charged Scalar Singlet

Leptophilic Z' Bosons

Conclusions

Leptophilic Z' Bosons: $M_{Z'} = 1$ TeV, $g_{\mu\tau}$ only

$g_{\mu\tau}^{L,R}$ only



Conclusions

Recent hints for **Lepton Flavour Universality Violation**, in particular

- The Cabibbo Angle Anomaly
- Deviations from the SM in $\tau \rightarrow \mu \bar{\nu} \nu / \tau(\mu) \rightarrow e \bar{\nu} \nu$

can be addressed by **leptophilic new physics**.

Conclusions

Recent hints for **Lepton Flavour Universality Violation**, in particular

- The Cabibbo Angle Anomaly
- Deviations from the SM in $\tau \rightarrow \mu \bar{\nu} \nu / \tau(\mu) \rightarrow e \bar{\nu} \nu$

can be addressed by **leptophilic new physics**.

Examples presented here:

- **Vectorlike Leptons**
- **Singly Charged Scalar Singlet**
- **Leptophilic Z' Bosons**

Conclusions

Recent hints for **Lepton Flavour Universality Violation**, in particular

- The Cabibbo Angle Anomaly
- Deviations from the SM in $\tau \rightarrow \mu \bar{\nu} \nu / \tau(\mu) \rightarrow e \bar{\nu} \nu$

can be addressed by **leptophilic new physics**.

Examples presented here:

- **Vectorlike Leptons**
- **Singly Charged Scalar Singlet**
- **Leptophilic Z' Bosons**

Lepton Flavour Violation

- Forbidden in the SM
- Strongly constrained by experiment

⇒ Clean discovery channels

For “The Details”:

- **Vectorlike leptons**
arXiv:2008.01113/ JHEP 12 (2020) 166, with Andreas Crivellin, FK, Claudio Andrea Manzari, Marc Montull
- **Singly charged scalar singlet, ϕ^+**
arXiv:2012.09845/ Phys.Rev.D 103 (2021), Andreas Crivellin, FK, Claudio Andrea Manzari, Luca Panizzi
- **Leptophilic Z' bosons**
arXiv:2104.07680/ JHEP06 (2021) 068, Andrzej J. Buras, Andreas Crivellin, FK, Claudio Andrea Manzari and Marc Montull

Questions?

Backup slides


Possible Signatures of Leptophilic New Physics

- Cabibbo Angle Anomaly
- Anomalous magnetic moments of charged leptons Δa_ℓ
- Electric dipole moments of charged leptons d_ℓ
- Flavour violating decays of charged leptons $\ell \rightarrow 3\ell'$, $\ell \rightarrow \ell' \bar{\nu} \nu$
- Radiative leptonic decays $\ell \rightarrow \ell' \gamma$
- $\mu \rightarrow e$ conversion in nuclei
- Neutrino trident production $\nu N \rightarrow \nu \bar{\mu} \mu N$
- Contact interactions $e^+ e^- \rightarrow \ell^+ \ell^-$ (LEP)
- $Z - Z'$ mixing \Rightarrow modified Z -pole observables


The Cabibbo Angle

The Cabibbo angle can be determined from

- V_{ud} from superallowed β -decays ($0^+ - 0^+$ -transitions)



- V_{us} from τ -decays & from K -decays




- V_{cd} from $D \rightarrow \mu\nu$

New Physics Interpretations of the CAA arXiv: 2008.03261

- Direct contributions to β -decays
- Modified Wud -coupling
- Modified $Wl\nu$ -coupling
- Direct contributions to μ -decays
 \Rightarrow Modified Fermi constant, G_F

$$\delta(\mu \rightarrow e\bar{\nu}\nu) = \frac{\mathcal{A}_{NP}(\mu \rightarrow e\bar{\nu}\nu)}{\mathcal{A}_{SM}(\mu \rightarrow e\bar{\nu}\nu)}$$

$$\Rightarrow \boxed{G_F = G_F^{\text{SM}} (1 + \delta(\mu \rightarrow e\bar{\nu}\nu))}$$



Resolution of the Cabibbo Angle Anomaly

Assume

$$|V_{ud}|^2(1 - \delta(\mu \rightarrow e\nu\nu))^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9985(5)$$

$$\begin{aligned}\Rightarrow V_{us}^\beta &\equiv \sqrt{1 - (V_{ud}^\beta)^2 - |V_{ub}|^2} \\ &\simeq V_{us}^{\mathcal{L}} \left[1 + \left(\frac{V_{ud}^{\mathcal{L}}}{V_{us}^{\mathcal{L}}} \right)^2 \delta(\mu \rightarrow e\nu\nu) \right].\end{aligned}$$

V_{us}^β : determined via CKM unitarity from V_{ud}^β (β decays)

$V_{us}^{\mathcal{L}}$: values in the CKM matrix

Leptophilic Z' Bosons: The Pragmatic Approach

- Agnostic about origin of Z' boson
- Simply assume $M_{Z'} > M_{EW}$
- Leptophilic Z' boson \Rightarrow couplings to SM leptons ℓ, ν
- Can allow for $Z - Z'$ -mixing \Rightarrow correction to m_Z (destructive)


Lagrangian:

$$\mathcal{L} \supset \underbrace{-i g_{Z'}^\varphi Z'^\mu \varphi^\dagger \overleftrightarrow{D}_\mu \varphi}_{\substack{\Rightarrow Z - Z'\text{-mixing} \\ \Rightarrow \text{mod. pole mass}}} + \underbrace{\bar{\ell}_i \left(g_{ij}^L \gamma_\mu P_L + g_{ij}^R \gamma_\mu P_R \right) \ell_j Z'^\mu}_{Z'\ell\ell\text{-couplings}} + \underbrace{\bar{\nu}_i g_{ij}^L \gamma_\mu P_L \nu_j Z'^\mu}_{Z'\nu\nu\text{-couplings}}$$

$$\overleftrightarrow{D}_\mu = D_\mu - \overleftarrow{D}_\mu, \quad g_{Z'}^\varphi \in \mathbb{R}, \quad g_{ij}^{L/R} \text{ are Hermitian in flavour space}$$

Vectorlike Leptons: Best Fit

- one generation of singlet N coupling **only** to electrons
- one generation of triplet Σ_1 coupling **only** to muons



- 68% CL - - - 68% CL NNC
- 95% CL - - - 95% CL NNC
- 99.7% CL - - - 99.7% CL NNC

Observable	Measurement	SM Posterior	NP Posterior	Pull
M_W [GeV]	80.379(12)	80.363(4)	80.369(6)	0.56
$R \left[\frac{K \rightarrow \mu\nu}{K \rightarrow e\nu} \right]$	0.9978 ± 0.0020	1	1.00168(39)	-0.80
$R \left[\frac{\pi \rightarrow \mu\nu}{\pi \rightarrow e\nu} \right]$	1.0010 ± 0.0009	1	1.00168(39)	0.42
$R \left[\frac{\tau \rightarrow \mu\nu\bar{\nu}}{\tau \rightarrow e\nu\bar{\nu}} \right]$	1.0018 ± 0.0014	1	1.00168(39)	1.2
$ V_{us}^{K\mu 3} $	0.22345(67)	0.22573(35)	0.22519(39)	0.77
$ V_{ud}^{\beta} $	0.97365(15)	0.97419(8)	0.97378(13)	2.52

IC value: 73 (vs. IC_{SM} : 93)

IC-value

Information criterion (IC): allows for a comparison between different models within a Bayesian approach

$$IC = -2 \log L + 4 \sigma_{\log L}^2$$

where

$\log L$: average of the log-likelihood

$\sigma_{\log L}$: variance of the log-likelihood

\Rightarrow *The smaller the IC-value, the better the NP-model.*

Pull


Pull of an observable \mathcal{O}_i :

$$P(\mathcal{O}_i) = \left| \frac{\mathcal{O}_i^{\text{exp}} - \mathcal{O}_i^{\text{SM}}}{\sqrt{(\sigma_i^{\text{exp}})^2 + (\sigma_i^{\text{SM}})^2}} \right| - \left| \frac{\mathcal{O}_i^{\text{exp}} - \mathcal{O}_i^{\text{NP}}}{\sqrt{(\sigma_i^{\text{exp}})^2 + (\sigma_i^{\text{NP}})^2}} \right|$$

\Rightarrow *The larger the pull, the better the NP-model.*

LEP Searches

DM searches with mono-photon signatures at LEP:



DELPHI analyses

(arXiv:0406019, 0901.4486

[hep-ex])

& 1103.0240 [hep-ph]

$$\Rightarrow \frac{|\lambda_{12,13}|^2}{m_\phi^2} \lesssim \frac{1}{(175 \text{ GeV})^2}$$