



Contribution ID: 296

Type: Talk

## **【356】 UHECR Acceleration in FR-0 Jetted Active Galaxies**

*Wednesday, September 1, 2021 6:15 PM (15 minutes)*

Fanaroff Riley (FR) 0 radio galaxies form a low luminosity extension of the well-established ultrahigh energy cosmic ray (UHECR) accelerators FR-1 and FR-2 galaxies. Their higher number density makes them interesting candidate sources for an isotropic contribution to the observed UHECR flux. Here, acceleration and survival of UHECR in prevailing conditions of the FR-0 environment are discussed.

The photon target fields are composed of a jet and a host galaxy component, based on multi-wavelength data from the *FR0CAT*. This allows to simulate all relevant UHECRs loss processes.

We show that FR-0 galaxies can contribute to the UHECR flux in a hybrid scenario based on Fermi-I order and gradual shear acceleration.

**Authors:** Dr MERTEN, Lukas (University of Innsbruck); BOUGHELLIBA, Margot (University of Innsbruck); REIMER, Anita (University of Innsbruck); DA VELA, Paolo (University of Innsbruck); VOROBIOV, Serguei (University of Nova Gorica); TAVECCHIO, Fabrizio (Astronomical Observatory of Brera); BONNOLI, Giacomo (Instituto de Astrofísica de Andalucía); LUNDQUIST, Jon Paul (University of Nova Gorica); RIGHI, Chiara (Astronomical Observatory of Brera)

**Presenter:** Dr MERTEN, Lukas (University of Innsbruck)

**Session Classification:** Nuclear, Particle- & Astrophysics

**Track Classification:** Nuclear, Particle- and Astrophysics (FAKT - TASK)