



Contribution ID: 388

Type: Talk

[337] Increasing the ionization yield for the detection of ^{236}U and ^{233}U by AMS

Wednesday 1 September 2021 18:30 (15 minutes)

The detection efficiency of Accelerator Mass Spectrometry for long lived uranium isotopes (^{236}U or ^{233}U) is mainly limited by the rather low yield of the corresponding negative ions extracted from a caesium sputter ion source ($\approx 10^{-4}$). With our new sample preparation method environmental U is embedded in only 200 μg Fe_2O_3 matrix which is then mixed with PbF_2 . Extracting U as UF_5^- instead of UO^- yields an improvement in detection efficiency by more than a factor 10. UF_5^- extraction seems advantageous for the suppression of molecular isobaric background ($^{232}\text{ThH}^{3+}$, $^{235}\text{UH}^{3+}$) and allows operation at lower He stripper gas pressure.

Author: KERN, Michael (University of Vienna, Faculty of Physics - Isotope Physics)

Co-authors: GOLSER, Robin (University of Vienna, Faculty of Physics - Isotope Physics); HAIN, Karin (University of Vienna, Faculty of Physics - Isotope Physics); SCHMID, Clemens (University of Vienna, Faculty of Physics - Isotope Physics); STEIER, Peter (University of Vienna, Faculty of Physics - Isotope Physics); WIEDERIN, Andreas (Universität Wien, Faculty of Physics - Isotope Physics)

Presenter: KERN, Michael (University of Vienna, Faculty of Physics - Isotope Physics)

Session Classification: Nuclear, Particle- & Astrophysics

Track Classification: Nuclear, Particle- and Astrophysics (FAKT - TASK)