



Contribution ID: 392

Type: Talk

☒338☒ Determination of ^{135}Cs and ^{137}Cs by Accelerator Mass Spectrometry

Wednesday, September 1, 2021 6:45 PM (15 minutes)

The radionuclides ^{135}Cs and ^{137}Cs are present in the environment with an isotopic ratio $^{135,137}\text{Cs}/\text{Cs}$ ranging below 10^{-10} . The isotopic ratio $^{135}\text{Cs}/^{137}\text{Cs}$ can be used for source assessment of anthropogenic cesium input into the environment and finds applications in geology, nuclear forensics and oceanography. The combination of low concentration, low beta-decay energy and long half-life prevents the determination of ^{135}Cs via radiometric methods with the demanded sensitivity. Therefore, we established a measurement procedure for cesium by Accelerator Mass Spectrometry at VERA. First results on isobar suppression, abundance sensitivity and $^{135}\text{Cs}/^{137}\text{Cs}$ ratios of environmental samples will be presented.

Primary author: WIESER, Alexander (University of Vienna)

Co-authors: LACHNER, Johannes (University of Vienna); MARTSCHINI, Martin (Universität Wien); PRILLER, Alfred; STEIER, Peter (University of Vienna, Faculty of Physics - Isotope Physics); Mr ZOK, Dorian (Gottfried Wilhelm Leibniz Universität Hannover - Institut für Radioökologie und Strahlenschutz); GOLSER, Robin (University of Vienna, Faculty of Physics - Isotope Physics)

Presenter: WIESER, Alexander (University of Vienna)

Session Classification: Nuclear, Particle- & Astrophysics

Track Classification: Nuclear, Particle- and Astrophysics (FAKT - TASK)