

New Physics Beyond the Standard Model

Known knowns (= SM)
Known unknowns (e.g., DM)

Unknown unknowns

Lepton flavour violation in B decays?

$g_\mu - 2 ?$

John Ellis

KING'S
College
LONDON

$g_{\mu} - 2$:

from Dirac and Schwinger to Fermilab and Beyond



A story of 94 years,
8 experiments
and many theorists

John Ellis

KING'S
College
LONDON

It began with Dirac ...



- Two fundamental papers:

- “The quantum theory of the electron” (1928)

The Dirac equation: $(i\gamma_\mu \partial^\mu - m)\psi = 0$ predicts that the electron's magnetic moment $g = 2$

“That was really an unexpected bonus for me, completely unexpected”

- “The Quantum Theory of the Emission and Absorption of Radiation” (1927)

The basis for QED (and all of quantum field theory) enables the calculations of the anomaly: $g \neq 2$

The Quantum Theory of the Electron.

By P. A. M. DIRAC, St. John's College, Cambridge.

(Communicated by R. H. Fowler, F.R.S.—Received January 2, 1928.)

The new quantum mechanics, when applied to the problem of the structure of the atom with point-charge electrons, does not give results in agreement with experiment. The discrepancies consist of "duplexity" phenomena, the observed number of stationary states for an electron in an atom being twice the number given by the theory. To meet the difficulty, Goudsmit and Uhlenbeck have introduced the idea of an electron with a spin angular momentum of half a quantum and a magnetic moment of one Bohr magneton. This model for the electron has been fitted into the new mechanics by Pauli,* and Darwin,† working with an equivalent theory, has shown that it gives results in agreement with experiment for hydrogen-like spectra to the first order of accuracy.

The question remains as to why Nature should have chosen this particular model for the electron instead of being satisfied with the point-charge. One would like to find some incompleteness in the previous methods of applying quantum mechanics to the point-charge electron such that, when removed, the whole of the duplexity phenomena follow without arbitrary assumptions. In the present paper it is shown that this is the case, the incompleteness of the previous theories lying in their disagreement with relativity, or, alternately, with the general transformation theory of quantum mechanics. It appears that the simplest Hamiltonian for a point-charge electron satisfying the requirements of both relativity and the general transformation theory leads to an explanation of all duplexity phenomena without further assumption. All the same there is a great deal of truth in the spinning electron model, at least as a first approximation. The most important failure of the model seems to be that the magnitude of the resultant electron moving in an orbit in a central field model leads one to expect.

* Pauli, 'Z. f. Physik,' vol. 43, p. 601 (1927).

† Darwin, 'Roy. Soc. Proc.,' A, vol. 116, p. 227 (1927).

(2) Observations have been made of the electric fields and field changes associated with 18 distant and 5 near thunderstorms. The sudden changes of field due to distant lightning discharges (> 8 km.) were predominantly negative in sign, those due to near discharges (< 6 km.) predominantly positive. The relative frequencies of positive and negative changes were 1 : 5 in the former case and 4·3 : 1 in the latter. The steady electric fields below the 5 near storms were all strongly negative.

(3) It is shown that these results indicate that the thunderclouds were bi-polar in nature and that the polarity was generally, if not always, positive, the upper pole being positive and the lower pole negative. It is doubtful if any active storms of opposite polarity were observed at all.

(4) The electric moments of the charges removed by 82 lightning discharges have been measured. The mean value is 94 coulomb-kilometres.

The Quantum Theory of the Emission and Absorption of Radiation.

By P. A. M. DIRAC, St. John's College, Cambridge, and Institute for Theoretical Physics, Copenhagen.

(Communicated by N. Bohr, For. Mem. R.S.—Received February 2, 1927.)

§ 1. *Introduction and Summary.*

The new quantum theory, based on the assumption that the dynamical variables do not obey the commutative law of multiplication, has by now been developed sufficiently to form a fairly complete theory of dynamics. One can treat mathematically the problem of any dynamical system composed of a number of particles with instantaneous forces acting between them, provided it is describable by a Hamiltonian function, and one can interpret the mathematics physically by a quite definite general method. On the other hand, hardly anything has been done up to the present on quantum electrodynamics. The system in which the forces are propagated instantaneously, of the production of electron, and of the reaction of this field. In addition, there is a serious difficulty in making the theory satisfy all the requirements of the restricted

$$\left[p_0 + \frac{e}{c} A_0 + \rho_1 \left(\sigma, \mathbf{p} + \frac{e}{c} \mathbf{A} \right) + \rho_3 mc \right] \psi = 0$$

... and then Schwinger

- First calculation of leading-order contribution to $g - 2$ in QED: $\frac{\alpha}{2\pi}$ (1947)
- Inscribed on his tombstone



¹ Walke, Thompson, and Holt, *Phys. Rev.* **57**, 171 (1940).
² Solomon, Gould, and Anfinson, *Phys. Rev.* **72**, 1097 (1947).
³ Feather, *Proc. Camb. Phil. Soc.* **35**, 599 (1938).
⁴ Glendenin, *Nucleonica*, in press for January, 1948.
⁵ Marshall and Ward, *Can. J. Research* **15**, 29 (1939).
⁶ This result is in good agreement with a value of 250 kev, given in *Radioisotopes, Catalog and Price List No. 2*, revised September, 1947, distributed by Isotopes Branch, United States Atomic Energy Commission. Unfortunately, the Atomic Energy Commission's result is not supported by any published experimental evidence.

On Quantum-Electrodynamics and the Magnetic Moment of the Electron

JULIAN SCHWINGER
Harvard University, Cambridge, Massachusetts
 December 30, 1947

ATTEMPTS to evaluate radiative corrections to electron phenomena have heretofore been beset by divergence difficulties, attributable to self-energy and vacuum polarization effects. Electrodynamics unquestionably requires revision at ultra-relativistic energies, but is presumably accurate at moderate relativistic energies. It would be desirable, therefore, to isolate those aspects of the current theory that essentially involve high energies, and are subject to modification by a more satisfactory theory, from aspects that involve only moderate energies and are thus relatively trustworthy. This goal has been achieved by transforming the Hamiltonian of current hole theory electrodynamics to exhibit explicitly the logarithmically divergent self-energy of a free electron, which arises from

the virtual emission and absorption of light quanta. The electromagnetic self-energy of a free electron can be ascribed to an electromagnetic mass, which must be added to the mechanical mass of the electron. Indeed, the only meaningful statements of the theory involve this combination of masses, which is the experimental mass of a free electron. It might appear, from this point of view, that the divergence of the electromagnetic mass is unobjectionable, since the individual contributions to the experimental mass are unobservable. However, the transformation of the Hamiltonian is based on the assumption of a weak interaction between matter and radiation, which requires that the electromagnetic mass be a small correction ($\sim (e^2/hc)m_0$) to the mechanical mass m_0 .

The new Hamiltonian is superior to the original one in essentially three ways: it involves the experimental electron mass, rather than the unobservable mechanical mass; an electron now interacts with the radiation field only in the presence of an external field, that is, only an accelerated electron can emit or absorb a light quantum;⁶ the interaction energy of an electron with an external field is now subject to a *finite* radiative correction. In connection with the last point, it is important to note that the inclusion of the electromagnetic mass with the mechanical mass does not avoid all divergences; the polarization of the vacuum produces a logarithmically divergent term proportional to the interaction energy of the electron in an external field. However, it has long been recognized that such a term is equivalent to altering the value of the electron charge by a constant factor, only the final value being properly identified with the experimental charge. Thus the interaction between matter and radiation produces a renormalization of the electron charge and mass, all divergences being contained in the renormalization factors.

The simplest example of a radiative correction is that for the energy of an electron in an external magnetic field. The detailed application of the theory shows that the radiative correction to the magnetic interaction energy corresponds to an additional magnetic moment associated with the electron spin, of magnitude $\delta\mu/\mu = (\frac{1}{2}\pi)e^2/hc = 0.001162$. It is indeed gratifying that recently acquired experimental data confirm this prediction. Measurements on the hyperfine splitting of the ground states of atomic hydrogen and deuterium¹ have yielded values that are definitely larger than those to be expected from the directly measured nuclear moments and an electron moment of one Bohr magneton. These discrepancies can be accounted for by a small additional electron spin magnetic moment.² Recalling that the nuclear moments have been calibrated in terms of the electron moment, we find the additional moment necessary to account for the measured hydrogen and deuterium hyperfine structures to be $\delta\mu/\mu = 0.00126 \pm 0.00019$ and $\delta\mu/\mu = 0.00131 \pm 0.00025$, respectively. These values are not in disagreement with the theoretical prediction. More precise conformation is provided by measurement of the g values for the $^2S_{1/2}$, $^2P_{1/2}$, and $^2P_{3/2}$ states of sodium and gallium.³ To account for these results, it is necessary to ascribe the following additional spin magnetic moment to the electron, $\delta\mu/\mu = 0.00118 \pm 0.00003$.

$\mathcal{O}\left(\frac{\alpha}{\pi}\right)^2$: André Petermann

- Co-inventor (with Stueckelberg) of the renormalisation group
- First to submit a paper proposing quarks (a few days before Gell-Mann and Zweig), not widely known because written in French, and publication delayed > year!

8.B

Nuclear Physics 63 (1965) 349–352; © North-Holland Publishing Co., Amsterdam
Not to be reproduced by photoprint or microfilm without written permission from the publisher



PROPRIÉTÉS DE L'ÉTRANGETÉ ET UNE FORMULE DE MASSE POUR LES MÉSONS VECTORIELS

A. PETERMANN
CERN, Genève 23

Reçu le 30 décembre 1963

Abstract: A mass-formula for vector mesons is proposed, and the role of strangeness in mass-formulae discussed.

- Pioneer of $g_\mu - 2$ calculations: $a_\mu \equiv \frac{(g_\mu - 2)}{2} = 1 + \frac{\alpha}{2\pi} + 0.75\left(\frac{\alpha}{\pi}\right)^2$ (1957)
- First correct calculation of $\mathcal{O}(\alpha^2)$ contributions to $g_e - 2$, $g_\mu - 2$

(also Sommerfeld; Suura & Wichmann, previous work by Karplus & Kroll)

Fourth order magnetic moment of the electron

by A. Petermann.

CERN. Theoretical Study Division. Institute for theoretical Physics. Copenhagen.
(17. VIII. 1957.)

In connection with the upper and lower bounds analysis done by the author¹⁾, which indicated a clear discrepancy with the Karplus and Kroll's result for the 4th order magnetic moment²⁾, we have performed an analytic evaluation of the five independent diagrams contributing to this moment in fourth order³⁾. The results are the following:

$$\mu_I = \frac{\alpha^2}{\pi^2} \left(\frac{1}{6} + \frac{13}{36} \pi^2 + \frac{5}{4} \zeta(3) - \frac{5}{6} \pi^2 \text{Log } 2 \right) = -0.467 \frac{\alpha^2}{\pi^2}. \quad (1)$$

$$\mu_{II_a} = \frac{\alpha^2}{\pi^2} \left(\frac{11}{48} + \frac{\pi^2}{18} \right) = 0.778 \frac{\alpha^2}{\pi^2}. \quad (2)$$

$$\mu_{II_c} = \frac{\alpha^2}{\pi^2} \left(-\frac{67}{24} + \frac{\pi^2}{18} - \frac{1}{2} \zeta(3) + \frac{1}{3} \pi^2 \text{Log } 2 - \frac{1}{2} \text{Log } \frac{\lambda^2}{m^2} \right) = -0.564 \frac{\alpha^2}{\pi^2} - \frac{1}{2} \frac{\alpha^2}{\pi^2} \text{Log } \frac{\lambda^2}{m^2}. \quad (3)$$

$$\mu_{II_d} = \frac{\alpha^2}{\pi^2} \left(\frac{11}{24} - \frac{\pi^2}{18} + \frac{1}{2} \text{Log } \frac{\lambda^2}{m^2} \right) = -0.090 \frac{\alpha^2}{\pi^2} + \frac{1}{2} \frac{\alpha^2}{\pi^2} \text{Log } \frac{\lambda^2}{m^2}. \quad (4)$$

$$\mu_{II_e} = \frac{\alpha^2}{\pi^2} \left(\frac{119}{36} - \frac{\pi^2}{3} \right) = 0.016 \frac{\alpha^2}{\pi^2}. \quad (5)$$

$$\mu_{\text{total}}^{(4)} = \frac{\alpha^2}{\pi^2} \left(\frac{197}{144} + \frac{\pi^2}{12} + \frac{3}{4} \zeta(3) - \frac{1}{2} \pi^2 \text{Log } 2 \right) = -0.328 \frac{\alpha^2}{\pi^2}. \quad (6)$$

Compared with the values given in their original paper by KARPLUS and KROLL, one can see that two terms were in error: μ_I differs by

$$\frac{\alpha^2}{\pi^2} \frac{1}{32} = 0.031 \frac{\alpha^2}{\pi^2};$$

$$\mu_{II_c} \text{ by } \frac{\alpha^2}{\pi^2} \left(\frac{32}{3} - \frac{61}{8} \pi^2 + \frac{17}{2} \pi^2 \text{Log } 2 - \frac{109}{4} \zeta(3) \right) - 2.614 \frac{\alpha^2}{\pi^2}.$$

The three other terms check. The error in μ_I remained of course undetected in the upper and lower bound analysis owing to its small-

*) The terminology of ref. 2 is used throughout this paper.

ness. But the large discrepancy in μ_{II_c} was that pin-pointed out in the previous paper.

A summary of the most important electromagnetic observables, the theoretical values of which are modified by the new value of the magnetic moment, is now given:

$$\text{Moment of the electron: } \frac{\mu_e}{\mu_0} = 1.0011596 = 1 + \frac{\alpha}{2\pi} - 0.328 \frac{\alpha^2}{\pi^2}.$$

FRANKEN and LIEBES' value for it: $\mu_e/\mu_0 = 1.001167 \pm 0.000005^*$). g -factor of the μ -meson (electromagnetic):

$$2(1.0011654) = 2 \left(1 + \frac{\alpha}{2\pi} + 0.75 \frac{\alpha^2}{\pi^2} \right).$$

Last Lederman's value: $2(1.0021 \pm 0.0008)^*$.

$$2^2 S_{1/2} - 2^2 P_{1/2} \text{ (Hydrogen): } (1057.94 \pm 0.15) \text{ Mc/s; observed: } (1057.77 \pm 0.10) \text{ Mc/s.}$$

$$2^2 S_{1/2} - 2^2 P_{1/2} \text{ (Deuterium): } (1059.22 \pm 0.15) \text{ Mc/s; observed: } (1059.00 \pm 0.10) \text{ Mc/s.}$$

Fine structure constant: $1/\alpha = 137.0384$; (previously: 137.0365).

The new fourth order correction given here is in agreement with:

a) The upper and lower bounds given by the author¹⁾.

b) A calculation using a different method, performed by C. SOMMERFIELD³⁾.

c) A recalculation done by N. M. KROLL and collaborators⁴⁾.

The author thanks Prof. NIELS BOHR for the hospitality at the Institute.

References.

- 1) A. PETERMANN, Nuclear Physics **3**, 689 (1957) and Nuclear Physics in the press.
- 2) R. KARPLUS and N. M. KROLL, Phys. Rev. **77**, 536 (1950).
- 3) C. SOMMERFIELD, Phys. Rev. In the press.

*) Private Communication.

Lederman *et al.*: First Measurement of $g_\mu - 2$

PHYSICAL REVIEW

VOLUME 109, NUMBER 3

FEBRUARY 1, 1958

Magnetic Moment of the Free Muon*†

T. COFFIN, R. L. GARWIN,† S. PENMAN, L. M. LEDERMAN, AND A. M. SACHS

Columbia University,§ New York, New York

(Received October 1, 1957)

The magnetic moment of the positive μ meson has been measured in several target materials by a magnetic resonance technique. Muons were brought to rest with their spins parallel to a magnetic field. A radio-frequency pulse was applied to effect a spin reorientation which was detected by counting the decay electrons emerging after the pulse in a fixed direction. Results are expressed in terms of a g factor which for a spin $\frac{1}{2}$ particle is the ratio of the actual moment to $e\hbar/2m_\mu c$. The most accurate result obtained in a CHBr_3 target, is that $g = 2(1.0026 \pm 0.0009)$ compared to the theoretical prediction of $g = 2(1.0012)$. Less accurate measurements yielded $g = 2.005 \pm 0.005$ in a copper target and $g = 2.00 \pm 0.01$ in a lead target.

I. INTRODUCTION

THE μ meson has often been described as one of the more baffling of elementary particles. It alone, among the unstable particles, has no strong interaction. Aside from its usefulness as a tool in the study of nuclear structure and the details of parity violation in weak interactions it appears to play no essential role in any organization of fundamental particles. A precise measurement of the magnetic moment of the muon offers some promise for clarification of this situation.

The Dirac equation predicts precisely 2 for the g value of a spin $\frac{1}{2}$ particle. Including corrections due to the interaction of the particle with its radiation field, one obtains^{1,2}

$$g_\mu = 2 \left(1 + \frac{\alpha}{2\pi} + 0.75 \frac{\alpha^2}{\pi^2} + \dots \right) \quad (1)$$
$$= 2(1.0012).$$

an energy λ would alter the g value as follows



$$g = 2 \left\{ 1 + \left[1 - \frac{2}{3} \left(\frac{m_\mu}{\lambda} \right)^2 \right] \frac{\alpha}{2\pi} + \dots \right\}. \quad (2)$$

It might be remarked⁴ that the model used in reference 3 implies a modification in the scattering of one Dirac particle by another. Such a modification can be described by a mean square radius, the appropriate relation being $\langle r^2 \rangle_e = 6(\hbar/\lambda c)^2$. Qualitatively, at least, the measured proton radius should constitute an upper limit for such an "electrodynamic radius." Hence the fractional alteration of the muon moment from such a presumed breakdown of quantum electrodynamics should not exceed $\sim 0.02(\alpha/2\pi)$.

- Columbia Nevis and Carnegie Institute of Technology cyclotrons
- Agreement between theory and experiment

First $g_\mu - 2$ Experiment at CERN

(1958 - 1962)



$$\delta a \equiv \delta \left(\frac{g_\mu - 2}{2} \right) = \pm 0.4 \%$$


Georges Charpak

Francis Farley

Hans Sens

Theo Muller

Nino Zichichi




(Suggested by Leon Lederman)



(Also experiment at Berkeley Cyclotron)

Principle of Storage Ring Experiments


LIFE OF A MUON: THE g-2 EXPERIMENT




First Storage Ring Experiment at CERN

(1962 - 1968)

Design



Under construction



$$\delta a = \pm 270 \text{ ppm}$$

Agreement with theory after inclusion of light-by-light scattering
(Aldins, Kinoshita, Brodsky, Dufner)




(Kinoshita, 1967)

$\mathcal{O}\left(\frac{\alpha}{\pi}\right)^3$ Calculations



(Lautrup, 1968)
(+ De Rafael)



$$\begin{aligned}
 A_2^{(6)}(m_\mu/m_e) &= \frac{2}{9} \log^2 x - \left(\zeta(3) - \frac{2}{3} \pi^2 \log 2 + \frac{7\pi^2}{9} + \frac{31}{27} \right) \log x + \frac{97\pi^4}{360} \\
 &\quad - \frac{2}{9} \pi^2 \log^2 2 - \frac{8}{3} a_4 - \frac{\log^4 2}{9} - 6\zeta(3) + \frac{5}{3} \pi^2 \log 2 - \frac{85\pi^2}{18} + \frac{1219}{216} \\
 &\quad + x \left(-\frac{4}{3} \pi^2 \log x - \frac{604}{9} \pi^2 \log 2 + \frac{54079\pi^2}{1080} - \frac{13\pi^3}{18} \right) \\
 &\quad + x^2 \left[\frac{2}{3} \log^3 x + \left(\frac{\pi^2}{9} - \frac{10}{3} \right) \log^2 x + \left(\frac{16\pi^4}{135} + 4\zeta(3) - \frac{32\pi^2}{9} + \frac{194}{9} \right) \log x \right. \\
 &\quad \left. + \frac{4}{3} \zeta(3) \pi^2 - \frac{61\pi^4}{270} + \zeta(3) + \frac{197\pi^2}{36} - \frac{2809}{108} - \frac{14}{3} \pi^2 \log 2 \right] + \mathcal{O}(x^3) \\
 &= 22.868\,379\,98(20),
 \end{aligned}$$




(Kinoshita)

$\mathcal{O}\left(\frac{\alpha}{\pi}\right)^4$ Calculations





(Lautrup, 1972)



$$A_2^{(8)}(m_\mu/m_e) = 123.785\,51(44) + 8.8997(59) = 132.6852(60)$$

Second Storage Ring Experiment at CERN

(1969 - 1976)



Precession frequency

$$\vec{\omega}_a \equiv \vec{\omega}_s - \vec{\omega}_c = -\frac{q}{m_\mu} \left[a_\mu \vec{B} - a_\mu \left(\frac{\gamma}{\gamma+1} \right) (\beta \cdot \vec{B}) \vec{\beta} \right]$$

~~$\beta \cdot \mathbf{B} = 0$~~

Precession frequency $\propto a_\mu$
 $\delta a_\mu = 8 \text{ ppm}$

Magic energy
 $E = 3.094 \text{ GeV}$
 $\gamma = 29.3$

$$- \left(a_\mu - \frac{1}{\gamma^2 - 1} \right) \frac{\vec{\beta} \times \vec{E}}{c}$$

(Emilio Picasso \rightarrow LEP)

$g_\mu - 2$ in Supersymmetry

SPIN-ZERO LEPTONS AND THE ANOMALOUS MAGNETIC MOMENT OF THE MUON

John ELLIS, John HAGELIN and D.V. NANOPOULOS
CERN, Geneva, Switzerland

Received 14 June 1982

The anomalous magnetic moment of the muon $(g - 2)_\mu$ imposes constraints on the masses and mixing of spin-zero leptons (sleptons). We develop the predictions of models of spontaneous supersymmetry breaking for the slepton mass matrix, and show that they are comfortably consistent with the $(g - 2)_\mu$ constraints.

During the present resurgence of interest in supersymmetry broken at low energies [1] new significance is attached to the classical phenomenological playgrounds of gauge theories such as the anomalous magnetic moments of the electron and muon [2], flavour-changing neutral interactions [3,5] parity [6] and CP violation [7,8] in the strong interactions. The three latter phenomena make life rather difficult [3,7] for the most general form of soft supersymmetry breaking, whereas simple models [9-11] of spontaneously broken supersymmetry naturally [3,4,7] respect the $\Delta F \neq 0, P$ and CP violation constraints. As for the anomalous magnetic moments of the leptons, it has long been known that they vanish in an exactly supersymmetric theory [12], and Fayet [2] showed that in his model of supersymmetry breaking $(g - 2)_\mu$ would be compatible with experiment if the spin-zero muon (smuon) masses were heavier than 15 GeV. Direct experimental searches [13] now exclude the existence of lighter smuons. Fayet's analysis [2] was in the context of a model with a very light photino $\tilde{\gamma}$ (see fig. 1a), and Grifols and Méndez [14] have recently made the interesting observation that his analysis is significantly altered for massive gauginos (see figs. 1b, 1c). They show that there are potentially nontrivial constraints on the smuon masses in models of broken supersymmetry.




Fig. 1. One-loop diagrams contributing to $(g - 2)_\mu$: (a) essentially massless photino ($\tilde{\gamma}$) exchange, (b) \tilde{W} and sneutrino ($s\nu$) exchange, and (c) \tilde{B} or \tilde{Z} exchange.

right transition operator there is a GIM [15]-like cancellation between the smuon mass eigenstates in fig. 1c which provides a potential suppression mechanism. We analyze recent models [10,11] of spontaneous supersymmetry breaking originating in the D and F sectors, respectively. We show that in the former case $(g - 2)_\mu$ is suppressed by near degeneracy between the smuon mass eigenstates, while in the latter case $(g - 2)_\mu$ is suppressed by small mixing angles between the left- and right-handed smuons. We close with some remarks about $(g - 2)_e$ and about parity violation in the strong interactions.

When they examined figs. 1a, 1b and 1c, Grifols and Méndez [14] realized that there was a fundamental difference between the (almost ?) massless $\tilde{\gamma}$ diagram of fig. 1a and the \tilde{W} diagram of fig. 1b as compared to the massive \tilde{B} or \tilde{Z} diagram of fig. 1c. The

- One-loop contribution from smuon/neutralino loop

$$\Delta(g - 2)_\mu = -ab(\cos \alpha \sin \alpha / 4\pi^2)(m_\mu / m_{\tilde{G}})$$

$$\times \{1/(1 - \eta_1) + 2\eta_1/(1 - \eta_1)^2$$


$$+ [2\eta_1/(1 - \eta_1)^3] \log \eta_1 - (\eta_1 \leftrightarrow \eta_2)\},$$

- where $\eta_i \equiv (m_{s\mu_i}^2 / m_{\tilde{G}}^2)$


- and $\mathcal{L} = a\sqrt{2} s_\mu \bar{\mu}_L \tilde{G} + b\sqrt{2} t_\mu \bar{\mu}_R \tilde{G}$

BNL Experiment

(1984 - 2003)



Possible Discrepancy with Theory?



$$\delta a = \pm 0.47 \text{ ppm}$$

$g_\mu - 2$ in Supersymmetry v2: the CMSSM

Combining the muon anomalous magnetic moment with other constraints on the CMSSM

John Ellis^a, D.V. Nanopoulos^{b,c,d}, Keith A. Olive^{a,e}

^a TH Division, CERN, Geneva, Switzerland

^b Department of Physics, Texas A&M University, College Station, TX 77843, USA


^c Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, TX 77381, USA

^d Chair of Theoretical Physics, Academy of Athens, Division of Natural Sciences, 28 Panepistimiou Avenue, Athens 10679, Greece


^e Theoretical Physics Institute, School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA

Received 16 March 2001; accepted 10 April 2001

Editor: R. Gatto



Sparticle masses a few hundred GeV



$\tan \beta$

(2001)

Abstract

We combine the constraint suggested by the recent BNL E821 measurement of the anomalous magnetic moment of the muon on the parameter space of the constrained MSSM (CMSSM) with those provided previously by LEP, the measured rate of $b \rightarrow s\gamma$ decay and the cosmological relic density $\Omega_\chi h^2$. Our treatment of $\Omega_\chi h^2$ includes carefully the direct-channel Higgs poles in annihilation of pairs of neutralinos χ and a complete analysis of $\chi - \tilde{\ell}$ coannihilation. We find excellent consistency between all the constraints for $\tan \beta \gtrsim 10$ and $\mu > 0$, for restricted ranges of the CMSSM parameters m_0 and $m_{1/2}$. All the preferred CMSSM parameter space is within reach of the LHC, but may not be accessible to the Tevatron collider, or to a first-generation e^+e^- linear collider with centre-of-mass energy below 1.2 TeV. © 2001 Published by Elsevier Science B.V.

$\mathcal{O}\left(\frac{\alpha}{\pi}\right)^5$ Calculations

Complete Tenth-Order QED Contribution to the Muon $g - 2$

Tatsumi Aoyama,^{1,2} Masashi Hayakawa,^{3,2} Toichiro Kinoshita,^{4,2} and Makiko Nio²

¹*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI), Nagoya University, Nagoya, 464-8602, Japan*

²*Nishina Center, RIKEN, Wako, Japan 351-0198*



³*Department of Physics, Nagoya University, Nagoya, Japan 464-8602*

⁴*Laboratory for Elementary Particle Physics, Cornell University, Ithaca, New York, 14853, U.S.A*

(Dated: August 21, 2012)


We report the result of our calculation of the complete tenth-order QED terms of the muon $g - 2$. Our result is $a_\mu^{(10)} = 753.29 (1.04)$ in units of $(\alpha/\pi)^5$, which is about 4.5 s.d. larger than the leading-logarithmic estimate 663 (20). We also improved the precision of the eighth-order QED term of a_μ , obtaining $a_\mu^{(8)} = 130.8794 (63)$ in units of $(\alpha/\pi)^4$. The new QED contribution is $a_\mu(\text{QED}) = 116\,584\,718\,951 (80) \times 10^{-14}$, which does not resolve the existing discrepancy between the standard-model prediction and measurement of a_μ .

PACS numbers: 13.40.Em,14.60.Ef,12.20.Ds



(2012)


Electroweak Contributions



$$a_{\mu}^{\text{EW}(1)} = \frac{G_{\text{F}}}{\sqrt{2}} \frac{m_{\mu}^2}{8\pi^2} \left[\frac{5}{3} + \frac{1}{3}(1 - 4s_{\text{W}}^2)^2 \right] = 194.79(1) \times 10^{-11}$$


- Leading one-loop order

- Two-loop contributions




$$\begin{aligned} a_{\mu}^{\text{EW}(2), \text{logs}} &= -4 \frac{\alpha}{\pi} \log \frac{M_{\text{Z}}}{m_{\mu}} a_{\mu}^{\text{EW}(1)} \\ &+ \frac{G_{\text{F}} m_{\mu}^2}{8\pi^2 \sqrt{2}} \frac{\alpha}{\pi} \log \frac{M_{\text{Z}}}{m_{\mu}} \left[-\frac{47}{9} - \frac{11}{9}(1 - 4s_{\text{W}}^2)^2 \right] \\ &+ \frac{G_{\text{F}} m_{\mu}^2}{8\pi^2 \sqrt{2}} \frac{\alpha}{\pi} \sum_f \log \frac{M_{\text{Z}}}{\max(m_f, m_{\mu})} \left[-6g_{\text{A}}^{\mu} g_{\text{A}}^f N_f Q_f^2 + \frac{4}{9} g_{\text{V}}^{\mu} g_{\text{V}}^f N_f Q_f \right] \\ &= -41.2(1.0) \times 10^{-11} \end{aligned}$$

Combined result




Hadronic Contribution to Light-by-Light Scattering



Contribution	PdRV(09) [475]	N/JN(09) [476, 596]	J(17) [27]	Our estimate
π^0, η, η' -poles	114(13)	99(16)	95.45(12.40)	93.8(4.0)
π, K -loops/boxes	-19(19)	-19(13)	-20(5)	-16.4(2)
S -wave $\pi\pi$ rescattering	-7(7)	-7(2)	-5.98(1.20)	-8(1)
subtotal	88(24)	73(21)	69.5(13.4)	69.4(4.1)
scalars	-	-	-	} - 1(3)
tensors	-	-	1.1(1)	
axial vectors	15(10)	22(5)	7.55(2.71)	
u, d, s -loops / short-distance	-	21(3)	20(4)	15(10)
c -loop	2.3	-	2.3(2)	3(1)
total	105(26)	116(39)	100.4(28.2)	92(19)

Theory Initiative

- Comprehensive review of calculations of the Standard Model contributions to $g_\mu - 2$
- Including discussion of the uncertainties
- Particularly in calculation of leading-order vacuum polarisation



Aoyama et al, arXiv:2006.04822



The anomalous magnetic moment of the muon in the Standard Model

T. Aoyama^{1,2,3}, N. Asmussen⁴, M. Benayoun⁵, J. Bijnens⁶, T. Blum^{7,8}, M. Bruno⁹, I. Caprini¹⁰, C.M. Carloni Calame¹¹, M. Cè^{9,12,13}, G. Colangelo^{14,*}, F. Curciarello^{15,16}, H. Czyż¹⁷, I. Danilkin¹², M. Davier^{18,*}, C.T.H. Davies¹⁹, M. Della Morte²⁰, S.I. Eidelman^{21,22,*}, A.X. El-Khadra^{23,24,*}, A. Gérardin²⁵, D. Giusti^{26,27}, M. Golterman²⁸, Steven Gottlieb²⁹, V. Gülpers³⁰, F. Hagelstein¹⁴, M. Hayakawa^{31,2}, G. Herdoíza³², D.W. Hertzog³³, A. Hoecker³⁴, M. Hoferichter^{14,35,*}, B.-L. Hoid³⁶, R.J. Hudspith^{12,13}, F. Ignatov²¹, T. Izubuchi^{37,8}, F. Jegerlehner³⁸, L. Jin^{7,8}, A. Keshavarzi³⁹, T. Kinoshita^{40,41}, B. Kubis³⁶, A. Kupich²¹, A. Kupś^{42,43}, L. Laub¹⁴, C. Lehner^{26,37,*}, L. Lellouch²⁵, I. Logashenko²¹, B. Malaescu⁵, K. Maltman^{44,45}, M.K. Marinković^{46,47}, P. Masjuan^{48,49}, A.S. Meyer³⁷, H.B. Meyer^{12,13}, T. Mibe^{1,*}, K. Miura^{12,13,3}, S.E. Müller⁵⁰, M. Nio^{2,51}, D. Nomura^{52,53}, A. Nyffeler^{12,*}, V. Pascalutsa¹², M. Passera⁵⁴, E. Perez del Rio⁵⁵, S. Peris^{48,49}, A. Portelli³⁰, M. Procura⁵⁶, C.F. Redmer¹², B.L. Roberts^{57,*}, P. Sánchez-Puertas⁴⁹, S. Serednyakov²¹, B. Schwartz²¹, S. Simula²⁷, D. Stöckinger⁵⁸, H. Stöckinger-Kim⁵⁸, P. Stoffer⁵⁹, T. Teubner^{60,*}, R. Van de Water²⁴, M. Vanderhaeghen^{12,13}, G. Venanzoni⁶¹, G. von Hippel¹², H. Wittig^{12,13}, Z. Zhang¹⁸, M.N. Achasov²¹, A. Bashir⁶², N. Cardoso⁴⁷, B. Chakraborty⁶³, E.-H. Chao¹², J. Charles²⁵, A. Crivellin^{64,65}, O. Deineka¹², A. Denig^{12,13}, C. DeTar⁶⁶, C.A. Dominguez⁶⁷, A.E. Dorokhov⁶⁸, V.P. Druzhinin²¹, G. Eichmann^{69,47}, M. Fael⁷⁰, C.S. Fischer⁷¹, E. Gámiz⁷², Z. Gelzer²³, J.R. Green⁹, S. Guellati-Khelifa⁷³, D. Hatton¹⁹, N. Hermansson-Truedsson¹⁴, S. Holz³⁶, B. Hörz⁷⁴, M. Knecht²⁵, J. Koponen¹, A.S. Kronfeld²⁴, J. Laiho⁷⁵, S. Leupold⁴², P.B. Mackenzie²⁴, W.J. Marciano³⁷, C. McNeile⁷⁶, D. Mohler^{12,13}, J. Monnard¹⁴, E.T. Neil⁷⁷, A.V. Nesterenko⁶⁸, K. Ottnad¹², V. Pauk¹², A.E. Radzhabov⁷⁸, E. de Rafael²⁵, K. Raya⁷⁹, A. Risch¹², A. Rodríguez-Sánchez⁶, P. Roig⁸⁰, T. San José^{12,13}, E.P. Solodov²¹, R. Sugar⁸¹, K. Yu. Todyshev²¹, A. Vainshtein⁸², A. Vaquero Avilés-Casco⁶⁶, E. Weil⁷¹, J. Wilhelm¹², R. Williams⁷¹, A.S. Zhevlakov⁷⁸

¹ Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan

² Nishina Center, RIKEN, Wako 351-0198, Japan

³ Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI), Nagoya University, Nagoya 464-8602, Japan

⁴ School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, United Kingdom

⁵ LPNHE, Sorbonne Université, Université de Paris, CNRS/IN2P3, Paris, France

* Corresponding authors.

E-mail address: MUON-GM2-THEORY-SC@fnal.gov (G. Colangelo, M. Davier, S.I. Eidelman, A.X. El-Khadra, M. Hoferichter, C. Lehner, T. Mibe, A. Nyffeler, B.L. Roberts, T. Teubner).

<https://doi.org/10.1016/j.physrep.2020.07.006>


0370-1573/© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Hadronic Vacuum Polarization

- Most important contribution is from low energies $\lesssim 1$ GeV, dominated by ρ and ω peaks, taking account of interference effects
- Uncertainties dominated by ρ and ω region, and by region between 1 and 2 GeV (ϕ , etc.)
- High energies under good control from perturbative QCD

$$\begin{aligned}
 a_{\mu}^{\text{HVP, LO}} &= 693.1(2.8)_{\text{exp}}(2.8)_{\text{sys}}(0.7)_{\text{DV+QCD}} \times 10^{-10} \\
 &= 693.1(4.0) \times 10^{-10}.
 \end{aligned}$$

Aoyama et al, arXiv:2006.04822





τ Decays?

- Relation between τ decays and $I = 1$ portion of hadronic vacuum polarization:


$$\sigma_{X^0}^{I=1}(s) = \frac{4\pi\alpha^2}{s} v_{1,X^-}(s)$$

$$v_{1,X^-}(s) = \frac{m_\tau^2}{6|V_{ud}|^2} \frac{\mathcal{B}_{X^-}}{\mathcal{B}_e} \frac{1}{N_X} \frac{dN_X}{ds} \times \left[\left(1 - \frac{s}{m_\tau^2}\right)^2 \left(1 + \frac{2s}{m_\tau^2}\right) \right]^{-1} \frac{R_{IB}(s)}{S_{EW}}$$


- **BUT** what about $I = 0$ portion?
- **AND** what about isospin breaking?
- **AND** uncertainties in τ decay data?
- **NOT INCLUDED** by Theory Initiative



Comparison of Calculations of Hadronic Vacuum Polarization





$$a_{\mu}^{\text{HVP}} + [a_{\mu}^{\text{QED}} + a_{\mu}^{\text{Weak}} + a_{\mu}^{\text{HLbL}}] \rightarrow a_{\mu}^{\text{SM}}$$



RBC/UKQCD Hybrid Method




Replace lattice data at very short and long distances
by experimental e+e- scattering data

- Convert R-ratio data to Euclidean correlation function (via the dispersive integral) and compare with lattice results for windows in Euclidean time
- intermediate window:
expect reduced FV effects and discretization errors



BUT



BMW Lattice Calculation



High statistics, accurate continuum extrapolation


$$a_{\mu}^{\text{EXP}} - a_{\mu}^{\text{BMW}} = 107(70) \times 10^{-11}$$

How to Accommodate BMW?




- Analyticity and unitarity constrain increase in $\pi^+\pi^-$ cross section < 1 GeV
- Maximum allowed conflicts with data, does not change greatly prediction for a_μ
- Increase in cross section at higher energies affects electroweak observables

Fermilab Experiment



Does the magnet look familiar?

Fermilab Data




Fermilab Measurement

FNAL result: $a_\mu(\text{FNAL}) = 116\,592\,040(54) \times 10^{-11}$ (0.46 ppm)

Combined result: $a_\mu(\text{Exp}) = 116\,592\,061(41) \times 10^{-11}$ (0.35 ppm)

Difference from Standard Model: $a_\mu(\text{Exp}) - a_\mu(\text{SM}) = (251 \pm 59) \times 10^{-11}$





Interpretation Papers

2104.05685	Vector LQ	B	Du		
5656	$L_\mu - L_\tau$	DM	Borah		
5006	$B_q - L_\mu$	B	Cen		Leptoquarks
4494	LFV	LFV	Li		
4503	Pseudoscalar	DM, H decays	Lu		Extra U(1)
4456	2HDM	DM	Arcadi		
3542	B-LSSM	H decays	Yang		Extra Higgs
3701	Leptophilic spin 0	H factory	Chun		
3839	SUSY	HL-LHC	Aboubrahim		Supersymmetry
3691	Survey	DM, LHC	Athron		
3705	Seesaw	g_e	Escribano		Axion
3699	Gauged 2HDM	B	Chen		
3239	SUSY	Gravitino DM	Gu		
3284	NMSSM	DM	Cao		
3262	GUT-constrained SUSY	DM, LHC	Wang		
3292	MSSM	CPV	Han		
3296	lepton mass matrix	Flavour	Calibbi		
3280	Z_d	Cs weak charge	Cadeddu		
3334	E_6 3-3-1	H stability	Li		
3242	μ - τ -philic H	τ decays, LHC	Wang		
3259	Anomaly mediation	DM	Yin		
3245	pMSSM	DM, fine-tuning	Van Beekveld		
3274	NMSSM	DM, AMS-02 pbar	Abdughani		
3290	MSSM	DM	Cox		
3367	2HDM	V-like leptons	Ferreira		
3267	Axion	Low-scale	Buen-Abad		
3340	$L_\mu - L_\tau$	AMS-02 positrons	Zu		
3282	ALP	V-like fermions	Brdar		
3301	Lepton portal	DM	Bai		
3276	Dark axion portal	Dark photon	Ge		
3491	GmSUGRA	LHC	Ahmed		
3227	2HDM	LHC	Han		
3302	SUSY	small μ	Baum		
3238	Scalar	DM, p radius	Zhu		
3489	μ ν SSM	B, H decays	Zhang		
3287	pMSSM	ILC	Chakraborti		
3228	DM	B, H decays	Arcadi		

890	Radiative seesaw				Chiang
2103.13991	Scalar LQ	B, H decays			Greljo
2012.11766	DM				D'Agnolo
2012.07894	Axions				Darmé
1812.06851	Charmphilic LQ				Kowalska
2104.04458	GUT-constrained SUSY	DM			Chakraborti
5730	LQ + charged singlet	B, Cabibbo			Marzocca
6320	L-R symmetry				Boyarkin
6858	$L_\mu - L_\tau$	ν masses			Zhou
6854	D-brane	U(1), Regge			Anchordoqui
6656	vector LQ	B			Ban
7597	SUSY	LHC, landscape			Baer
7047	3HDM	Fermion masses			Carcamo
7680	Leptophilic Z'	Global analysis			Buras
8289	Custodial symmetry	Light scalar + pseudoscalar			Balkin
9205	U(1)D	Neutrino mass			Dasgupta
8819	Lepton non-universality	Naturalness			Cacciapaglia
8640	$2 \times 2 \times 1$	Higgses, heavy nus			Boyarkina
8293	Multi-TeV sleptons in FSSM	Extended H, tau decays			Altmannshofer
10114	SO(10)	Yukawa unification			Aboubrahim
7681	U(1)B-L	DUNE			Dev
10324	Gauged lepton number	Dark matter			Ma
10175	2HDM	Lighter Higgs?			Jueid
11229	LQ	Matter unification			Fileviez
15136	U(1)	HE neutrinos, H tension			Alonso
2105.00903	Anomalous 3-boson vertex	W mass			Arbuzov
7655	U(1)T3R	RK(*)			Dutta
8670	Leptoquark	ν mass, LFV			Zhang



Scalar Leptoquarks

- Consider two types of leptoquarks with couplings



$$-\mathcal{L}_{QL}^Y \supset Y_2 Q_L \Phi_3 (\nu^c)_L + Y_2 \ell_L \Phi_4 (u^c)_L + Y_4 \ell_L \Phi_3^\dagger (d^c)_L + Y_4 Q_L \Phi_4^\dagger (e^c)_L + \text{h.c.}$$



- Consider constraints from $B_s \rightarrow \mu^+\mu^-$, $\mathcal{R}_K, \mathcal{R}_{K^*}$



Scalar Leptoquarks

- Consider 2 scenarios for mixing between leptoquarks:


$$\begin{aligned}
 -\mathcal{L} \supset & \bar{e}^i \left(-\sin \theta_{LQ} c_L^{ij} P_L + \cos \theta_{LQ} c_R^{ij} P_R \right) d^j \phi_A^{-2/3} \\
 & + \bar{e}^i \left(\cos \theta_{LQ} c_L^{ij} P_L + \sin \theta_{LQ} c_R^{ij} P_R \right) d^j \phi_B^{-2/3} + \text{h.c.}
 \end{aligned}$$



- Constraints from $g_\mu - 2$, \mathcal{R}_K , $B_s \rightarrow \mu^+ \mu^-$

Leptophilic Z' Gauge Boson

- LHC sets strong bounds only if Z' boson couples to quarks
- Weaker constraints on Z' bosons coupled to leptons only
- $\ell \rightarrow \ell' \nu \bar{\nu}$, $\ell \rightarrow \ell' \gamma$, $\ell \rightarrow 3\ell'$, mixing with Z and anomalous magnetic moments
- Can explain $g_\mu - 2$ with Z' coupled to $L_\mu - L_\tau$
- Search for lepton universality violation in $\tau \rightarrow \mu \nu \bar{\nu} / \tau \rightarrow e \nu \bar{\nu}$




$$\frac{\mathcal{A}[\tau \rightarrow \mu \nu \bar{\nu}]}{\mathcal{A}[\mu \rightarrow e \nu \bar{\nu}]_{\text{EXP}}} = 1.0029 \pm 0.0014$$

$$\frac{\mathcal{A}[\tau \rightarrow \mu \nu \bar{\nu}]}{\mathcal{A}[\tau \rightarrow e \nu \bar{\nu}]_{\text{EXP}}} = 1.0018 \pm 0.0014$$

$$\frac{\mathcal{A}[\tau \rightarrow e \nu \bar{\nu}]}{\mathcal{A}[\mu \rightarrow e \nu \bar{\nu}]_{\text{EXP}}} = 1.0010 \pm 0.0014$$

Leptophilic Z' Gauge Boson

- Scenario with no $Z - Z'$ mixing, left- and right-handed couplings to μ, τ only



$g_\mu - 2$ in Supersymmetry



- Muon ψ_f , 4 neutralinos ψ_i , 2 smuons ϕ_k ($\tilde{\mu}_{L,R}$)




$$- \mathcal{L}_{int} = \sum_{ik} \bar{\psi}_f \left(K_{ik} \frac{1 - \gamma_5}{2} + L_{ik} \frac{1 + \gamma_5}{2} \right) \psi_i \phi_k + H.c.$$

- One-loop contributions from smuon/neutralino loops:

- Left-right mixing: $a_f^{11} = \sum_{ik} \frac{m_f}{8\pi^2 m_i} \text{Re}(K_{ik} L_{ik}^*) I_1\left(\frac{m_f^2}{m_i^2}, \frac{m_k^2}{m_i^2}\right)$

- Unmixed: $a_f^{12} = \sum_{ik} \frac{m_f^2}{16\pi^2 m_i^2} (|K_{ik}|^2 + |L_{ik}|^2) I_2\left(\frac{m_f^2}{m_i^2}, \frac{m_k^2}{m_i^2}\right)$



$g_\mu - 2$ in Phenomenological Supersymmetry (pMSSM11)



No relation between squark & gluino masses and sleptons and neutralino
 No problem accommodating BNL/FNAL result
 Neutralino DM, smuon masses $\sim 300/400$ GeV

LHC vs Supersymmetry


- LHC does not exclude (relatively) light electroweakly-interacting particles, e.g., sleptons



- LHC favours squarks & gluinos $> 2 \text{ TeV}$ (but loopholes)


Supersymmetry

- g_μ – 2-friendly scenario with light neutralino, chargino & slepton





- **Red star** points include all relevant LHC and direct scattering constraints
- Prospects for the ILC?

$g_\mu - 2$ in Supersymmetric SU(5) GUT (CMSSM)



Scenario relates squark & gluino masses
to sleptons and neutralino
Cannot accommodate BNL/FNAL result
Smuon masses $\gtrsim 4$ TeV



$g_\mu = 2$ in Flipped SU(5) GUT

- Extend GUT SU(5) with additional U(1)

- “Flipped” fermion assignments to representations:

$$\bar{f}_i(\bar{\mathbf{5}}, -3) = \{U_i^c, L_i\} \quad , \quad F_i(\mathbf{10}, 1) = \{Q_i, D_i^c, N_i^c\} \quad , \quad l_i(\mathbf{1}, 5) = E_i^c \quad , \quad i = 1, 2, 3$$

- Break GUT symmetry with 10-dimensional Higgses, electroweak symmetry with 5-dimensional Higgses:

$$H(\mathbf{10}, 1) = \{Q_H, D_H^c, N_H^c\} \quad , \quad \bar{H}(\bar{\mathbf{10}}, -1) = \{\bar{Q}_H, \bar{D}_H^c, \bar{N}_H^c\}$$

$$h(\mathbf{5}, -2) = \{T_{H_c}, H_d\} \quad , \quad \bar{h}(\bar{\mathbf{5}}, 2) = \{\bar{T}_{\bar{H}_c}, H_u\}$$


- Superpotential:

$$W = \lambda_1^{ij} F_i F_j h + \lambda_2^{ij} F_i \bar{f}_j \bar{h} + \lambda_3^{ij} \bar{f}_i \ell_j^c h + \lambda_4 H H h + \lambda_5 \bar{H} \bar{H} \bar{h} \\ + \lambda_6^{ia} F_i \bar{H} \phi_a + \lambda_7^a h \bar{h} \phi_a + \lambda_8^{abc} \phi_a \phi_b \phi_c + \mu_\phi^{ab} \phi_a \phi_b \quad ,$$


- Scan free parameters of model:

$$M_5, M_{X1}, m_{10}, m_5, m_1, \mu, M_A, A_0, \tan \beta$$

$g_\mu = 2$ in Flipped SU(5)



$g_\mu - 2$ in CMSSM & Flipped SU(5) vs Lattice, Data-Driven Calculation



Δa_μ ($\times 10^{11}$): GUT models vs Standard Model calculations

$g_\mu = 2$ in Flipped SU(5)

Parameters & predictions at best-fit point



Input GUT parameters (masses in units of 10^{16} GeV)		
$M_{GUT} = 1.00$	$M_X = 0.79$	$V = 1.13$
$\lambda_4 = 0.1$	$\lambda_5 = 0.3$	$\lambda_6 = 0.001$
$g_5 = 0.70$	$g_X = 0.70$	$m_{\nu_3} = 0.05$ eV
Input supersymmetry parameters (masses in GeV units)		
$M_5 = 2460$	$M_1 = 240$	$\mu = 4770$
$m_{10} = 930$	$m_{\bar{5}} = 450$	$m_1 = 0$
$M_A = 2100$	$A_0/M_5 = 0.67$	$\tan \beta = 35$
MSSM particle masses (in GeV units)		
$m_\chi = 84$	$m_{\tilde{t}_1} = 4030$	$m_{\tilde{g}} = 5090$
$m_{\chi_2} = 2160$	$m_{\chi_3} = 5080$	m_{χ_4}
$m_{\tilde{\mu}_R} = 101$	$m_{\tilde{\mu}_L} = 1600$	$m_{\tilde{\tau}_1}$
$m_{\tilde{q}_L} = 4470$	$m_{\tilde{d}_R} = 4250$	$m_{\tilde{u}_R}$
$m_{\tilde{t}_2} = 4410$	$m_{\tilde{b}_1} = 4170$	$m_{\tilde{b}_2}$
$m_{\chi^\pm} = 2160$	$m_{H,A} = 2100$	$m_{H^\pm} = 2100$
Other observables		
$\Delta a_\mu = 150 \times 10^{-11}$	$\Omega_\chi h^2 = 0.13$	$m_h = 122$ GeV
Normal-ordered ν masses:	$\tau_{p \rightarrow e^+ \pi^0} _{\text{NO}} = 1.1 \times 10^{36}$ yrs	$\tau_{p \rightarrow \mu^+ \pi^0} _{\text{NO}} = 1.1 \times 10^{37}$ yrs
Inverse-ordered ν masses:	$\tau_{p \rightarrow e^+ \pi^0} _{\text{IO}} = 3.2 \times 10^{37}$ yrs	$\tau_{p \rightarrow \mu^+ \pi^0} _{\text{IO}} = 2.3 \times 10^{36}$ yrs

Opportunities to search for light smuon, neutralino at LHC
Other sparticles too heavy?


MuonE: Proposed CERN Experiment to Measure HVP in Space-Like Region

Scattering of 150 GeV muons on electrons at CERN SPS

$$a_{\mu}^{\text{HLO}} = \frac{\alpha}{\pi} \int_0^1 dx (1-x) \Delta\alpha_{\text{had}}[t(x)] \quad \alpha(t) = \frac{\alpha(0)}{1 - \Delta\alpha(t)} \quad t(x) = -\frac{x^2 m_{\mu}^2}{1-x} < 0$$





J-PARC Experiment



Different technique: ultra-cold muon beam from muonium, accelerate to 300 MeV, inject into storage ring with radius 66cm

Magnetic Dipole Moment of the Electron

- Discrepancies between determinations of α from atomic measurements and $a_e \equiv (g_e - 2)/2 + \text{QED}$
- New determination of α via rubidium recoil measurement allows BSM contribution to a_e in range $-3.4 \times 10^{-13} < \delta a_e < 9.8 \times 10^{-13}$, comparable to $\delta a_\mu \times (m_e/m_\mu)^2$
- Experiment underway to improve precision on a_e by order of magnitude



Quo Vadis $g_\mu - 2$?

- **Never forget:** the (near-) consistency between theory and experiment for $g_\mu - 2$ (and $g_e - 2$) is among the greatest successes of particle physics, particularly quantum field theory
- **Need no reminder:** the discrepancy between theory and experiment for $g_\mu - 2$ may be a window on physics beyond the Standard Model
- Still some **debate** about Standard Model calculation (lattice?)
- **Plenty** of theoretical interpretations proposed: many possible connections to other physics areas (B decays, dark matter, ...)
- **More experimental results** on the way: FNAL, J-PARC, a_e , MuonE, ...
- **We live in interesting times!**

Summary

Visible matter

Standard Model

$$g_{\mu} - 2?$$