Annual Modulation Results from DAMA/LIBRA

R. Cerulli
INFN-Roma Tor Vergata

ICNFP 2021
Kolymbari, Crete, Greece
Aug 23 – Sept 2, 2021

DAMA set-ups

an observatory for rare processes @ LNGS

- DAMA/LIBRA (DAMA/NaI)
- DAMA/LXe
- DAMA/R&D
- DAMA/Crys
- DAMA/Ge

Collaboration:

Roma Tor Vergata, Roma La Sapienza, LNGS, IHEP/Beijing
+ by-products and small scale expts.: INR-Kiev
+ neutron meas.: ENEA-Frascati, ENEA-Casaccia
+ in some studies on ββ decays (DST-MAE project): IIT Kharagpur and Ropar, India

Web Site: http://people.roma2.infn.it/dama

Some direct detection processes:

- Scatterings on nuclei

  → detection of nuclear recoil energy



- Inelastic Dark Matter: **W + N → W* + N**

  → W has 2 mass states χ+ , χ- with δ mass splitting

  → Kinematical constraint for the inelastic scattering of χ- on a nucleus

$$\frac{1}{2}\mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Excitation of bound electrons in scatterings on nuclei

  → detection of recoil nuclei + e.m. radiation

- Conversion of particle into e.m. radiation

  → detection of γ, X-rays, e^-



- Interaction only on atomic electrons

  → detection of e.m. radiation



... even WIMPs

- Interaction of light DMp (LDM) on e^- or nucleus with production of a lighter particle

  → detection of electron/nucleus recoil energy

e.g. sterile ν



e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the e.m. component of their rate

… also other ideas …

The annual modulation: a model independent signature for the investigation of DM particles

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions can point out its presence.

Requirements:

1) Cosine-like modulation of the rate
2) In low energy range
3) Period of 1 year
4) Phase at about June 2nd
5) For single-hit events in a multi-detector set-up
6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios

Drukier, Freese, Spergel PRD86; Freese et al. PRD88



$v_{\oplus}(t) = v_{sun} + v_{orb} \cos\gamma \cos[\omega(t-t_0)]$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

the DM annual modulation signature has a different origin and peculiarities (e.g. the phase) than those effects correlated with the seasons

To mimic this signature, spurious effects and side reactions must be able to account for the whole observed modulation amplitude, and also to satisfy simultaneously all the requirements

Highly radiopure NaI(Tl) experiment in DAMA

DAMA/NaI

Concluded on July 2002; 7 annual cycles collected; exposure 0.29 ton×yr



Model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

+ many results on other rare processes



DAMA/LIBRA

New NaI(Tl) detectors with better radiopurity features

Residual contaminations: ^{232}Th, ^{238}U and ^{40}K at level of 10^{-12} g/g

- DAMA/LIBRA-phase1: 7 annual cycles, 1.04 ton × yr
- Model independent evidence of a particle DM component in the galactic halo at 9.3σ C.L.
- DAMA/LIBRA-phase2: lowering software energy threshold below 2 keV; 8 annual cycles released so far (1.53 ton × yr)

The pioneer DAMA/NaI: ≈100 kg highly radiopure NaI(Tl)

Performances:

N.Cim.A112(1999)545-575, EPJC18(2000)283, Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

Results on rare processes:

- Possible Pauli exclusion principle violation — PLB408(1997)439
- CNC processes — PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) — PLB460(1999)235
- Search for solar axions — PLB515(2001)6
- Exotic Matter search — EPJdirect C14(2002)1
- Search for superdense nuclear matter — EPJA23(2005)7
- Search for heavy clusters decays — EPJA24(2005)51

Results on DM particles:

- PSD — PLB389(1996)757
- Investigation on diurnal effect — N.Cim.A112(1999)1541
- Exotic Dark Matter search — PRL83(1999)4918
- **Annual Modulation Signature** — PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23, EPJC18(2000)283, PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205, PRD77(2008)023506, MPLA23(2008)2125



Model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

total exposure (7 annual cycles) 0.29 ton × yr

The DAMA/LIBRA set-up ~250 kg NaI(Tl) (Large sodium **I**odide **B**ulk for **RA**re processes)

As a result of a 2nd generation R&D for more radiopure NaI(Tl) by exploiting new chemical/physical radiopurification techniques (all operations involving - including photos - in HP Nitrogen atmosphere)

Residual contaminations in the new DAMA/LIBRA NaI(Tl) detectors: ^{232}Th, ^{238}U and ^{40}K at level of 10^{-12} g/g



- Radiopurity, performances, procedures, etc.: NIMA592(2008)297, JINST 7 (2012) 03009
- Results on DM particles,
  - Annual Modulation Signature: EPJC56(2008)333, EPJC67(2010)39, EPJC73(2013)2648.
  - Related results: PRD84(2011)055014, EPJC72(2012)2064, IJMPA28(2013)1330022, EPJC74(2014)2827, EPJC74(2014)3196, EPJC75(2015)239, EPJC75(2015)400, IJMPA31(2016) dedicated issue, EPJC77(2017)83
- Results on rare processes:
  - PEPv: EPJC62(2009)327, arXiv1712.08082;
  - CNC: EPJC72(2012)1920;
  - IPP in ^{241}Am: EPJA49(2013)64

DAMA/LIBRA–phase1 (7 annual cycles, 1.04 ton×yr) confirmed the model-independent evidence of DM: reaching 9.3σ C.L.

DAMA/LIBRA–phase2

JINST 7(2012)03009

Upgrade on Nov/Dec 2010: all PMTs replaced with new ones of higher Q.E.



Q.E. of the new PMTs:
33 – 39% @ 420 nm
36 – 44% @ peak

DAMA/LIBRA–phase2

JINST 7(2012)03009

Lowering software energy threshold below 2 keV:

- to study the nature of the particles and features of astrophysical, nuclear and particle physics aspects, and to investigate 2nd order effects
- special data taking for *other rare processes*



The resolution:



The contaminations:

	^{226}Ra (Bq/kg)	^{235}U (mBq/kg)	^{228}Ra (Bq/kg)	^{228}Th (mBq/kg)	^{40}K (Bq/kg)
Mean Contamination	0.43	47	0.12	83	0.54
Standard Deviation	0.06	10	0.02	17	0.16

The light responses:

DAMA/LIBRA-phase1: 5.5 – 7.5 ph.e./keV
DAMA/LIBRA-phase2: 6-10 ph.e./keV

DAMA/LIBRA-phase2 data taking

Upgrade at end of 2010: all PMTs replaced with new ones of higher Q.E.



Energy resolution @ 60 keV mean value:

prev. PMTs 7.5% (0.6% RMS)
new HQE PMTs 6.7% (0.5% RMS)

Annual Cycles	Period	Mass (kg)	Exposure (kg x d)	$(\alpha-\beta^2)$
I	Dec 23, 2010 – Sept. 9, 2011	commissioning		
II	Nov. 2, 2011 – Sept. 11, 2012	242.5	62917	0.519
III	Oct. 8, 2012 – Sept. 2, 2013	242.5	60586	0.534
IV	Sept. 8, 2013 – Sept. 1, 2014	242.5	73792	0.479
V	Sept. 1, 2014 – Sept. 9, 2015	242.5	71180	0.486
VI	Sept. 10, 2015 – Aug. 24, 2016	242.5	67527	0.522
VII	Sept. 7, 2016 – Sept. 25, 2017	242.5	75135	0.480
VIII	**Sept. 25, 2017 – Aug. 20, 2018**	**242.5**	**68759**	**0.557**
IX	**Aug. 24, 2018 – Oct. 3, 2019**	**242.5**	**77213**	**0.446**

New data release July 2021

- ✓ Fall 2012: new preamplifiers installed + special trigger modules.
- ✓ Calibrations 8 a.c.: $\approx 1.6 \times 10^8$ events from sources
- ✓ Acceptance window eff. 8 a.c.: $\approx 4.2 \times 10^6$ events ($\approx 1.7 \times 10^5$ events/keV)

Exposure with this data release of DAMA/LIBRA-phase2: **1.53 ton × yr**

Exposure DAMA/NaI+DAMA/LIBRA-phase1+phase2: **2.86 ton × yr**

Model Independent Annual Modulation Result

DAMA/LIBRA-phase2 (8 a.c. , 1.53 ton × yr)

experimental residuals of the single-hit scintillation events rate vs time and energy

1-2 keV



1-3 keV



1-6 keV



2-6 keV



Absence of modulation? No

χ^2/dof = 130/69 (1-2 keV); 176/69 (1-3 keV); 202/69 (1-6 keV); 157/69 (2-6 keV)

Fit on DAMA/LIBRA-phase2

$A\cos[\omega(t-t_0)]$; t_0 = 152.5 d, T = 1.00 y

1-2 keV

A=(0.0224±0.0030) cpd/kg/keV

χ^2/dof = 75.8/68 **7.4 σ C.L.**

1-3 keV

A=(0.0191±0.0020) cpd/kg/keV

χ^2/dof = 81.6/68 **9.7 σ C.L.**

1-6 keV

A=(0.01048±0.00090) cpd/kg/keV

χ^2/dof = 66.2/68 **11.6 σ C.L.**

2-6 keV

A=(0.00933±0.00094) cpd/kg/keV

χ^2/dof = 58.2/68 **9.9 σ C.L.**

The data of DAMA/LIBRA-phase2 favor the presence of a modulated behavior with proper features at 11.6σ C.L.

Rate behaviour above 6 keV

- **No Modulation above 6 keV**



Mod. Ampl. (6-14 keV): cpd/kg/keV

(0.0032 ± 0.0017) DAMA/LIBRA-ph2_2
(0.0016 ± 0.0017) DAMA/LIBRA-ph2_3
(0.0024 ± 0.0015) DAMA/LIBRA-ph2_4
-(0.0004 ± 0.0015) DAMA/LIBRA-ph2_5
(0.0001 ± 0.0015) DAMA/LIBRA-ph2_6
(0.0015 ± 0.0014) DAMA/LIBRA-ph2_7
-(0.0005 ± 0.0013) DAMA/LIBRA-ph2_8
-(0.0003 ± 0.0014) DAMA/LIBRA-ph2_9

→ statistically consistent with zero

DAMA/LIBRA-phase2_2_9



σ ≈ 1%, fully accounted by statistical considerations

- **No modulation in the whole energy spectrum:** studying integral rate at higher energy, R_{90}

- R_{90} percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA running periods

- Fitting the behaviour with time, adding a term modulated with period and phase as expected for DM particles:

Period	Mod. Ampl.
DAMA/LIBRA-ph2_2	(0.12±0.14) cpd/kg
DAMA/LIBRA-ph2_3	-(0.08±0.14) cpd/kg
DAMA/LIBRA-ph2_4	(0.07±0.15) cpd/kg
DAMA/LIBRA-ph2_5	-(0.05±0.14) cpd/kg
DAMA/LIBRA-ph2_6	(0.03±0.13) cpd/kg
DAMA/LIBRA-ph2_7	-(0.09±0.14) cpd/kg
DAMA/LIBRA-ph2_8	-(0.18±0.13) cpd/kg
DAMA/LIBRA-ph2_9	(0.08±0.14) cpd/kg

consistent with zero

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region → R_{90} ~ tens cpd/kg → ~ 100 σ far away

No modulation above 6 keV

This accounts for all sources of bckg and is consistent with the studies on the various components

DM model-independent Annual Modulation Result

DAMA/LIBRA-phase2 (8 a.c., 1.53 ton × yr)

Multiple hits events = Dark Matter particle "switched off"





Single hit residual rate (red)

vs

Multiple hit residual rate (green)

- Clear modulation in the single hit events;
- No modulation in the residual rate of the multiple hit events

This result offers an additional strong support for the presence of DM particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background

The analysis in frequency

(according to PRD75 (2007) 013010)

To perform the Fourier analysis of the data in a wide region of frequency, the single-hit scintillation events have been grouped in 1 day bins

The whole power spectra up to the Nyquist frequency





DAMA/NaI + DAMA/LIBRA-(ph1+ph2) (22 yr)
total exposure: 2.86 ton×yr

Zoom around the 1 y^{-1} peak



Green area: 90% C.L. region calculated taking into account the signal in (2-6) keV

Clear annual modulation in (2-6) keV + only aliasing peaks far from signal region

Energy distribution of the modulation amplitudes

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

$T=2\pi/\omega=1$ yr t_0= 152.5 day

DAMA/NaI + DAMA/LIBRA-phase1
+ DAMA/LIBRA-phase2 (2.86 ton×yr)

max-likelihood analysis of the single hit scintillation events



A clear modulation is present in the (1-6) keV energy interval, while S_m values compatible with zero are present just above

- The S_m values in the (6–14) keV energy interval have random fluctuations around zero with χ^2 equal to 20.3 for 16 degrees of freedom (upper tail probability 21%).
- In (6–20) keV χ^2/dof = 42.2/28 (upper tail probability 4%). The obtained χ^2 value is rather large due mainly to two data points, whose centroids are at 16.75 and 18.25 keV, far away from the (1–6) keV energy interval. The P-values obtained by excluding only the first and either the points are 14% and 23%.

S_m for each annual cycle



DAMA/LIBRA-phase1 +
DAMA/LIBRA-phase2
total exposure: **2.57 ton×yr**

Energy bin (keV)	run test probability	
	Lower	Upper
1-2	89%	37%
2-3	87%	30%
3-4	17%	94%
4-5	17%	94%
5-6	30%	85%

The signal is well distributed over all the annual cycles in each energy bin

A sinusoidal contribution in the signal? Phase ≠ 152.5 day?

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)] = S_0 + Y_m \cos[\omega(t - t^*)]$$

For Dark Matter signals:

- $|Z_m| \ll |S_m| \approx |Y_m|$
- $t^* \approx t_0 = 152.5d$
- $\omega = 2\pi/T$
- $T = 1$ year

DAMA/NaI + DAMA/LIBRA-phase1 + DAMA/LIBRA-phase2 (8 a.c.) [2.86 ton × yr]



Slight differences from 2nd June are expected in case of contributions from non thermalized DM components (as e.g. the SagDEG stream)



E (keV)	S_m (cpd/kg/keV)	Z_m (cpd/kg/keV)	Y_m (cpd/kg/keV)	t* (day)
DAMA/NaI + DAMA/LIBRA-ph1 + DAMA/LIBRA-ph2				
2-6	0.0097 ± 0.0007	- 0.0003 ± 0.0007	0.0097 ± 0.0007	150.5 ± 4.0
6-14	0.0003 ± 0.0005	-0.0006 ± 0.0005	0.0007 ± 0.0010	undefined
1-6	0.0104 ± 0.0007	0.0002 ± 0.0007	0.0104 ± 0.0007	153.5 ± 4.0

Efforts towards lower software energy threshold

- decreasing the software energy threshold down to 0.75 keV
- using the same technique to remove the noise pulses
- evaluating the efficiency by dedicated studies

New data point with the 8 a.c. of **DAMA/LIBRA-phase2 (1.53 ton×yr)**



- A clear modulation is also present below 1 keV, from 0.75 keV, while S_m values compatible with zero are present just above 6 keV
- This preliminary result suggests the necessity to lower the software energy threshold and to improve the experimental error on the first energy bin

Stability parameters of DAMA/LIBRA–phase2

Modulation amplitudes obtained by fitting the time behaviours of main running parameters, acquired with the production data, when including a DM-like modulation

Running conditions stable at a level better than 1% also in the new running periods

	DAMA/LIBRA-phase2_2	DAMA/LIBRA-phase2_3	DAMA/LIBRA-phase2_4	DAMA/LIBRA-phase2_5	DAMA/LIBRA-phase2_6	DAMA/LIBRA-phase2_7	DAMA/LIBRA-phase2_8	DAMA/LIBRA-phase2_9
Temperature (°C)	(0.0012 ± 0.0051)	$-(0.0002 \pm 0.0049)$	$-(0.0003 \pm 0.0031)$	(0.0009 ± 0.0050)	(0.0018 ± 0.0036)	$-(0.0006 \pm 0.0035)$	$-(0.0029 \pm 0.0039)$	(0.0014 ± 0.0033)
Flux N_2 (l/h)	$-(0.15 \pm 0.18)$	$-(0.02 \pm 0.22)$	$-(0.02 \pm 0.12)$	$-(0.02 \pm 0.14)$	$-(0.01 \pm 0.10)$	$-(0.01 \pm 0.16)$	(0.05 ± 0.25)	(0.014 ± 0.092)
Pressure (mbar)	$(1.1 \pm 0.9) \times 10^{-3}$	$(0.2 \pm 1.1)) \times 10^{-3}$	$(2.4 \pm 5.4) \times 10^{-3}$	$(0.6 \pm 6.2) \times 10^{-3}$	$(1.5 \pm 6.3) \times 10^{-3}$	$(7.2 \pm 8.6) \times 10^{-3}$	$(3 \pm 12) \times 10^{-3}$	$(3.5 \pm 4.9) \times 10^{-3}$
Radon (Bq/m^3)	(0.015 ± 0.034)	$-(0.002 \pm 0.050)$	$-(0.009 \pm 0.028)$	$-(0.044 \pm 0.050)$	(0.082 ± 0.086)	(0.06 ± 0.11)	$-(0.046 \pm 0.076)$	(0.002 ± 0.035)
Hardware rate above single ph.e. (Hz)	$-(0.12 \pm 0.16) \times 10^{-2}$	$(0.00 \pm 0.12) \times 10^{-2}$	$-(0.14 \pm 0.22) \times 10^{-2}$	$-(0.05 \pm 0.22) \times 10^{-2}$	$-(0.06 \pm 0.16) \times 10^{-2}$	$-(0.08 \pm 0.17) \times 10^{-2}$	$(0.04 \pm 0.20) \times 10^{-2}$	$-(0.19 \pm 0.18) \times 10^{-2}$

All the measured amplitudes well compatible with zero

+ none can account for the observed effect

(to mimic such signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also simultaneously satisfy all the 6 requirements)

Summary of the results obtained in the additional investigations of possible systematics or side reactions – DAMA/LIBRA

NIMA592(2008)297, EPJC56(2008)333, J. Phys. Conf. ser. 203(2010)012040, arXiv:0912.0660, S.I.F.Atti Conf.103(211), Can. J. Phys. 89 (2011) 11, Phys.Proc.37(2012)1095, EPJC72(2012)2064, arxiv:1210.6199 & 1211.6346, IJMPA28(2013)1330022, EPJC74(2014)3196, IJMPA31(2017)issue31, Universe4(2018)116, Bled19(2018)27, NPAE19(2018)307, PPNP114(2020)103810

Source	Main comment	Cautious upper limit (90%C.L.)
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5\times10^{-6}$ cpd/kg/keV
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
NOISE	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
ENERGY SCALE	Routine + intrinsic calibrations	$<1\text{-}2\times10^{-4}$ cpd/kg/keV
EFFICIENCIES	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV *multiple-hits* events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
SIDE REACTIONS	Muon flux variation measured at LNGS	$<3\times10^{-5}$ cpd/kg/keV

+ they cannot satisfy all the requirements of annual modulation signature

Thus, they cannot mimic the observed annual modulation effect

About ANAIS result

- ANAIS has ≈ 5 times larger counting rate in [1,2] keV than DAMA/LIBRA-phase2
- High counting rate in ROI explained as populations, other than bckg, "*which could be leaking at the lowest energies in the ROI*" being the trigger rate "*dominated by other events, some of them with origin in the PMTs, others still unexplained*"
- Even **a 0.3% instability** of the ANAIS counting rate in the [1-6] keV region is enough to hide the annual modulation signal detected by DAMA: A ≈ 0.01 cpd/kg/keV (green line in the plot)
- In ANAIS the detection efficiencies of the applied cuts are **not** periodically evaluated with dedicated calibrations at very low energy as in DAMA/LIBRA
- The only check on stability of the cut-efficiencies is a fit on the counting rate of low energy events induced by the ^{22}Na or ^{40}K contaminations, selected in double coincidences, and with cuts applied
- But statistics is low: ≈ 100 events in bin of 10 days, i.e. a 10% error/bin
- A fit of these data including a modulated components shows that they cannot exclude an effect at the level of 2-3%, much higher than the needed stability
- while the searched effect requires a stability of the efficiency at the level of 0.4% in [1,2] keV





- Similar result can be obtained for the [2-5] keV region (studying ^{40}K double) the sensitivity is ≈ 1% (needed: <0.4%)

Moreover:

- Different quenching factors are expected and measured for different NaI(Tl) crystals (they depends, e.g., on the used growing technique, on the different thallium doping concentration, …)
- A clear evidence is offered by the different α/β light ratio measured with DAMA and COSINE crystals
- As mentioned also in the ANAIS paper, this effect introduce a systematic uncertainty in the comparison with DAMA/LIBRA

About Interpretation: is an "universal" and "correct" way to approach the problem of DM and comparisons?



No, it isn't. This is just a largely arbitrary/partial/incorrect exercise



see e.g.: Riv.N.Cim. 26 n.1(2003)1, IJMPD13(2004) 2127, EPJC47(2006)263, IJMPA21(2006)1445, EPJC56(2008)333, PRD84 (2011)055014, IJMPA28 (2013)1330022, NPAE20(4) (2019)317, PPNP114(2020) 103810

...models...

- Which particle?
- Which interaction coupling?
- Which Form Factors for each target-material?
- Which Spin Factor?
- Which nuclear model framework?
- Which scaling law?
- Which halo model, profile and related parameters?
- Streams?
- ...

...and experimental aspects...

- Exposures
- Energy threshold
- Calibrations
- Stability of all the operating conditions.
- Efficiencies
- Definition of fiducial volume and non-uniformity
- Detector response (phe/keV)
- Energy scale and energy resolution
- Selections of detectors and of data.
- Subtraction/rejection procedures and stability in time of all the selected windows and related quantities
- Quenching factors, channeling, ...
- ...

Uncertainty in experimental parameters, and necessary **assumptions** on various related astrophysical, nuclear and particle-physics aspects, affect all the results at various extent, both in terms of exclusion plots and in terms of allowed regions/volumes. Thus comparisons with **a fixed set** of assumptions and parameters' values **are intrinsically strongly uncertain**.

No direct model-independent comparison among expts with different target-detectors and different approaches

Model-independent evidence by DAMA/NaI and DAMA/LIBRA-ph1, -ph2

well compatible with several candidates in many astrophysical, nuclear and particle physics scenarios

- Just few examples of interpretation of the annual modulation in terms of candidate particles in some scenarios
- E_{th}=1 keV; old data release



Examples of model-dependent analyses

NPAE 20(4) (2019) 317
PPNP114(2020)103810

A large (but not exhaustive) class of halo models and uncertainties are considered

E_{th}=1 keV; old data release



DM particles elastically scattering off target nuclei – SI interaction

$$\sigma_{SI}(A,Z) \propto m_{red}^2(A,DM)\left[f_p Z + f_n(A-Z)\right]^2$$

Case of isospin violating SI coupling: $f_p \neq f_n$



1. Constants q.f.
2. Varying q.f.(E_R)
3. With channeling effect



Even a relatively small SD (SI) contribution can drastically change the allowed region in the (m_{DM}, $\xi\sigma_{SI(SD)}$) plane

- Two bands at low mass and at higher mass;
- Good fit for low mass DM candidates at $f_n/f_p \approx -53/74 = -0.72$ (signal mostly due to ^{23}Na recoils).
- The inclusion of the uncertainties related to halo models, quenching factors, channeling effect, nuclear form factors, etc., can also support for f_n/f_p=1 low mass DM candidates either including or not the channeling effect.
- The case of isospin-conserving f_n/f_p=1 is well supported at different extent both at lower and larger mass.

Running phase2 with lower software energy threshold below 1 keV with high efficiency

Enhancing experimental sensitivities and improving DM corollary aspects, other DM features, second order effects and other rare processes

- After a dedicated R&D on new high Q.E. PMTs with increased radio-purity
- After the study of possible new protocols for possible modifications of the detectors



Alternative strategy has been chosen, upgrading the hardware:

- new miniaturized low background **pre-amps** directly installed on the low-background supports of the **voltage dividers** of the low background high Q.E. **PMTs** of phase2
- higher vertical resolution 14bit digitizers



Design of the voltage divider and preamplifier mounted on the Pyralux support

The features of the voltage divider+preamp system:

- S/N improvement ≈3.0-9.0;
- discrimination of the single ph.el. from electronic noise: 3 - 8;
- the Peak/Valley ratio: 4.7 - 11.6;
- residual radioactivity lower than that of single PMT

Conclusions

- Model-independent evidence for a signal that satisfies all the requirements of the DM annual modulation signature at 13.7σ C.L. (22 independent annual cycles with 3 different set-ups: 2.86 ton × yr)
- Modulation parameters determined with increasing precision
- New investigations on different peculiarities of the DM signal in progress
- Full sensitivity to many kinds of DM candidates and interactions types (both inducing recoils and/or e.m. radiation), full sensitivity to low and high mass candidates





- Model-dependent analyses improve the C.L. and restrict the allowed parameters' space for the various scenarios
- DAMA/LIBRA–phase2 continuing data taking
- Preliminary efforts towards 0.75 keV software energy threshold done
- DAMA/LIBRA–phase2 towards lower software energy threshold of 0.5 keV. New divider/amp systems and new 14bit digitizers
- Continuing investigations of rare processes other than DM
- Other pursued ideas: $ZnWO_4$ anisotropic scintillator for DM directionality. Response to nuclear recoils measured.