



Contribution ID: 59

Type: Talk

Triple nuclear collisions and new perspectives to explore the QCD matter properties under new extreme conditions

Wednesday, 1 September 2021 11:00 (30 minutes)

Kyryl Bugaev^{a,b}, Oleksandr Vitiuk^c, Valery Pugatch^c, Vasyl Dobishuk^c, Sergiy Chernyshenko^c, Boris Grinyuk^b, Pavlo Panasiuk^a, Nazar Yakovenko^a, Elizaveta Zherebtsova^d, Larissa Bravina^e, Arkadiy Taranenko^d, Evgeny Zabrodin^{e,f} and Marcus Bleicher^g

^aDepartment of Physics, Taras Shevchenko National University of Kyiv, 03022 Kyiv, Ukraine

^bBogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Metrologichna str. 14-B, 03680 Kyiv, Ukraine

^cInstitute for Nuclear Research, National Academy of Sciences of Ukraine, Prospekt Nauki av. 47, 03680 Kyiv, Ukraine

^dNational Research Nuclear University "MEPhI" (Moscow Engineering Physics Institute), Kashirskoe Shosse 31, 115409 Moscow, Russia

^eUniversity of Oslo, POB 1048 Blindern, N-0316 Oslo, Norway

^fSkobeltsyn Institute of Nuclear Physics, Moscow State University, 119899 Moscow, Russia

^gInstitute for Theoretical Physics, Goethe University, Max-von-Laue-Str. 1, 60438 Frankfurt am Main, Germany

We suggest to explore an entirely new method to experimentally and theoretically study the phase diagram of str

We simulated the TNC using the UrQMD 3.4 model [3, 4] at the beam center-of-mass collision energies $\sqrt{s} = 200$ GeV and $\sqrt{s} = 2.76$ TeV. It is found that in the most central and simultaneous TNC the initial baryonic charge density is about 3 times higher than the one achieved in the usual binary nuclear collisions at the same energies [5]. As a consequence, a production of protons and Λ -hyperons is increased by 2 and 1.5, respectively, while a sizable suppression of their antiparticles is observed.

At the beam center-of-mass collision energies of 10-40 GeV, the production of protons as well as of Λ -hyperons is enhanced approximately by a factor of 2.2 compared to the binary collisions, while the positive kaons are enhanced by 1.5. Hence we conclude that in the TNC method it is possible to create substantially denser strange matter than in the binary collisions. It is argued that this method at lower energies can be of principal importance for searching the (tri)critical endpoint of the QCD phase diagram [5].

References:

- [1] V. Pugatch, International Conference "CERN-Ukraine co-operation: current state and prospects" Kharkiv, 15-May-2018; LHCb-TALK-2018-557.
- [2] LHCb Collaboration. SMOG2 Technical Design Report. CERN-LHCC-2019-0051.
- [3] S.A. Bass et al., Prog. Part. Nucl. Phys. 41 (1998), 225-370.
- [4] M. Bleicher et al., J. Phys. G 25 (1999), 1859-1896.
- [5] K. Bugaev et al., talk at the Online «Strangeness in Quark Matter» Conference 2021, Brookhaven, May 17-22, 2021; <https://indico.cern.ch/event/985652/sessions/392917/#20210521>

Is this abstract from experiment?

No

Name of experiment and experimental site

N/A

Is the speaker for that presentation defined?

Yes

Details

Kyrill Bugaev, Prof., Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine,
Metrologichna str. 14-B, 03680 Kyiv, Ukraine,
<http://bitp.kiev.ua/en/about>

Internet talk

Yes

Primary authors: BUGAEV, Kyrill (Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine); VITIUK, Oleksandr (Taras Shevchenko National University of Kyiv)

Co-authors: Prof. BRAVINA, Larissa (University of Oslo, POB 1048 Blindern, N-0316 Oslo, Norway); ZABRODIN, Evgeny (University of Oslo (NO)); BLEICHER, Marcus (Uni Frankfurt)

Presenters: BUGAEV, Kyrill (Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine); VITIUK, Oleksandr (Taras Shevchenko National University of Kyiv)

Session Classification: B Heavy Ion Collisions and Critical Phenomena