

Contribution ID: 9

Type: Poster

## Simulations for a High-Current Electron Gun for the EBIT Charge Breeder of the ReA Post-Accelerator at FRIB

The Facility for Rare Isotope Beams (FRIB) at Michigan State University is nearing completion. In some cases, the RIB rates at FRIB are expected to exceed  $10^{10}$  particles/s. The ReA EBIT charge breeder operates with an electron-beam current of 300 –600 mA for a density of 170 –340 A/cm<sup>2</sup>. This current corresponds to a maximum capacity of  $10^{10}$  elementary charges, which can be insufficient to handle high FRIB rates. To increase the EBIT electron-beam current and density, electron-beam simulations have been performed for a new electron-gun insert with a dispenser cathode having a larger emitting area. An electron beam of 2 A was the target current to reach a capacity of  $5 \times 10^{10}$  elementary charges. Parameter studies have been conducted to optimize beam transmission. A simulated current density at the trap was obtained to be 432 A/cm<sup>2</sup>. The basic design of the insert and result of the simulations for the ReA EBIT will be presented.

## E-mail for contact person

berekha@gmail.com

## **Funding Information**

Work supported by National Science Foundation, under contract No. PHY-1565546

**Primary authors:** Dr SON, Hyock-Jun (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr LAPIERRE, Alain (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr VILLARI, Antonio (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University); Dr SCHWARZ, Stefan (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michiga

**Presenter:** Dr SON, Hyock-Jun (Facility for Rare Isotope Beams, National Superconducting Cyclotron Laboratory, Michigan State University)

Session Classification: Poster Session 2

Track Classification: Production of highly charged ion beams