TIPP 2011 - 2nd International Conference on Technology and Instrumentation in Particle Physics



Contribution ID: 391

Type: Oral Presentation

Search for neutrinoless double beta decay with the NEMO-3 detector and R&D for SuperNEMO

Monday 13 June 2011 15:20 (20 minutes)

The NEMO-3 (Neutrino Ettore Majorana Observatory) experiment, located in the Modane Underground Laboratory, searches for neutrinoless double beta decay. The experiment has been taking data since 2003 with seven double beta isotopes and completed data acquisition in late 2010. Two neutrino double beta decay results for the main isotopes (7 kg of 100Mo and 1 kg of 82Se), new results for 150Nd and 130Te, as well as results for 96Zr, 48Ca, and 116Cd are presented. NEMO-3 uses a unique technique that allows for the in situ measurement of background contamination. No evidence for neutrinoless double beta decay has been found to date. The data are also interpreted in terms of alternative models such as weak right-handed currents and Majoron emission. In this talk, I will discuss the measurements made with NEMO-3 and discuss the status, research and design of the next generation experiment, SuperNEMO.

Authors: PAHLKA, Benton (The University of Texas at Austin); PAHLKA, Benton (Fermilab)Presenter: PAHLKA, Benton (The University of Texas at Austin)Session Classification: Detector for Neutrinos

Track Classification: Detectors for neutrino physics