

# The Higgs $p_T$ Spectrum and Total Cross Section with Fiducial Cuts at $N^3LL' + N^3LO$

Johannes Michel

MIT Center for Theoretical Physics

Higgs 2021

Stony Brook, October 21



# The Higgs $p_T$ Spectrum and Total Cross Section with Fiducial Cuts at $N^3LL' + N^3LO$

based on

[PRL 127 (2021) 7, 072001, 2102.08039]

in collaboration with

G. Billis, B. Dehnadi, M. Ebert, F. Tackmann



- Measure fiducial & differential Higgs cross sections at the LHC
  - ▶ Most model-independent way we have to search for BSM in the Higgs sector
- Total fiducial cross section measures deviations from SM gluon-fusion rate

$$\begin{array}{c}
 g \\
 \text{wavy line} \\
 \nearrow \\
 \text{triangle} \\
 \text{t} \\
 \searrow \\
 g \\
 \text{wavy line} \\
 \text{---} H
 \end{array}
 +
 \begin{array}{c}
 g \\
 \text{wavy line} \\
 \nearrow \\
 \text{red hatched circle} \\
 \searrow \\
 g \\
 \text{wavy line} \\
 \text{---} H
 \end{array}
 = \left( \frac{\alpha_s}{12\pi v} C_t + \frac{v}{\Lambda^2} C_{HG} \right) H G_{\mu\nu}^a G^{a,\mu\nu}$$

- $p_T^H$  spectrum is the most important differential quantity
- High  $p_T^H \sim \sqrt{\hat{s}} \gg m_H$  increases sensitivity to new operators  
 [See talk by R. Abraham in Wednesday session for an interesting case study]
- Focus of this talk:  $p_T^H \lesssim m_H \sim \sqrt{\hat{s}} \ll 2m_t$  (or  $p_T^H$  integrated over)
  - ▶ Measure or put bounds on anomalous  $b$ ,  $c$ , and light quark Yukawa couplings  
 [Bishara, Haisch, Monni, Re '16; Soreq, Zhu, Zupan '16; see e.g. ATLAS-CONF-2019-029]

- Uncertainty  $\Delta\sigma$  on SM prediction translates into discovery reach:

$$\frac{\Delta\sigma}{\sigma} \sim \frac{v^2}{\Lambda_{\text{BSM}}^2} \Leftrightarrow \Lambda_{\text{BSM}} \sim v \sqrt{\frac{\sigma}{\Delta\sigma}}$$

## Challenges for theory

- QCD corrections to  $gg \rightarrow H$  are large:  $\sigma/\sigma_{\text{LO}} \approx 3$ 
  - ▶ Calculation of inclusive cross section has been pushed to N<sup>3</sup>LO  
[Anastasiou, Duhr, Dulat, Furlan, Gehrmann, Herzog, Lazopoulos, Mistlberger '15-'18]
- But LHC experiments apply kinematic selection cuts on Higgs decay products
  - ▶ Need complete interplay of QCD corrections and  $\mathcal{O}(1)$  fiducial acceptance

# Fiducial predictions for $gg \rightarrow H \rightarrow \gamma\gamma$ at three loops



Consider  $gg \rightarrow H \rightarrow \gamma\gamma$  with ATLAS fiducial cuts:

$$p_T^{\gamma 1} \geq 0.35 m_H, \quad p_T^{\gamma 2} \geq 0.25 m_H, \quad |\eta^\gamma| \leq 2.37, \quad |\eta^\gamma| \notin [1.37, 1.52]$$

## Focus of this talk

[Billis, Dehnadi, Ebert, JM, Tackmann, PRL 127 (2021) 7, 072001, 2102.08039]

- Compute fiducial spectrum for  $q_T \equiv p_T^H = p_T^{\gamma\gamma}$  at  $N^3LL'+N^3LO$
- Compute total fiducial cross section at  $N^3LO$ , and improved by resummation



# Fiducial predictions for $gg \rightarrow H \rightarrow \gamma\gamma$ at three loops

Consider  $gg \rightarrow H \rightarrow \gamma\gamma$  with ATLAS fiducial cuts:

$$p_T^{\gamma 1} \geq 0.35 m_H, \quad p_T^{\gamma 2} \geq 0.25 m_H, \quad |\eta^\gamma| \leq 2.37, \quad |\eta^\gamma| \notin [1.37, 1.52]$$

## Focus of this talk

[Billis, Dehnadi, Ebert, JM, Tackmann, PRL 127 (2021) 7, 072001, 2102.08039]

- Compute fiducial spectrum for  $q_T \equiv p_T^H = p_T^{\gamma\gamma}$  at  $N^3\text{LL}' + N^3\text{LO}$
- Compute total fiducial cross section at  $N^3\text{LO}$ , and improved by resummation

- Previous state of the art was  $N^3\text{LL}(+\text{NNLO}_1)$  and NNLO, respectively

[Chen et al. '18; Bizoń et al. '18; Gutierrez-Reyes et al. '19; Becher, Neumann '20]

Kicked off a recent push for fiducial color singlet at complete three-loop accuracy:

- Complementary  $N^3\text{LO}$  results for fiducial  $Y_{\gamma\gamma}, \eta_{\gamma 1}, \Delta\eta_{\gamma\gamma}$  (with different method)  
[Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607; see talk by B. Mistlberger in this session]
- Fiducial  $N^3\text{LL}'$  results for Higgs  $q_T$  spectrum

[Re, Rottoli, Torrielli, 2104.07509; see talk by E. Re in this session]

[For Drell-Yan,  $\gamma\gamma$ , see also 2103.04974, 2106.11260, 2107.12478]

# Fiducial predictions for $gg \rightarrow H \rightarrow \gamma\gamma$ at three loops

Consider

$$p_T^{\gamma\gamma}$$



Focus

- Con

- Con

- Prev

[Chen



Kicked off a recent push for fiducial color singlet at complete three-loop accuracy:

- Complementary  $N^3$ LO results for fiducial  $Y_{\gamma\gamma}$ ,  $\eta_{\gamma 1}$ ,  $\Delta\eta_{\gamma\gamma}$  (with different method) [Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607; see talk by B. Mistlberger in this session]
- Fiducial  $N^3$ LL' results for Higgs  $q_T$  spectrum [Re, Rottoli, Torrielli, 2104.07509; see talk by E. Re in this session] [For Drell-Yan,  $\gamma\gamma$ , see also 2103.04974, 2106.11260, 2107.12478]


# Fiducial predictions for $gg \rightarrow H \rightarrow \gamma\gamma$ at three loops

Consider  $gg$

$$p_T^{\gamma\gamma} \geq \epsilon$$

Focus of the

- Compute
- Compute
- Previous [Chen et al.



[37, 1.52]

[2001, 2102.08039]

Information

Kicked off a recent push for fiducial color singlet at complete three-loop accuracy:

- Complementary  $N^3\text{LO}$  results for fiducial  $Y_{\gamma\gamma}, \eta_{\gamma\gamma}, \Delta\eta_{\gamma\gamma}$  (with different method) [Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607; see talk by B. Mistlberger in this session]
- Fiducial  $N^3\text{LL}'$  results for Higgs  $q_T$  spectrum [Re, Rottoli, Torrielli, 2104.07509; see talk by E. Re in this session] [For Drell-Yan,  $\gamma\gamma$ , see also 2103.04974, 2106.11260, 2107.12478]



## Structure of the $q_T$ spectrum

Compute cross section from  $\sigma = \int dq_T \frac{d\sigma}{dq_T}$  and *power expand* around IR,  $q_T \rightarrow 0$ :

$$\begin{aligned}\frac{d\sigma}{dq_T} &= \frac{d\sigma^{(0)}}{dq_T} + \frac{d\sigma^{(1)}}{dq_T} + \frac{d\sigma^{(2)}}{dq_T} + \dots \\ &\sim \frac{1}{q_T} \left[ \mathcal{O}(1) + \mathcal{O}\left(\frac{q_T}{m_H}\right) + \mathcal{O}\left(\frac{q_T^2}{m_H^2}\right) + \dots \right]\end{aligned}$$

$$\frac{d\sigma^{(0)}}{dq_T} = \sigma_{\text{LO}} \delta(q_T) + \sum_n \alpha_s^n \left\{ \sigma_n^V \delta(q_T) + \sum_m \sigma_{n,m}^{(0)} \left[ \frac{\ln^m(q_T/m_H)}{q_T} \right]_+ \right\}$$

- ▶ Contains LO contribution, virtual corrections, and log-enhanced singular terms

$$\frac{d\sigma^{(1)}}{dq_T} = \sum_n \alpha_s^n \sum_m \sigma_{n,m}^{(1)} \frac{1}{m_H} \ln^m(q_T/m_H)$$

- ▶ Still logarithmically divergent, only present if decay products are resolved

## Structure of the $q_T$ spectrum

Compute cross section from  $\sigma = \int dq_T \frac{d\sigma}{dq_T}$  and *power expand* around IR,  $q_T \rightarrow 0$ :

$$\frac{d\sigma}{dq_T} = \frac{d\sigma^{(0)}}{dq_T} + \frac{d\sigma^{(1)}}{dq_T} + \frac{d\sigma^{(2)}}{dq_T} + \dots$$
$$\sim \frac{1}{q_T} \left[ \mathcal{O}(1) + \mathcal{O}\left(\frac{q_T}{m_H}\right) + \mathcal{O}\left(\frac{q_T^2}{m_H^2}\right) + \dots \right]$$

$$\frac{d\sigma^{(2)}}{dq_T} = \sum_n \alpha_s^n \sum_m \sigma_{n,m}^{(2)} \frac{q_T}{m_H^2} \ln^m(q_T/m_H) + \dots$$

- ▶ Finite as  $q_T \rightarrow 0$ , extract by fitting known functional form to  $H + 1j$  calculation
  - ▶ For the experts: Use a differential  $q_T$  subtraction accounting for fiducial power corrections
  - ▶ Avoids shortcomings of slicing [e.g. 1807.11501, 2103.04974] or Projection to Born [e.g. 2102.07607]
  - ▶ See backup for details


Set up some notation, use that production and [decay \(acceptance\)](#) factorize:

$$\frac{d\sigma}{dq_T} = \int dY A(q_T, Y; \Theta) W(q_T, Y), \quad A_{\text{incl}} = 1, \quad W(q_T, Y) = \frac{d\sigma_{\text{incl}}}{dq_T dY}$$

## Leading-power factorization & resummation to $N^3LL'$


At leading power in  $q_T \ll m_H$ , the hadronic dynamics factorize as:

$$W^{(0)}(q_T, Y) = H(m_H^2, \mu) \int d^2\vec{k}_a d^2\vec{k}_b d^2\vec{k}_s \delta(q_T - |\vec{k}_a + \vec{k}_b + \vec{k}_s|) \\ \times B_g^{\mu\nu}(x_a, \vec{k}_a, \mu, \nu) B_{g\mu\nu}(x_b, \vec{k}_b, \mu, \nu) S(\vec{k}_s, \mu, \nu)$$



## Leading-power factorization & resummation to $N^3LL'$

- Factorization predicts singular structure of  $\frac{d\sigma}{dq_T}$  as  $q_T \rightarrow 0$
- Enables all-order resummation  $\Rightarrow$  Sudakov peak
- Resummation at  $N^3LL'$  involves a host of three and four-loop QCD ingredients [see backup for a complete list of ingredients and references]



*But if this were the end of the story, it'd be pretty boring indeed!*

... are the power corrections coming from the  $q_T$ -dependent acceptance:

$$\frac{d\sigma^{\text{fpc}}}{dq_T} \equiv \int dY \left[ A(q_T, Y; \Theta) - A^{(0)}(Y; \Theta) \right] W^{(0)}(q_T, Y)$$

- Uniquely predict all linear power corrections  $d\sigma^{(1)}$  because

$$A(q_T, Y; \Theta) = A^{(0)}(Y; \Theta) \left[ 1 + \mathcal{O}\left(\frac{q_T}{m_H}\right) \right]$$

$$W(q_T, Y) = W^{(0)}(q_T, Y) \left[ 1 + \mathcal{O}\left(\frac{q_T^2}{m_H^2}\right) \right]$$

- Resummed to the same  $N^3\text{LL}'$  accuracy as leading-power terms by resumming  $W^{(0)}$  and keeping exact  $A(q_T, Y; \Theta)$

[Presence of linear terms pointed out in Ebert, Tackmann, 1911.08486]

[Factorization & resummation demonstrated in Ebert, JM, Stewart, Tackmann, 2006.11382]

[See also Alekhin, Kardos, Moch, Trócsányi, 2104.02400; Salam, Slade, 2106.08329]

## Key point

Fiducial power corrections induce resummation effects *in the total cross section*

Compare fixed-order series, isolating the effect of  $\int dq_T \frac{d\sigma^{\text{fpc}}}{dq_T}$ :

$$\sigma_{\text{incl}}^{\text{FO}} = 13.80 [1 + 1.291 + 0.783 + 0.299] \text{ pb}$$

$$\begin{aligned} \sigma_{\text{fid}}^{\text{FO}} &= 6.928 [1 + 1.429 + 0.723 + 0.481] \text{ pb} \\ &= 6.928 [1 + (1.300 + 0.129_{\text{fpc}}) + (0.784 - 0.061_{\text{fpc}}) + (0.331 + 0.150_{\text{fpc}})] \text{ pb} \end{aligned}$$



- Fiducial power corrections show no convergence, remainder is similar to inclusive

## Key point

Fiducial power corrections induce resummation effects *in the total cross section*

Two ways to understand this:

1. Acceptance acts as a weight under the  $q_T$  integral



$$\sigma_{\text{incl}} = \int dq_T \mathbf{W}(q_T) \quad \sigma_{\text{fid}} = \int dq_T \mathbf{A}(q_T) \mathbf{W}(q_T)$$



# Resummation effects in the total cross section

## Key point

Fiducial power corrections induce resummation effects *in the total cross section*

Two ways to understand this:



1. Acceptance acts as a weight under the  $q_T$  integral
2. We're cutting on the resummation-sensitive photon  $p_T$



► Either way, effect is *predicted* by resummed perturbation theory




# Results: The fiducial $q_T$ spectrum at $N^3LL'+N^3LO$



- Total uncertainty is  $\Delta_{tot} = \Delta_{q_T} \oplus \Delta_{\varphi} \oplus \Delta_{match} \oplus \Delta_{FO} \oplus \Delta_{nons}$   
[See also Ebert, JM, Stewart, Tackmann, 2006.11382 for details]
- Observe excellent perturbative convergence & uncertainty coverage
  - Crucial to consider *every* variation to probe all parts of the prediction
- Divide  $H \rightarrow \gamma\gamma$  branching ratio  $\mathcal{B}_{\gamma\gamma}$  out of data [LHC Higgs Cross Section WG, 1610.07922]
- Data are corrected for other production channels, photon isolation efficiency [ATLAS, 1802.04146]

# Results: The total fiducial cross section at $N^3$ LO and $N^3$ LL'+ $N^3$ LO




- Large  $N^3$ LO correction to fiducial cross section (worse than inclusive)
  - Caused by fiducial power corrections, *not* captured by rescaling inclusive  $N^3$ LO result
  - Recently, proposals for elaborate cuts to eliminate fiducial power corrections [Salam, Slade, 2106.08329]
- Resummation restores convergence, gives detailed handle on uncertainty
- ▶ Today's message: Theory can deal with it!

# Outlook: Resummation effects in other $H \rightarrow \gamma\gamma$ observables



- “Infrared sensitivity” observed also in other Higgs observables at  $N^3\text{LO}$

[Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607]

↔ Precisely the fiducial power corrections we can analytically deal with and resum



Note: Plots on the right show only  $\sigma^{\text{fpc}}$ .




# Outlook: Resummation effects in other $H \rightarrow \gamma\gamma$ observables


- “Infrared sensitivity” observed also in other Higgs observables at N<sup>3</sup>LO

[Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607]

⇒ Precisely the fiducial power corrections we can analytically deal with and resum



Note: Plots on the right show only  $\sigma^{\text{fpc}}$ .




# Outlook: Resummation effects in other $H \rightarrow \gamma\gamma$ observables


- “Infrared sensitivity” observed also in other Higgs observables at N<sup>3</sup>LO

[Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607]


↔ Precisely the fiducial power corrections we can analytically deal with and resum



Note: Plots on the right show only  $\sigma^{\text{fpc}}$ .




- Presented  $N^3LL'+N^3LO$  and  $N^3LO$  predictions for fiducial  $p_T^H$  spectrum and total fiducial cross section for  $gg \rightarrow H \rightarrow \gamma\gamma$  at the LHC
  - First direct comparison to LHC data at genuine three-loop order



- Resummed large fiducial power corrections induced by experimental acceptance
  - Even *total* fiducial cross sections are sensitive to  $q_T$  resummation effects
  - Enables best-possible combined predictions for other  $H \rightarrow \gamma\gamma$  observables

- Presented  $N^3LL'+N^3LO$  and  $N^3LO$  predictions for fiducial  $p_T^H$  spectrum and total fiducial cross section for  $gg \rightarrow H \rightarrow \gamma\gamma$  at the LHC
- First direct comparison to LHC data at genuine three-loop order





- Resummed large fiducial power corrections induced by experimental acceptance
- Even *total* fiducial cross sections are sensitive to  $q_T$  resummation effects
- Enables best-possible combined predictions for other  $H \rightarrow \gamma\gamma$  observables

Thank you for your attention!

Backup



# Uncertainty breakdown



$$N^3LO: \quad \sigma_{fid}/\mathcal{B}_{\gamma\gamma} = (25.16 \pm 1.78_{FO} \pm 0.12_{nons}) \text{ pb}$$

$$N^3LL'+N^3LO: \quad \sigma_{fid}/\mathcal{B}_{\gamma\gamma} = (25.41 \pm 0.59_{FO} \pm 0.21_{q_T} \pm 0.17_{\varphi} \pm 0.06_{match} \pm 0.20_{nons}) \text{ pb}$$

$\Delta_{q_T}$  36 independent scale variations in  $W^{(0)}$  factorization

$\Delta_{\varphi}$  Vary phase of hard scale over  $\arg \mu_H \in \{\pi/4, 3\pi/4\}$

$\Delta_{match}$  Vary transition points governing resummation turn-off

$\Delta_{FO}$  Vary  $\mu_R/m_H \in \{1/2, 2\}$  (dominates over  $\mu_F$  due to overall  $\alpha_s^2$ )

$\Delta_{nons}$  Uncertainty on nonsingular extraction


At leading power in  $q_T \ll m_H$ , the hadronic dynamics factorize as:

$$W^{(0)}(q_T, Y) = H(m_H^2, \mu) \int d^2\vec{k}_a d^2\vec{k}_b d^2\vec{k}_s \delta(q_T - |\vec{k}_a + \vec{k}_b + \vec{k}_s|) \\ \times B_g^{\mu\nu}(x_a, \vec{k}_a, \mu, \nu) B_{g\mu\nu}(x_b, \vec{k}_b, \mu, \nu) S(\vec{k}_s, \mu, \nu)$$

Ingredients satisfy 2D renormalization group equations, e.g. soft function:

$$\mu \frac{d}{d\mu} \ln \tilde{S}(\vec{b}_T, \mu, \nu) = \tilde{\gamma}_S^g(\mu, \nu) \quad \nu \frac{d}{d\nu} \ln \tilde{S}(\vec{b}_T, \mu, \nu) = \tilde{\gamma}_\nu^g(b_T, \mu)$$

- Solve recursively at fixed order
  - ▶ Complete log structure of  $d\sigma^{(0)}$
- Closed-form all-order solution
  - ▶ Resummed Sudakov peak
- Resummation order specified by perturbative order of anom. dims. and boundary conditions



At leading power in  $q_T \ll m_H$ , the hadronic dynamics factorize as:

$$W^{(0)}(q_T, Y) = H(m_H^2, \mu) \int d^2\vec{k}_a d^2\vec{k}_b d^2\vec{k}_s \delta(q_T - |\vec{k}_a + \vec{k}_b + \vec{k}_s|) \\ \times B_g^{\mu\nu}(x_a, \vec{k}_a, \mu, \nu) B_{g\mu\nu}(x_b, \vec{k}_b, \mu, \nu) S(\vec{k}_s, \mu, \nu)$$

To reach  $N^3LL'$  for  $W^{(0)}$ , implemented in SCETlib:

- Three-loop **soft** and **hard** function ... includes in particular the three-loop virtual form factor [Li, Zhu, '16] [Baikov et al. '09; Lee et al. '10; Gehrmann et al. '10]
- Three-loop **unpolarized** and two-loop **polarized beam** functions [Ebert, Mistlberger, Vita '20; Luo, Yang, Zhu, Zhu '20] [Luo, Yang, Zhu, Zhu '19; Gutierrez-Reyes, Leal-Gomez, Scimemi, Vladimirov '19]
- Four-loop cusp, three-loop noncusp anomalous dimensions [Brüser, Grozin, Henn, Stahlhofen '19; Henn, Korchemsky, Mistlberger '20; v. Manteuffel, Panzer, Schabinger '20] [Li, Zhu, '16; Moch, Vermaseren, Vogt '05; Idilbi, Ma, Yuan '06; Vladimirov '16]
- $N^3LL$  solutions to virtuality/rapidity RGEs in  $b_T$  space
- Hybrid profile scales for fixed-order matching [Lustermans, JM, Tackmann, Waalewijn '19]

## Differential $q_T$ subtractions

$$\sigma = \int_0^{q_T^{\text{off}}} dq_T \frac{d\sigma^{\text{sing}}}{dq_T} + \int_0^{q_T^{\text{off}}} dq_T \frac{d\sigma_{\text{FO}}^{\text{nons}}}{dq_T} + \int_{q_T^{\text{off}}} dq_T \frac{d\sigma_{\text{FO}1}}{dq_T}$$

Include  $d\sigma^{\text{fpc}}$  in differential subtraction:

$$\frac{d\sigma^{\text{sing}}}{dq_T} = \int dY A(q_T, Y; \Theta) W^{(0)}(q_T, Y) = \frac{d\sigma^{(0)}}{dq_T} + \frac{d\sigma^{\text{fpc}}}{dq_T}$$

Remaining (nonsingular) terms:

$$\frac{d\sigma_{\text{FO}}^{\text{nons}}}{dq_T} = \int dY A(q_T, Y; \Theta) \left[ W_{\text{FO}}^{(2)}(q_T, Y) + \dots \right] = \left[ \frac{d\sigma_{\text{FO}1}}{dq_T} - \frac{d\sigma_{\text{FO}}^{\text{sing}}}{dq_T} \right]_{q_T > 0}$$

Challenges:

- Obtaining stable  $H + 1j$  results for  $q_T \rightarrow 0$  is *hard* ...in particular at NNLO<sub>1</sub>
- Dropping the nonsingular below  $q_T \leq q_T^{\text{cut}}$  is not viable, either ...as we'll see shortly
  - Crucial to use differential subtraction, not slicing

## Differential $q_T$ subtractions

$$\sigma = \int_0^{q_T^{\text{off}}} dq_T \frac{d\sigma^{\text{sing}}}{dq_T} + \int_0^{q_T^{\text{off}}} dq_T \frac{d\sigma_{\text{FO}}^{\text{nons}}}{dq_T} + \int_{q_T^{\text{off}}} dq_T \frac{d\sigma_{\text{FO}1}}{dq_T}$$

Include  $d\sigma^{\text{fpc}}$  in differential subtraction:

$$\frac{d\sigma^{\text{sing}}}{dq_T} = \int dY A(q_T, Y; \Theta) W^{(0)}(q_T, Y) = \frac{d\sigma^{(0)}}{dq_T} + \frac{d\sigma^{\text{fpc}}}{dq_T}$$

Remaining (nonsingular) terms:

$$\frac{d\sigma_{\text{FO}}^{\text{nons}}}{dq_T} = \int dY A(q_T, Y; \Theta) \left[ W_{\text{FO}}^{(2)}(q_T, Y) + \dots \right] = \left[ \frac{d\sigma_{\text{FO}1}}{dq_T} - \frac{d\sigma_{\text{FO}}^{\text{sing}}}{dq_T} \right]_{q_T > 0}$$



### Key idea

Fit nonsingular data to known form at subleading power and integrate *analytically*:

$$q_T \frac{d\sigma_{\text{FO}}^{\text{nons}}}{dq_T} \Big|_{\alpha_s^n} = \frac{q_T^2}{m_H^2} \sum_{k=0}^{2n-1} \left( a_k + b_k \frac{q_T}{m_H} + c_k \frac{q_T^2}{m_H^2} + \dots \right) \ln^k \frac{q_T^2}{m_H^2}$$

- Include higher-power  $b_k, c_k$  to get unbiased  $a_k$
- ▶ Allows us to use more precise data at higher  $q_T$  as lever arm in the fit



## Fit results at (N)NLO



### Fixed-order inputs:

- NLO contribution to  $W(q_T, Y)$  at  $q_T > 0$  (LO<sub>1</sub>) is easy
- At NNLO (NLO<sub>1</sub>), renormalize & implement bare analytic results for  $W(q_T, Y)$   
[Dulat, Lionetti, Mistlberger, Pelloni, Specchia '17]



# Fit results at (N)LO



## Fit procedure:

- Perform separate  $\chi^2$  fits of  $\{a_k^{\text{incl, fid}}\}$  to inclusive and fiducial nonsingular data [generated by our analytic implementation]
- Increase fit window to larger  $q_T$  until  $p$  value decreases
- Include subleading log coefficients at next higher power until  $p$  value decreases
- Also test intermediate combinations to ensure fit is stable [procedure follows Moutl, Rothen, Stewart, Tackmann, Zhu '15-'16]

# Fit results at (N)LO





## Fit procedure:

- Perform separate  $\chi^2$  fits of  $\{a_k^{\text{incl, fid}}\}$  to inclusive and fiducial nonsingular data [generated by our analytic implementation]
- Increase fit window to larger  $q_T$  until  $p$  value decreases
- Include subleading log coefficients at next higher power until  $p$  value decreases
- Also test intermediate combinations to ensure fit is stable [procedure follows Moutl, Rothen, Stewart, Tackmann, Zhu '15-'16]





# Fit results at (N)NLO



## Fit procedure:

- Perform separate  $\chi^2$  fits of  $\{a_k^{\text{incl, fid}}\}$  to inclusive and fiducial nonsingular data [generated by our analytic implementation]
- Increase fit window to larger  $q_T$  until  $p$  value decreases
- Include subleading log coefficients at next higher power until  $p$  value decreases
- Also test intermediate combinations to ensure fit is stable [procedure follows Moutl, Rothen, Stewart, Tackmann, Zhu '15-'16]



# Fit results at (N)NLO



## Fit procedure:

- Perform separate  $\chi^2$  fits of  $\{a_k^{\text{incl, fid}}\}$  to inclusive and fiducial nonsingular data [generated by our analytic implementation]
- Increase fit window to larger  $q_T$  until  $p$  value decreases
- Include subleading log coefficients at next higher power until  $p$  value decreases
- Also test intermediate combinations to ensure fit is stable [procedure follows Moutl, Rothen, Stewart, Tackmann, Zhu '15-'16]

# Fit results at (N)NLO





- Check the purely hadronic  $a_k^{\text{fid}}$  by directly fitting them to

$$q_T \int dY A^{(0)}(Y; \Theta) [W - W^{(0)}] = \frac{q_T^2}{m_H^2} \sum_{k=0}^{2n-1} \left( a_k^{\text{fid}} + c'_k \frac{q_T^2}{m_H^2} + \dots \right) \ln^k \frac{q_T^2}{m_H^2} \quad \checkmark$$

- Recover analytic (N)NLO coefficient of  $\sigma_{\text{incl}}$  at  $10^{-5}$  ( $10^{-4}$ )  $\checkmark$
- Analytic implementation gives us awesome precision on *all* NLP coefficients (all logs at NLO *and* NNLO, also differential in  $Y$ , broken down by color structure, ...)
- ▶ Can serve as benchmark for  $q_T$  resummation at subleading power


# Fit results at N<sup>3</sup>LO



## Setup:

- Combined fit to existing binned inclusive and fiducial NNLO<sub>1</sub> data from NNLOjet [Chen, Cruz-Martinez, Gehrmann, Glover, Jaquier '15-16; as used in Chen et al. '18; Bizoń et al. '18]
- Empirically find  $0.4 \leq a_k^{\text{fid}}/a_k^{\text{incl}} \leq 0.55$  at (N)NLO  $\Rightarrow$  use as weak 1 $\sigma$  constraint
- Add  $\sigma_{\text{incl}}(q_T \leq q_T^{\text{cut}}) = \sigma_{\text{incl}}^{\text{N}^3\text{LO}} - \sigma_{\text{incl}}(q_T > q_T^{\text{cut}})$  as additional incl. data point [Mistlberger '18]

## Comparison to other methods: $q_T$ slicing




### Slicing approach to $q_T$ subtractions:

[used e.g. in Cieri, Chen, Gehrmann, Glover, Huss, 1807.11501; Camarda, Cieri, Ferrera, 2103.04974]

$$\sigma = \sigma^{(0)}(q_T^{\text{cut}}) + \sigma^{\text{fpc}}(q_T^{\text{cut}}) + \sigma^{\text{nons}}(q_T^{\text{cut}}) + \int_{q_T^{\text{cut}}} dq_T \frac{d\sigma_{\text{FO1}}}{dq_T}$$

- Slicing uses finite  $q_T^{\text{cut}} \sim 2$  GeV and neglects both  $\sigma^{\text{fpc}}(q_T^{\text{cut}})$ ,  $\sigma^{\text{nons}}(q_T^{\text{cut}}) \approx 0$
- This is a catastrophic approximation even at  $\alpha_s^2$ , and definitely at  $\alpha_s^3$
- Even without  $\sigma^{\text{fpc}}$  (e.g., without cuts), this is a bad approximation at  $\alpha_s^3$ 
  - $q_T^{\text{cut}}$  variations only scan local maximum around 2 GeV ...

## Comparison to other methods: Projection to Born



### Projection-to-Born method:

[used e.g. in Chen, Gehrmann, Glover, Huss, Mistlberger, Pelloni, 2102.07607]

$$\frac{d\sigma}{dY} = A(0, Y) \frac{d\sigma_{\text{incl}}}{dY} + \int_{\approx q_T^{\text{cut}}} dq_T [A(q_T, Y) - A(0, Y)] W(q_T, Y)$$

- First term from analytic (threshold expansion of) inclusive rapidity spectrum
- Second term numerically from  $H + 1j$  MC, dominated by  $\sigma^{\text{fpc}}$  at small  $q_T$
- ▶ Need to integrate down to  $q_T^{\text{cut}} \ll 0.1 \text{ GeV}$  to get error below 10% of  $\sigma_{\text{LO}}^{\text{fid}}$ !  
[See also Salam, Slade, 2106.08329 for an explicit/analytic estimate at double-logarithmic level]