Contribution ID: 306

Type: talk

## A Novel Scintillator Detector for the Mu2e-II Experiment and a Muon Tomography Probe of the Interior of the Great Pyramid

Tuesday 13 July 2021 16:45 (15 minutes)

The Mu2e experiment is designed to search for the charged-lepton-flavor-violating process,  $\mu^-$  to a  $e^-$ , with unprecedented sensitivity. The single 105-MeV electron that results from this process can be mimicked by electrons produced by cosmic-ray muons traversing the detector. An active veto detector surrounding the apparatus is used to detect incoming cosmic-ray muons. To reduce the backgrounds to the required level it must have an efficiency of about 99.99\% as well as excellent hermeticity. The detector consists of four layers of scintillator counters, each with two embedded wavelength-shifting fibers, whose light is detected by silicon photomultipliers. An upgrade of the experiment, Mu2e-II, that will provide an order of magnitude more sensitivity is under design. The cosmic-ray veto detector is being redesigned to handle the higher rates. This redesign is also being used for a proposed high-resolution probe of the interior of the Great Pyramid of Khufu. The design and expected performance of the detector will be described.

## Are you are a member of the APS Division of Particles and Fields?

Yes

Author:DUKES, EdmondPresenter:DUKES, EdmondSession Classification:Particle Detectors

Track Classification: Particle Detectors