Contribution ID: 205

Type: talk

## Higgs to two b's or not two b's: The measurement for a Standard Model Higgs boson, prooduced in association with a W or Z boson, decaying into a pair of b-quarks in pp collisions at 13 TeV with the ATLAS detector

Monday, 12 July 2021 15:15 (15 minutes)

Following the discovery of the Higg's boson in 2012 by both the ATLAS and CMS experiments, a wealth of papers have been published concerning measurements or observations of the Higgs' decay modes. However, the most dominant decay mode,  $H \rightarrow b\bar{b}$ , proved to be an elusive and challenging search due to the low signal-to-background environment, and a diverse range of backgrounds arising from multiple Standard Model processes. The backgrounds include W+jets, Z+jets, and  $t\bar{t}$  production amongst others. Measurements of the WH and ZH production, with the W or Z boson decaying into charged leptons (electrons or muons, including those produced from the leptonic decay of a tau lepton), in the  $H \rightarrow b\bar{b}$  decay channel in pp collisions at 13 TeV, corresponding to an integrated luminosity of 139 fb<sup>-1</sup>, with the ATLAS detector was performed. The production of a Higgs boson in association with a W or Z boson has been established with observed (expected) significances of 4.0 (4.1) and 5.3 (5.1) standard deviations, respectively.

## Are you are a member of the APS Division of Particles and Fields?

No

Primary author: Ms TEE, Amy (University of Wisconsin Madison)Presenter: Ms TEE, Amy (University of Wisconsin Madison)Session Classification: Higgs & Electroweak Physics

Track Classification: Higgs & Electroweak Physics