

Search for Neutrinoless Double Beta Decay of ^{76}Ge with GERDA - final latest results

Friday 14 January 2022 12:20 (15 minutes)

The GERDA experiment, located in the underground Laboratori Nazionali del Gran Sasso in Italy, has been designed to search for the neutrinoless double-beta ($0\nu\beta\beta$) decay in ^{76}Ge . It used in different stages of the project up to 44 kg of high purity germanium (HPGe) detectors enriched up to about 86% in the isotope ^{76}Ge . The bare detectors were operated in liquid argon, which served in the first phase as a passive, and later, in the second phase of the experiment, also as an active shield. The combination of powerful background suppression techniques (liquid argon veto and pulse shape discrimination) together with excellent energy resolution of HPGe detectors allowed GERDA to be the first background-free (less than 1 background event expected in the region of interest) $0\nu\beta\beta$ decay experiment. After about 5 years of data taking together in Phase I and Phase II more than 100 kg*yr of data has been accumulated, as foreseen in the original proposal. Combining all available data no signal has been observed, and a lower limit on the half-life of the $0\nu\beta\beta$ decay in ^{76}Ge is set at $T_{1/2} > 1.8 \times 10^{26}$ yr at 90% C.L, which is presently the strongest limitation. It may be translated into an upper limit of the effective neutrino mass of $m_{ee} < (79 - 120)$ meV, depending on the used matrix elements. In the talk the design of the experiment and details of the data analysis will be presented. A prospect for the successor Legend-200 experiment will be outlined as well.

Author: ZUZEL, Grzegorz (Jagiellonian University)

Presenter: ZUZEL, Grzegorz (Jagiellonian University)

Session Classification: Astrophysical neutrinos II