



Universität
Basel

Nucleon decay fingerprints from SUSY GUT models (using SusyTCProton)

Vasja Susič

University of Basel

with Stefan Antusch and Christian Hohl

Portorož 2021 — Physics of the flavourful Universe,

2021-09-24

Introduction

Motivation:

- Is there a process generically predicted by SUSY GUTs?
Can it be used to distinguish between models?

Nucleon (proton and neutron) decay!

- Need software for completely general computation (including all flavor dependence, etc.).

Based on arXiv:2011.15026.

Introduction

Motivation:

- Is there a process generically predicted by SUSY GUTs?
Can it be used to distinguish between models?

Nucleon (proton and neutron) decay!

- Need software for completely general computation (including all flavor dependence, etc.).


Contents of the talk:

- 1 A quick recap of $D = 5$ proton decay in SUSY
- 2 Example models and results
- 3 Software for proton decay calculation: `SusyTCP`Proton

Based on arXiv:2011.15026.


Notes on $D = 5$ proton decay — part 1

$D = 6$ decay (example):



$$\mathcal{A}_{D=6} \approx \frac{4\pi\alpha}{m_X^2}$$

$D = 5$ decay (example):




$$\mathcal{A}_{D=5} \approx \frac{\alpha}{4\pi} y_u y_d \frac{m_{\tilde{w}}}{m_T m_{\tilde{f}}^2}$$

- Generic form for proton decay operators: $qqql$ ($B - L = 0$)
- $D = 5$: **requires SUSY and triplets**... SM superpartners in loops
 \tilde{w} dominates, complicated **flavor** structure


Notes on $D = 5$ proton decay — part 1

$D = 6$ decay (example):



$$\mathcal{A}_{D=6} \approx \frac{4\pi\alpha}{m_X^2}$$

$D = 5$ decay (example):



$$\mathcal{A}_{D=5} \approx \frac{\alpha}{4\pi} y_u y_d \frac{m_{\tilde{w}}}{m_T m_f^2}$$

$$\Gamma \approx |\mathcal{A}|^2 m_p^5 \Rightarrow \Gamma_5 \sim \frac{y^2}{(4\pi)^2} \frac{m_p^5}{M_{\text{GUT}}^2 M_{\text{SUSY}}^2} > \Gamma_6 \sim \frac{m_p^5}{M_{\text{GUT}}^4}$$

- Generic form for proton decay operators: $qqql$ ($B - L = 0$)
- $D = 5$: requires SUSY and triplets... SM superpartners in loops
 \tilde{w} dominates, complicated flavor structure

Notes on $D = 5$ proton decay — part 2

- Yukawa operators in $SU(5)$: also contain triplets $T \sim (3, 1, -1/3)$

$$\begin{aligned} \mathbf{Y}_{\bar{5}} \mathbf{10}_F \bar{\mathbf{5}}_F \bar{\mathbf{X}} &\sim \mathbf{Y}_d Q d^c H_d + \mathbf{Y}_e L e^c H_d + \tilde{\mathbf{Y}}_{ql} Q L \bar{T} + \tilde{\mathbf{Y}}_{ud} u^c d^c \bar{T} \\ \mathbf{Y}_{10} \mathbf{10}_F \mathbf{10}_F \mathbf{X} &\sim \mathbf{Y}_u Q u^c H_u + \tilde{\mathbf{Y}}_{qq} Q Q T + \tilde{\mathbf{Y}}_{eu} e^c u^c T \end{aligned}$$

Above: $\mathbf{X} \in \{\mathbf{5}, \mathbf{45}\}$. Can also use non-renormalizable operators.

Notes on $D = 5$ proton decay — part 2

- Yukawa operators in $SU(5)$: also contain triplets $T \sim (3, 1, -1/3)$

$$\begin{aligned} \mathbf{Y}_{\bar{5}} \mathbf{10}_F \bar{\mathbf{5}}_F \bar{\mathbf{X}} &\sim \mathbf{Y}_d Q d^c H_d + \mathbf{Y}_e L e^c H_d + \tilde{\mathbf{Y}}_{ql} Q L \bar{T} + \tilde{\mathbf{Y}}_{ud} u^c d^c \bar{T} \\ \mathbf{Y}_{10} \mathbf{10}_F \mathbf{10}_F \mathbf{X} &\sim \mathbf{Y}_u Q u^c H_u + \tilde{\mathbf{Y}}_{qq} Q Q T + \tilde{\mathbf{Y}}_{eu} e^c u^c T \end{aligned}$$

Above: $\mathbf{X} \in \{\mathbf{5}, \mathbf{45}\}$. Can also use non-renormalizable operators.

- Effective Lagrangian for $D = 5$ nucleon decay:

$$\begin{aligned} W_5 &= \mathbf{C}^{5L} Q L Q Q & \mathbf{C}_{ijkl}^{5L} &= (\mathbf{M}_T^{-1})_{IJ} (\tilde{\mathbf{Y}}_{ql})_{Iij} (\tilde{\mathbf{Y}}_{qq})_{Jkl} \\ &+ \mathbf{C}^{5R} u^c d^c e^c u^c & \mathbf{C}_{ijkl}^{5R} &= (\mathbf{M}_T^{-1})_{IJ} (\tilde{\mathbf{Y}}_{ud})_{Iij} (\tilde{\mathbf{Y}}_{eu})_{Jkl} \end{aligned}$$

- To describe nucleon decay: add \mathbf{C}^{5L} and \mathbf{C}^{5R} to MSSM
- In each operator: some GUT relation between \mathbf{Y} and $\tilde{\mathbf{Y}}$

Nucleon decay channels — experimental bounds

Proton decay:


decay channel	τ/\mathcal{B} [year]
$p \rightarrow K^+ \bar{\nu}$	$> 6.6 \cdot 10^{33}$
$p \rightarrow K^0 e^+$	$> 1.1 \cdot 10^{33}$
$p \rightarrow K^0 \mu^+$	$> 1.6 \cdot 10^{34}$
$p \rightarrow \pi^+ \bar{\nu}$	$> 2.8 \cdot 10^{32}$
$p \rightarrow \pi^0 e^+$	$> 2.4 \cdot 10^{34}$
$p \rightarrow \pi^0 \mu^+$	$> 1.6 \cdot 10^{34}$
$p \rightarrow \eta^0 e^+$	$> 4.1 \cdot 10^{33}$
$p \rightarrow \eta^0 \mu^+$	$> 1.2 \cdot 10^{33}$

Neutron decay:

decay channel	τ/\mathcal{B} [year]
$n \rightarrow K^0 \bar{\nu}$	$> 1.2 \cdot 10^{32}$
$n \rightarrow \pi^0 \bar{\nu}$	$> 9.9 \cdot 10^{32}$
$n \rightarrow \pi^- e^+$	$> 2.1 \cdot 10^{33}$
$n \rightarrow \pi^- \mu^+$	$> 9.9 \cdot 10^{32}$
$n \rightarrow \eta^0 \bar{\nu}$	$> 5.6 \cdot 10^{32}$


- Table is from arXiv:2011.15026, bounds mostly from Super-K
- Generic form: **nucleon** \rightarrow **meson** + **lepton**
- $D = 5$ decay favors K over π (2nd family quarks involved)

Model setup: flavor SUSY GUT



- We are in $SU(5)$, we model Yukawa sector but not breaking sector
- Assume **single operator dominance**:
each Yukawa-matrix entry dominated by one non-renormalizable operator (generated by integrating out mediators, cf. 0902.4644)

Model setup: flavor SUSY GUT



- We are in $SU(5)$, we model Yukawa sector but not breaking sector
- Assume **single operator dominance**:
each Yukawa-matrix entry dominated by one non-renormalizable operator (generated by integrating out mediators, cf. 0902.4644)
- M_{GUT} :
 - GUT relations between \mathbf{Y} (and $\tilde{\mathbf{Y}}$ relevant for $\mathbf{C}^{5L,R}$)
 - mSUGRA boundary conditions
 - GUT-breaking sector: we assume all Higgs fields lie at M_{GUT}
- M_{SUSY} : scale of SM superpartners
- m_W : fit to low energy data
- We compare two example models (from 1405.6962)...

Example models — Yukawa textures

Model 1:

$$\mathbf{Y}_{10} = \begin{pmatrix} * & * & 0 \\ * & * & * \\ 0 & * & * \end{pmatrix}, \mathbf{Y}_{\bar{5}} = \begin{pmatrix} 0 & * & 0 \\ * & * & 0 \\ 0 & 0 & * \end{pmatrix}$$

\mathbf{Y} and $\tilde{\mathbf{Y}}$ relations:

$$\mathbf{Y}_e = \mathbf{Y}_d^T \cdot \begin{pmatrix} 0 & 6 & 0 \\ -1/2 & 6 & 0 \\ 0 & 0 & -3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ql} = \mathbf{Y}_d \cdot \begin{pmatrix} 0 & -1 & 0 \\ -1 & -1 & 0 \\ 0 & 0 & 3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ud} = \mathbf{Y}_d \cdot \begin{pmatrix} 0 & 2/3 & 0 \\ -4 & -4 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Model 2:

$$\mathbf{Y}_{10} = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix}, \mathbf{Y}_{\bar{5}} = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix}$$

\mathbf{Y} and $\tilde{\mathbf{Y}}$ relations:

$$\mathbf{Y}_e = \mathbf{Y}_d^T \cdot \begin{pmatrix} -1/2 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & -3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ql} = \mathbf{Y}_d \cdot \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ud} = \mathbf{Y}_d \cdot \begin{pmatrix} 2/3 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Example models — Yukawa textures

Model 1:

$$\mathbf{Y}_{10} = \begin{pmatrix} * & * & 0 \\ * & * & * \\ 0 & * & * \end{pmatrix}, \mathbf{Y}_{\bar{5}} = \begin{pmatrix} 0 & * & 0 \\ * & * & 0 \\ 0 & 0 & * \end{pmatrix}$$

\mathbf{Y} and $\tilde{\mathbf{Y}}$ relations:

$$\mathbf{Y}_e = \mathbf{Y}_d^T \cdot \begin{pmatrix} 0 & 6 & 0 \\ -1/2 & 6 & 0 \\ 0 & 0 & -3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ql} = \mathbf{Y}_d \cdot \begin{pmatrix} 0 & -1 & 0 \\ -1 & -1 & 0 \\ 0 & 0 & 3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ud} = \mathbf{Y}_d \cdot \begin{pmatrix} 0 & 2/3 & 0 \\ -4 & -4 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Model 2:

$$\mathbf{Y}_{10} = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix}, \mathbf{Y}_{\bar{5}} = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix}$$

\mathbf{Y} and $\tilde{\mathbf{Y}}$ relations:

$$\mathbf{Y}_e = \mathbf{Y}_d^T \cdot \begin{pmatrix} -1/2 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & -3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ql} = \mathbf{Y}_d \cdot \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 3/2 \end{pmatrix}$$

$$\tilde{\mathbf{Y}}_{ud} = \mathbf{Y}_d \cdot \begin{pmatrix} 2/3 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Also: $\tilde{\mathbf{Y}}_{qq} = -\mathbf{Y}_u$ and $\tilde{\mathbf{Y}}_{eu} = +\mathbf{Y}_u$ (both models). No neutrino sector.

- Factors come from choice of GUT operator in each entry.
- Construction for \mathbf{M}_T : multiple T , one only couples to fermions
 $\Rightarrow M_T^{\text{eff}} = ((\mathbf{M}_T^{-1})_{11})^{-1}$ effective T -mass (unrelated to a physical scale)

Numerical analysis of the models

- At $M_{\text{GUT}} = 2 \cdot 10^{16}$ GeV we impose boundary conditions:
 - Yukawa inputs: we only need to specify \mathbf{Y}_d and \mathbf{Y}_u
 - Fix CMSSM/mSUGRA parameters with $\text{sgn}(\mu) = +1$:

$$\tan \beta = 28.4, \quad m_0 = 4 \text{ TeV}, \quad M_{1/2} = 1.7 \text{ TeV}, \quad A_0 = -10 \text{ TeV}.$$

- Use SusyTCProton \Rightarrow obtain mass and mixing predictions at m_W
 \Rightarrow compute χ^2


Numerical analysis of the models

- At $M_{\text{GUT}} = 2 \cdot 10^{16}$ GeV we impose boundary conditions:
 - Yukawa inputs: we only need to specify \mathbf{Y}_d and \mathbf{Y}_u
 - Fix CMSSM/mSUGRA parameters with $\text{sgn}(\mu) = +1$:


$$\tan \beta = 28.4, \quad m_0 = 4 \text{ TeV}, \quad M_{1/2} = 1.7 \text{ TeV}, \quad A_0 = -10 \text{ TeV}.$$

- Use SusyTCProton \Rightarrow obtain mass and mixing predictions at m_W
 \Rightarrow compute χ^2
- Careful analysis of parameters shows:
 - 14(+3) inputs vs 14 outputs for χ^2
 - **inputs**: 10(+3) in Yukawa sector (varied), 4 soft parameters (fixed)
 - **outputs**: fermion masses and CKM mixings, SM Higgs mass
 - The +3 inputs: the “GUT phases” $\phi_1, \phi_2, M_T^{\text{eff}}$
Influence nucleon decay, no effect on SM observables!
- Best fit: $\chi^2 \approx 5$ for both models

Results — GUT-phase dependence of $p \rightarrow K^+ \bar{\nu}$ and $n \rightarrow K^0 \bar{\nu}$




Results — decay rates ($D = 5$ only)



- Perform MCMC around best fit point: vary Yukawa parameters.
- M_T^{eff} is an input: experimental bounds avoided if raised.
Benchmark value: $M_T^{\text{eff}} = 10^{19} \text{ GeV}$.


Results — decay rate ratios ($D = 5$ only)



- Perform MCMC around best fit point: vary Yukawa parameters.
- Decay ratios **fingerprints**: models can be distinguished!

Proton decay computation — procedure


Procedure: follow Goto-Nihei (also: review by Nath-Perez):



Step by step:

- (1) M_{GUT} : integrate out heavy triplets T, \bar{T} to obtain C^{5L} and C^{5R} .
- (2) RGE : run $C^{5L,R}$ from GUT to SUSY scale.
- (3) M_{SUSY} : dress the $C^{5L,R}$ diagrams with MSSM superpartners. One gets four-fermion operators C^6 .
- (4) RGE : evolve C^6 from SUSY scale to proton scale.
- (5) M_{proton} : computes decay rates from C^6 (chiral perturbation theory).

Proton decay computation — software



Our software: 2 Mathematica packages.

SusyTCProton:

- Improved version of SusyTC. Based on REAP.
- Inputs: values at GUT scale (including C^{5L} and C^{5R})
- Performs **RGE** for **MSSM + $C^{5L,R}$** , computes **SUSY spectrum**.

ProtonDecay:

- Input: from SusyTCProton or extended SLHA (add $C^{5L,R}$).
- Computes **decay rates**: **8 proton** channels, **5 neutron** channels.
- Extra feature: C^6 inputs. Can be used as a standalone.

Conclusions

- 1 **Nucleon decay**: an important process to consider in SUSY GUT
 - Limitation on models
 - Can be used to distinguish models
- 2 Decay ratios (aka **decay fingerprint**) can give important insight
- 3 **Example**: 2 flavor SUSY GUT models
(flavor structure can indeed have an important effect)
- 4 Public software for ($D = 5$) decay: **SusyTCProton**

Conclusions

- 1 **Nucleon decay**: an important process to consider in SUSY GUT
 - Limitation on models
 - Can be used to distinguish models
- 2 Decay ratios (aka **decay fingerprint**) can give important insight
- 3 **Example**: 2 flavor SUSY GUT models
(flavor structure can indeed have an important effect)
- 4 Public software for ($D = 5$) decay: **SusyTCProton**

Thank you for your attention!