

Interplay of New Physics Effects in $(g - 2)_\ell$ and $h \rightarrow \ell^+ \ell^-$ Lessons from SMEFT

Michele Tammaro


@ Portoroz, 21/09/2021

Based on 2103.10859 with J.F. Kamenik and S. Fajfer



RG running
←

$$\mathcal{L}_{EFT} = \sum_{d,i} \frac{C_i^{(d)}}{\Lambda^{d-4}} \mathcal{O}_i^{(d)}$$



LEFT
Only light fermions, gluons and photons (1709.04486)

5-flavor EFT
Top integrated out

SMEFT
Warsaw basis (1008.4884)

Complete one-loop RGE in SMEFT
(Jenkins et al.: 1312.2014, 1310.4838, 1312.2014)


One-loop SMEFT matching into LEFT
(Dekens, Stoffer: 1908.05295)

Spoilers

- Small set of operators in SMEFT describes the observables
- Correlations via one-loop RGE
- Chirally enhanced matching conditions at one-loop
- Pheno analysis can give hints on model building
- Tree-level matching + running can capture the dominant effects of NP


$$h \rightarrow l^+ l^-$$

$$\mathcal{O}_{1,pr} = (\varphi^\dagger \varphi) (\bar{l}_p e_r \varphi)$$



SMEFT

$$\mathcal{Q}_{1,\ell} = c_{1,\ell} h (\bar{l}_\ell l_\ell)$$




LEFT

Exp

$$\kappa_\ell^2 = \frac{\Gamma(h \rightarrow l^+ l^-)}{\Gamma_{SM}(h \rightarrow l^+ l^-)} = \left(1 + \frac{c_{1,\ell} v}{m_\ell} \right)^2$$


$$h \rightarrow l^+ l^-$$

$$\mathcal{O}_{1,pr} = (\varphi^\dagger \varphi) (\bar{l}_p e_r \varphi)$$




SMEFT

$$\mathcal{O}_{4,prst} = (\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$$



LEFT

$$\mathcal{Q}_{1,\ell} = c_{1,\ell} h (\bar{l}_\ell l_\ell)$$




Exp

$$\kappa_\ell^2 = \frac{\Gamma(h \rightarrow l^+ l^-)}{\Gamma_{SM}(h \rightarrow l^+ l^-)} = \left(1 + \frac{c_{1,\ell} v}{m_\ell} \right)^2$$

$$(g - 2)_\ell$$

$$\mathcal{O}_{2,pr} = (\bar{\ell}_p \sigma^{\mu\nu} e_r) \tau^a \varphi W_{\mu\nu}^a$$


$$\mathcal{O}_{3,pr} = (\bar{\ell}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$$



SMEFT

LEFT

$$Q_{2,\ell} = e \frac{c_{2,\ell}}{m_\ell} (\bar{l}_\ell \sigma_{\mu\nu} l_\ell) F^{\mu\nu}$$




Exp

$$\delta a_\ell = 4c_{2,\ell}(m_\ell)$$

$$(g - 2)_\ell$$


$$\mathcal{O}_{2,pr} = (\bar{\ell}_p \sigma^{\mu\nu} e_r) \tau^a \varphi W_{\mu\nu}^a$$

$$\mathcal{O}_{3,pr} = (\bar{\ell}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$$




SMEFT

$$\mathcal{O}_{5,prst} = (\bar{\ell}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$$



LEFT

$$\mathcal{Q}_{2,\ell} = e \frac{c_{2,\ell}}{m_\ell} (\bar{l}_\ell \sigma_{\mu\nu} l_\ell) F^{\mu\nu}$$



Exp

$$\delta a_\ell = 4c_{2,\ell}(m_\ell)$$

Pheno guidelines

LEFT

$$\mathcal{Q}_{1,\ell} = c_{1,\ell} h (\bar{l}_\ell l_\ell)$$

$$\mathcal{Q}_{2,\ell} = e \frac{c_{2,\ell}}{m_\ell} (\bar{l}_\ell \sigma_{\mu\nu} l_\ell) F^{\mu\nu}$$

SMEFT

$$\mathcal{O}_{1,pr} = (\varphi^\dagger \varphi) (\bar{l}_p e_r \varphi)$$

$$\mathcal{O}_{2,pr} = (\bar{l}_p \sigma^{\mu\nu} e_r) \tau^a \varphi W_{\mu\nu}^a$$

$$\mathcal{O}_{3,pr} = (\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$$

$$\mathcal{O}_{4,prst} = (\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$$

$$\mathcal{O}_{5,prst} = (\bar{l}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$$

1 - Assume initial condition for each Λ : $\mathcal{C}_i(\Lambda) = 1$, $\mathcal{C}_{j \neq i}(\Lambda) = 0$

2 - RGE running from heavy scale to weak scale

3 - Match SMEFT with LEFT including one-loop

4 - Compute NP effects in observables as function of Λ

$$\kappa_\ell^2 = \frac{\Gamma(h \rightarrow \ell^+ \ell^-)}{\Gamma_{SM}(h \rightarrow \ell^+ \ell^-)} = \left(1 + \frac{c_{1,\ell v}}{m_\ell} \right)^2$$


$$\delta a_\ell = 4c_{2,\ell}(m_\ell)$$

Muon

LHC (CMS) $\kappa_{\mu}^2 = 1.19 \pm 0.55$

HL-LHC $|\kappa_{\mu}^2 - 1| < 0.053$

FCC $|\kappa_{\mu}^2 - 1| < 0.0042$




Picture from Fodor's talk



Theory Initiative

$$\delta a_\mu = (251 \pm 59) \times 10^{-11}$$

BMW lattice

$$\delta a_\mu = (113 \pm 68) \times 10^{-11}$$





- $C_{1,\mu}(\Lambda)=1$
- $C_{2,\mu}(\Lambda)=1$
- $C_{3,\mu}(\Lambda)=1$
- $C_{4,\mu}(\Lambda)=1$
- $C_{5,\mu}(\Lambda)=1$

Top-down approach

or

How effective is EFT?



1 - Two-Higgs Doublet Model

2 - Scalar LeptoQuarks

2HDM

Two scalar doublets with hypercharge +1


$$\mathcal{L}_Y \supset \left(\frac{\eta + H^0}{\sqrt{2}} \right) [y'_t \bar{t} P_R t + y'_\mu \bar{\mu} P_R \mu] + H^+ [y'_t V_{ti} \bar{t} P_L d_i + y'_\mu \bar{\nu}_\mu P_R \mu] + \text{h.c.},$$



2HDM


(g-2) dominated by two-loop Barr-Zee diagrams (Ilisie: 1502.04199)

$$\varphi^0 = \{h, H^0, \eta\}$$



Tree-level matching to SMEFT

$$\hat{C}_{4,l} = \frac{y'_t y'_l}{m_H^2}$$




2HDM

(g-2) dominated by two-loop Barr-Zee diagrams (Ilisie: 1502.04199)


$$\varphi^0 = \{h, H^0, \eta\}$$

Tree-level matching to SMEFT




$$\hat{C}_{4,l} = \frac{y'_t y'_l}{m_H^2}$$

$$y'_t y'_\mu = 1$$




$$m_H = \Lambda = 10^4 \text{ GeV}$$





2HDM

Higgs decay at one-loop from neutral heavy scalar

Tree-level matching to SMEFT



$$\hat{C}_{4,l} = \frac{y'_t y'_l}{m_H^2}$$





$$S_1 = (\bar{\mathbf{3}}, \mathbf{1}, 1/3)$$

$$R_2 = (\mathbf{3}, \mathbf{2}, 7/6)$$

$$\mathcal{L}_{S_1} \supset y_{1,ij}^{LL} \bar{q}^{C,ia} S_1 \epsilon^{ab} \ell^{j,b} + y_{1,ij}^{RR} \bar{u}^{C,i} S_1 \epsilon^{ab} e^j + \text{h.c.}$$


$$\mathcal{L}_{R_2} \supset -y_{2,ij}^{RL} \bar{u}^i R_2^a \epsilon^{ab} \ell^{j,b} + y_{2,ij}^{LR} \bar{e}^i R_2^a * q^{j,a} + \text{h.c.}$$



Scalar LQ

Calculated in: 2008.02643


By Fierz transformation, will generate



$$\mathcal{L} \supset -\frac{4G_F}{\sqrt{2}} \left[g_{ij,ks}^{LL} \left(\bar{q}_L^i q_R^j \right) \left(\bar{\ell}_L^k \ell_R^s \right) + h_{ij,ks}^{LL} \left(\bar{q}_L^i \sigma_{\mu\nu} q_R^j \right) \left(\bar{\ell}_L^k \sigma^{\mu\nu} \ell_R^s \right) \right]$$

$$S_1 : g_{ij,ks}^{LL} = -4h_{ij,ks}^{LL} = \frac{v^2}{4m_{LQ}^2} y_{1,js}^{RR} \left(y_1^{LL} \right)_{ik}^*$$

$$R_2 : g_{ij,ks}^{LL} = 4h_{ij,ks}^{LL} = -\frac{v^2}{4m_{LQ}^2} y_{2,jk}^{RL} \left(y_2^{LR} \right)_{si}^*$$







Tree-level matching to SMEFT

$$\hat{\mathcal{C}}_{4,\mu} = \mp 4\hat{\mathcal{C}}_{5,\mu} = -\frac{4G_F}{\sqrt{2}} g^{LL} = \frac{v^2 G_F}{\sqrt{2} m_{LQ}^2} Y$$

Scalar LQ

Calculated in: 2008.02643

By Fierz transformation, will generate




Summary

- Small set of operators in SMEFT describes the observables
- Correlations via one-loop RGE
- Chirally enhanced matching conditions at one-loop
- Pheno analysis can give hints on model building
- Tree-level matching + running can capture the dominant effects of NP

Backup

$$R_{i,j}(\mu_w) = \frac{c_{i,\ell}(\mu_w)|_j}{c_{i,\ell}(m_t)|_L}, \quad i = 1, 2$$

$$C_{4,\ell}(10^4 \text{ GeV})=1$$



$$\begin{aligned}
\dot{\mathcal{C}}_{1,\ell} &= \left[-\frac{27}{4}g_2^2 - 3(3Y_\ell^2 + 3Y_e^2 - 4Y_\ell Y_e)g_1^2 + 3N_c y_t^2 + 24\lambda \right] \mathcal{C}_{1,\ell} + 4N_c y_t (y_t^2 - \lambda) \mathcal{C}_{4,\ell} \\
&\quad - 3(4g_1^2 g_2 Y_h (Y_e + Y_\ell) + 3g_2^3) \mathcal{C}_{2,\ell} - 6(4g_1^3 Y_h^2 (Y_e + Y_\ell) + g_2^2 g_1 Y_h) \mathcal{C}_{3,\ell}, \\
\dot{\mathcal{C}}_{2,\ell} &= \left[(3c_{F,2} - b_{0,2})g_2^2 + (-3Y_e^2 + 8Y_e Y_\ell - 3Y_\ell^2)g_1^2 + N_c y_t^2 \right] \mathcal{C}_{2,\ell} \\
&\quad + g_1 g_2 (3Y_\ell - Y_e) \mathcal{C}_{3,\ell} - 2g_2 N_c y_t \mathcal{C}_{5,\ell}, \\
\dot{\mathcal{C}}_{3,\ell} &= \left[-3c_{F,2}g_2^2 + (3Y_e^2 + 4Y_e Y_\ell + 3Y_\ell^2 - b_{0,1})g_1^2 + N_c y_t^2 \right] \mathcal{C}_{3,\ell} \\
&\quad + 4c_{F,2}g_1 g_2 (3Y_\ell - Y_e) \mathcal{C}_{2,\ell} + 4g_1 N_c y_t (Y_u + Y_q) \mathcal{C}_{5,\ell}, \\
\dot{\mathcal{C}}_{4,\ell} &= - \left[6(Y_e^2 + Y_e(Y_u - Y_q) + Y_q Y_u)g_1^2 + 3 \left(N_c - \frac{1}{N_c} \right) g_3^2 + y_t^2 (2N_c + 1) \right] \mathcal{C}_{4,\ell} \\
&\quad - \left[24(Y_q + Y_u)(2Y_e - Y_q + Y_u)g_1^2 - 18g_2^2 \right] \mathcal{C}_{5,\ell}, \\
\dot{\mathcal{C}}_{5,\ell} &= g_1(Y_q + Y_u)y_t \mathcal{C}_{3,\ell} - \frac{3}{2}g_2 y_t \mathcal{C}_{2,\ell} + \left[2(Y_e^2 - Y_e Y_q + Y_e Y_u - 2Y_q^2 + 5Y_q Y_u - 2Y_u^2)g_1^2 \right. \\
&\quad \left. - 3g_2^2 + \left(N_c - \frac{1}{N_c} \right) g_3^2 + y_t^2 \right] \mathcal{C}_{5,\ell} + \frac{1}{8} \left[-4(Y_q + Y_u)(2Y_e - Y_q + Y_u)g_1^2 + 3g_2^2 \right] \mathcal{C}_{4,\ell}.
\end{aligned}$$

2HDM

Two scalar doublets with hypercharge +1

$$\begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix} = R_\beta \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix}$$

$$\Phi_1 = \begin{pmatrix} G^+ \\ \frac{v + \rho_1 + iG_0}{\sqrt{2}} \end{pmatrix} \quad \Phi_2 = \begin{pmatrix} H^+ \\ \frac{\rho_2 + i\eta}{\sqrt{2}} \end{pmatrix}$$

$$\begin{pmatrix} h \\ H^0 \end{pmatrix} = R_\alpha \begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix}$$

$$v = \sqrt{v_1^2 + v_2^2}$$

Decoupling limit

$$\alpha, \beta \rightarrow 0$$

$$v_2 \rightarrow 0$$

$$g_{HW} = g_{HB} = g_{Hh} = 0$$

No mixing, only mass terms

$$m_H^2 \Phi_2^\dagger \Phi_2 = m_H^2 \left(|H^+|^2 + \frac{1}{2} |\eta|^2 + \frac{1}{2} |H^0|^2 \right)$$

Only diagonal coupling to up-quarks and leptons

$$\mathcal{L}_Y = Y_{st}^u \bar{q}_s \tilde{\Phi}_1 u_t + Y_{st}^d \bar{q}_s \Phi_1 d_t + Y_{pr}^\ell \bar{\ell}_p \Phi_1 e_r + Y_{pr}'^\ell \bar{\ell}_p \Phi_2 e_r + Y_{st}'^u \bar{q}_s \tilde{\Phi}_2 u_t + \text{h.c.}$$

2HDM

Two scalar doublets with hypercharge +1

$$\begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix} = R_\beta \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} \quad \Phi_1 = \begin{pmatrix} G^+ \\ \frac{v + \rho_1 + iG_0}{\sqrt{2}} \end{pmatrix} \quad \Phi_2 = \begin{pmatrix} H^+ \\ \frac{\rho_2 + i\eta}{\sqrt{2}} \end{pmatrix}$$

$$\begin{pmatrix} h \\ H^0 \end{pmatrix} = R_\alpha \begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix} \quad v = \sqrt{v_1^2 + v_2^2}$$

Decoupling limit

No mixing, only mass terms

$$\alpha, \beta \rightarrow 0$$

$$v_2 \rightarrow 0$$

$$g_{HW} = g_{HB} =$$

Tree-level matching to SMEFT

$$\hat{C}_{4,\ell} = \frac{y'_t y'_\ell}{m_H^2}$$


$$\mathcal{L}_Y = Y_{st}^u \bar{q}_s \tilde{\Phi}_1 u_t$$

$$\left(\frac{1}{2} |\eta|^2 + \frac{1}{2} |H^0|^2 \right)$$


$$s \tilde{\Phi}_2 u_t + \text{h.c.}$$

2HDM

(g-2) dominated by two-loop Barr-Zee diagrams (Ilisie: 1502.04199)



$$\varphi^0 = \{h, H^0, \eta\}$$



One-loop: chirally suppressed

Two-loop enhanced by heavy mass

Dominated by leading logs

$$\ln \left(\frac{m_t^2}{m_H^2} \right)$$