

# $H \rightarrow \gamma\gamma @ 3 \text{ TeV}$

Discussion on the paper to be submitted CLICdp-Draft-2021-003

CLICdp AWG, 17 May 2021

Goran Kacarevic,

Ivanka Bozovic Jelisavcic et al.

VINCA Belgrade & Uni. Kragujevac

# Status and Open issues

- PhD analyses started 2015
- Results presented:
  - CLICdp AWG: 21 Feb 2017, 22 Aug 2017, 24 Feb 2020, 27 Jan 2020
  - CLIC WS 2017, 2021
  - ALPS 2019
  - LCWS21
- Paper submitted to the Pub. Com. 25<sup>th</sup> of January 2021
- Draft number CLICdp-Draft-2021-003 assigned
- Resolved issue of systematic uncertainty estimate from the reconstruction of the luminosity spectrum

(P. Roloff commented that our estimate based of fraction of selected signal events at various center-of-mass energies is too conservative – leads to overestimate)

## **Are we double-counting background by including mono-photon processes ( $e\bar{e}\gamma$ , $\nu\bar{\nu}\gamma$ , $q\bar{q}\gamma$ )?**

- A bit of a history...
  - These processes were considered in the same analysis at 1.4 TeV (published in the Higgs paper)
  - WHIZARD V1.95 used in both analyses
  - ...under similar user's requirements

# 1.4 TeV

# samples

# 3 TeV

Process	$\sigma[\text{fb}]^1$	Events in $1.5 \text{ ab}^{-1}$	Events Available <sup>2</sup>	Event Weights
$e^+e^- \rightarrow \nu\bar{\nu}\gamma$	30	44000	50000	0.9
$e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$	17	26000	22000	1.2
$e^+e^- \rightarrow \gamma\gamma$	27	41000	32000	1.3
$e^+e^- \rightarrow e^+e^-\gamma$	290	430000	285000	1.5
$e^+e^- \rightarrow e^+e^-\gamma\gamma$	13	19000	5500	3.5
$e^+e^- \rightarrow q\bar{q}\gamma$	67	100000	92000	1.1
$e^+e^- \rightarrow q\bar{q}\gamma\gamma$	17	25000	16000	1.5

Signal process	$\sigma(\text{fb})$	$N@5\text{ab}^{-1}$	$N_{\text{simulatec}}$
$e^+e^- \rightarrow H\nu\nu, H \rightarrow \gamma\gamma$	0.95	4750	24550
Background processes		$\sigma(\text{fb})$	
$e^+e^- \rightarrow \gamma\gamma$	19	$9.5 \cdot 10^5$	$3 \cdot 10^4$
$e^+e^- \rightarrow e^+e^-\gamma$	797	$4.0 \cdot 10^6$	$3 \cdot 10^6$
$e^+e^- \rightarrow e^+e^-\gamma\gamma$	56	$2.8 \cdot 10^5$	$1.5 \cdot 10^5$
$e^+e^- \rightarrow \nu\bar{\nu}\gamma$	47	$2.4 \cdot 10^5$	$2 \cdot 10^5$
$e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$	49	$2.5 \cdot 10^5$	$1.6 \cdot 10^5$
$e^+e^- \rightarrow q\bar{q}\gamma$	363	$1.9 \cdot 10^6$	$1.2 \cdot 10^6$
$e^+e^- \rightarrow q\bar{q}\gamma\gamma$	59	$3.0 \cdot 10^5$	$3 \cdot 10^5$

Cuts on generated final states\*:

- At least two photons with  $E > 10 \text{ GeV}$ ,  $p_T > 5 \text{ GeV}$  and  $5^\circ < \theta < 175^\circ$
- At least one Higgs candidate with  $110 \text{ GeV} < M(\gamma\gamma) < 140 \text{ GeV}$
- No visible lepton or quark with  $10^\circ < \theta < 170^\circ$

\*Second photon comes from ISR or FSR (background)

Cuts on generated final states\*:

1. di-photon invariant mass in the window (100-150) GeV
2. at least two photons in event with polar angle between 5 deg and 175 deg
3.  $p_T > 10 \text{ GeV}$  of the photons
4. We haven't restricted  $q$  or  $l$  in the tracker – can be refined

Available samples at [MonteCarloSamplesForTheHiggsPaper](#)

Type	Energy	Detector	ProdID	Events planned	Events produced	$\sigma$ [fb]	Comments
ee -> qq $\gamma\gamma$	3 TeV	CLIC_ILD	6966	250000	299600	72.3	WHIZARD V57, $m(h) = 12 \text{ TeV}$
ee -> qq $\gamma$	3 TeV	CLIC_ILD	6969	1000000	1198600	583.6	WHIZARD V57, $m(h) = 12 \text{ TeV}$
ee -> $\gamma\nu\nu$	3 TeV	CLIC_ILD	6981	150000	194900	16806.4*	WHIZARD V57, $m(h) = 12 \text{ TeV}$ , preselection cuts for the H->gammagamma analysis ("cuts_h_gammagamma_3000.txt")
ee -> $\gamma\gamma ee$	3 TeV	CLIC_ILD	6984	120000	144900	4244.7*	WHIZARD V57, $m(h) = 12 \text{ TeV}$ , preselection cuts for the H->gammagamma analysis ("cuts_h_gammagamma_3000.txt")
ee -> $\gamma\nu\nu$	3 TeV	CLIC_ILD	7014	120000	155500	2616.6*	WHIZARD V57, $m(h) = 12 \text{ TeV}$ , preselection cuts for the H->gammagamma analysis ("cuts_h_gammagamma_3000.txt")
ee -> $\gamma\gamma$	3 TeV	CLIC_ILD	7017	20000	29900	866.9*	WHIZARD V57, $m(h) = 12 \text{ TeV}$ , preselection cuts for the H->gammagamma analysis ("cuts_h_gammagamma_3000.txt")
ee -> ee $\gamma$	3 TeV	CLIC_ILD	7020	3000000	3039000	185392.3*	WHIZARD V57, $m(h) = 12 \text{ TeV}$ , preselection cuts for the H->gammagamma analysis ("cuts_h_gammagamma_3000.txt")

# Discussion

## Are we double-counting background by including mono-photon processes ( $e^+e^- \gamma$ , $\nu\nu\gamma$ , $qq\gamma$ )?

• There are several questions that can be asked:

- 1) Are cross-sections for processes with a single photon in the final state correctly reproduced in WHIZARD V1.95?
- 2) Is kinematics of the final state photon properly described?

YES

If yes, than it is not justified to omit these processes because, apparently second photon can be found at the reconstruction level in a sufficient amount to mimic the signal

$N_{BDT}$ , for integrated luminosity of  $5 \text{ ab}^{-1}$

Process	$\epsilon_{pre}(\%)$	$\epsilon_{BDT}(\%)$	$N_{BDT}$
$e^+e^- \rightarrow H\nu\nu, H \rightarrow \gamma\gamma$	70	62.0	2062
$e^+e^- \rightarrow \gamma\gamma^*$	0.10	13.7	624
$e^+e^- \rightarrow e^+e^-\gamma^*$	0.03	9.8	26598
$e^+e^- \rightarrow e^+e^-\gamma\gamma^*$	0.05	7.4	788
$e^+e^- \rightarrow \nu\nu\gamma^*$	0.09	25.0	18585
$e^+e^- \rightarrow \nu\nu\gamma\gamma^*$	0.04	24.5	11628
$e^+e^- \rightarrow qq\gamma$	0.50	7.5	1089
$e^+e^- \rightarrow qq\gamma\gamma$	1	10.1	423

NO

If not, than it depends what is wrong:

- a) Only cross-section  $\rightarrow$  can be cured with scaling (in MVA)
- b) Kinematics  $\rightarrow$  processes in question should be reprocessed in WHIZARD (2 ?)

*Without clear answers to these questions can we be convinced that it is justified to neglect mono-photon processes?*

```

ISR

flag: 0x0
simulator status bits: [sbvtcls] s: created in simulation b: backscatter v: vertex is not endpoint

[ id ]index| PDG | px, py, pz | px_ep, py_ep, pz_ep | energy |gen|
| | spin | | colorflow | [parents] - [daughters]
[00000128] 0| 11| 0.00e+00, 0.00e+00, 1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [2,4]
[00000129] 1| -11| 0.00e+00, 0.00e+00, -1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [3,5]
[00000130] 2| 11| 2.75e-05, -5.57e-05, 1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0] - [6,7]
[00000131] 3| -11| 3.94e+00, 9.62e-01, -7.72e+02| 0.00e+00, 0.00e+00, 0.00e+00| 7.72e+02| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [1] - [6,7]
[00000132] 4| 22|-2.75e-05, 5.57e-05, 2.30e-04| 0.00e+00, 0.00e+00, 0.00e+00| 2.38e-04| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0] - []
[00000133] 5| 22|-3.94e+00, -9.62e-01, -7.28e+02| 0.00e+00, 0.00e+00, 0.00e+00| 7.28e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [1] - []
[00000134] 6| 13|-5.96e+02, -7.44e+02, 8.87e+02| 0.00e+00, 0.00e+00, 0.00e+00| 1.30e+03| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
[00000135] 7| -13| 6.00e+02, 7.45e+02, -1.59e+02| 0.00e+00, 0.00e+00, 0.00e+00| 9.70e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
-----

```

```

BEAMSTRAHLUNG

[ id ]index| PDG | px, py, pz | px_ep, py_ep, pz_ep | energy |gen|
| | spin | | colorflow | [parents] - [daughters]
[00000030] 0| 11| 0.00e+00, 0.00e+00, 1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [2,3]
[00000031] 1| -11| 0.00e+00, 0.00e+00, -1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [2,3]
[00000032] 2| 11| 0.00e+00, 0.00e+00, 1.11e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.11e+03| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - [4,5]
[00000033] 3| -11| 0.00e+00, 0.00e+00, -9.24e+01| 0.00e+00, 0.00e+00, 0.00e+00| 9.24e+01| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - [4,5]
[00000034] 4| 13| 2.25e+01, 3.03e+02, 3.19e+02| 0.00e+00, 0.00e+00, 0.00e+00| 4.41e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
[00000035] 5| -13|-2.25e+01, -3.03e+02, 6.96e+02| 0.00e+00, 0.00e+00, 0.00e+00| 7.59e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
-----

```

```

1 ISR + BS
2
3
4 [ id ]index| PDG | px, py, pz | px_ep, py_ep, pz_ep | energy |gen|
| | spin | | colorflow | [parents] - [daughters]
5
6 [00000020] 0| 11| 0.00e+00, 0.00e+00, 1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [2,3,4,5]
7 [00000021] 1| -11| 0.00e+00, 0.00e+00, -1.50e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.50e+03| 4 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [] - [2,3,4,5]
8 [00000022] 2| 11| 6.22e-15, -7.11e-15, 1.45e+03| 0.00e+00, 0.00e+00, 0.00e+00| 1.45e+03| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - [6,7]
9 [00000023] 3| -11| 1.23e+01, -1.79e+01, -4.77e+02| 0.00e+00, 0.00e+00, 0.00e+00| 4.78e+02| 3 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - [6,7]
0 [00000024] 4| 22| 0.00e+00, 0.00e+00, 2.20e-26| 0.00e+00, 0.00e+00, 0.00e+00| 2.20e-26| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - []
1 [00000025] 5| 22|-1.23e+01, 1.79e+01, -2.59e+02| 0.00e+00, 0.00e+00, 0.00e+00| 2.60e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [0,1] - []
2 [00000026] 6| 13|-6.11e+02, -4.60e+02, 8.62e+02| 0.00e+00, 0.00e+00, 0.00e+00| 1.15e+03| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
3 [00000027] 7| -13| 6.23e+02, 4.42e+02, 1.07e+02| 0.00e+00, 0.00e+00, 0.00e+00| 7.72e+02| 1 |
+00, 0.00e+00, 0.00e+00| (0, 0) | [2,3] - []
4

```

### Whizard 2.8.3: $ee \rightarrow \mu\mu$

- There are no BS photons when BS is included in production
- Beamstrahlung photons are not visible in MCParticleSKimmed collection, and we cannot know if the reconstructed photon is BS photon. But we can check it is ISR photon


# Background samples

Process	$N_{\text{preselection}}$	$N_{\text{ISR}}$	event loss(%)
$ee \rightarrow \gamma\gamma$	17578	3505	19.9
$ee \rightarrow ee\gamma$	16449	9528	57.9
$ee \rightarrow ee\gamma\gamma$	34717	14452	41.6
$ee \rightarrow qq\gamma$	1287	543	42.1
$ee \rightarrow qq\gamma\gamma$	13494	2801	20.8
$ee \rightarrow vv\gamma$	54914	44961	81.9
$ee \rightarrow vv\gamma\gamma$	54554	25830	47.3


If at least one candidate photon comes from ISR (RecoMCTruthLink) all backgrounds have significant loss.

# Di-photon mass distribution after MVA


**OLD - all background**



**no mono-photon**



**All backgrounds rescaled**




	$N_s$	$N_b$	Significance/ $\delta$ (%)
All backgrounds	2060	63000	8.3/12
No mono photon	2360	17000	17/5.9
All backgrounds rescaled	1926	27000	11.32/8.9




# Signal Fit


OLD



no mono-photon




All backgrounds rescaled



No relevant changes in mean and width



# Pull Distribution



The width of the pull distribution is reduced, resulting in a drop of statistical uncertainty from 8.3% , 4.3% and 6%.

# Summary

- After 5000 Toy-MC experiments uncertainty is 8.3%, 6 % and 4.3 % when including all backgrounds, rescaling all backgrounds, removing mono-photon backgrounds respectively.