

# The soft drop momentum sharing fraction $z_g$ beyond leading-logarithmic accuracy

*Tuesday, August 3, 2021 4:30 PM (15 minutes)*

Based on: <https://arxiv.org/abs/2106.04589>

Abstract:

Grooming techniques, such as soft drop, play a central role in reducing sensitivity of jets to e.g. underlying event and hadronization at current collider experiments. The momentum sharing fraction  $z_g$ , of the two branches in a jet that pass the soft drop condition, is one of the most important observables characterizing a collinear splitting inside the jet, and directly probes the QCD splitting functions. In this work, we present a factorization framework that enables a systematic calculation of the corresponding cross section beyond leading-logarithmic (LL) accuracy, showing that this measurement is not only sensitive to the QCD charge but also the spin of the parton that initiates the jet. Our results at next-to-leading logarithmic (NLL') accuracy include non-global logarithms, and provide a first meaningful assessment of the perturbative uncertainty. We present a comparison to the available experimental data from ALICE, ATLAS, and STAR and find excellent agreement.

**Authors:** CAL, Pedro (University of Amsterdam); WAALEWIJN, Wouter (University of Amsterdam); RINGER, Felix (Lawrence Berkeley National Laboratory); LEE, Kyle

**Presenter:** CAL, Pedro (University of Amsterdam)

**Session Classification:** Precision Calculations + Experimental Heavy-Ions Results