

# Collective flow of light nuclei and hyper-nuclei in Au+Au collisions at $\sqrt{s_{NN}} = 3, 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV}$ using the STAR detector

**Rishabh Sharma (*for the STAR Collaboration*)**  
**Indian Institute of Science Education and Research (IISER) Tirupati, India**



Supported in part by:



Office of  
Science


STAR Presentations: <https://drupal.star.bnl.gov/STAR/presentations>




# Outline



- ★ Motivation
- ★ The STAR experiment
- ★ Analysis method
- ★ Results
  - Elliptic flow of light nuclei
  - Directed flow of light (hyper-)nuclei
- ★ Summary




Nature 561, 321–330 (2018)



STAR, PRC 94, 034908 (2016)

- ★ Light (hyper-)nuclei production in heavy-ion collisions can be explained either by the **thermal model** or the **final-state coalescence** of nucleons
- ★  $v_2/A$  of light nuclei was observed to be close to  $v_2$  of protons for  $p_T/A < 1.5$  GeV/c in BES-I data
- ★ Higher statistics dataset in BES-II program will allow us to revisit and better understand the production mechanism of light (hyper-)nuclei

- ★ **Solenoidal Tracker at RHIC (STAR)** is one of the large detector systems at RHIC consisting of several sub-detectors
- ★  $dE/dx$  information from **Time Projection Chamber (TPC)** and  $m^2$  information from **Time of Flight (TOF)** are used for particle identification
- ★ Upgrade to **iTPC**
  - Large acceptance ( $|\eta| < 1.5$ )
  - Better track resolution
- ★ **Event Plane Detector (EPD)**:  $2.1 < |\eta| < 5.1$
- ★ Datasets:
  - **BES II**: Au+Au collisions at  $\sqrt{s_{NN}} = 3$  (FXT), 14.6, 19.6, 27, and 54.4 GeV (COL)



JINST 15 C07040 (2020)

- ★ The particle azimuthal distribution can be written as:

$$E \frac{d^3N}{d^3p} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left( 1 + \sum_{n=1}^{\infty} 2v_n \cos(n(\phi - \Psi_R)) \right) \quad v_n = \langle \cos[n(\phi - \psi_R)] \rangle$$

$v_1$ : Directed flow

$v_2$ : Elliptic flow


- ★  $n^{\text{th}}$  harmonic plane is calculated using the Q-vector:

$$Q_n \cos(n\Psi_n) = \sum_i w_i \cos(n\phi_i)$$

$$Q_n \sin(n\Psi_n) = \sum_i w_i \sin(n\phi_i)$$


$$\Psi_n = \left( \tan^{-1} \frac{\sum_i w_i \sin(n\phi_i)}{\sum_i w_i \cos(n\phi_i)} \right) / n$$

- ★  $\eta$ -sub event plane method is used




CMS, PRC 87 014902 (2013)

Rishabh Sharma - SQM 2022



STAR, PRL 103, 251601 (2009)




We observe an improvement of resolution by ~10% from BES I owing to higher TPC acceptance and track resolution


- ★ Particles are identified using **dE/dx information from TPC** in the range  $|\eta| \leq 1.0$

$$z_i = \ln \left( \frac{\langle dE/dx \rangle_{measured}}{\langle dE/dx \rangle_{theory}} \right)$$


- ★  $\langle dE/dx \rangle_{theory}$  is calculated using Bichsel function
- ★ Double Gaussian fit is done to calculate yield in each  $p_T$  and  $\phi - \Psi_2$  bin




Au+Au  $\sqrt{s_{NN}} = 19.6$  GeV COL (2019), d




Au+Au  $\sqrt{s_{NN}} = 19.6$  GeV COL (2019), t



Au+Au  $\sqrt{s_{NN}} = 19.6$  GeV COL (2019),  $^3\text{He}$



$\sqrt{s_{NN}} = 3 \text{ GeV (FXT) Au+Au Collisions at RHIC (BES II)}$




○
**KFParticle package has been used for signal reconstruction**

*I. Kisel (CBM), J. Phys. Conf. Ser. 1070, 012015 (2018)*

Rishabh Sharma - SQM 2022

# Elliptic flow ( $v_2$ )






The  $v_2(p_T)$  for all nuclei species increases with increasing  $p_T$  for all collision energies

Statistical errors have reduced significantly compared to the BES I results




# Centrality dependence



$v_2$  of deuterons shows a strong centrality dependence


Peripheral collisions have relatively larger  $v_2$  due to their larger initial spatial anisotropy



A systematic deviation of around 20-30% from mass number scaling is observed for all light nuclei species at all measured energies

# Directed flow ( $v_1$ ) vs rapidity

$\sqrt{s_{NN}} = 3 \text{ GeV (FXT) Au+Au Collisions at RHIC (BES II)}$




○ First observation of hyper-nuclei directed flow ( $v_1$ ) in high-energy heavy-ion collisions

○ Hyper-nuclei  $v_1$  seems to follow the mass number scaling

STAR, PLB 827 136941 (2022)

# v<sub>1</sub>-slope vs particle mass



**○ Within statistical uncertainties, the slopes of  $v_1$  of hypernuclei seem to follow the mass number scaling**

STAR, PLB 827 136941 (2022)

- ★  $v_2$  of d, t, and  $^3\text{He}$  is measured in Au+Au collisions at  $\sqrt{s_{\text{NN}}} = 14.6, 19.6, 27$  and,  $54.4$  GeV (COL)
  - 20-30% deviation of light nuclei  $v_2$  from mass number scaling is observed
  - Clear centrality dependence is observed for deuterons for all collision energies
- ★  $v_1$  of  $\Lambda$ ,  $^3_{\Lambda}H$ , and  $^4_{\Lambda}H$  is presented in Au+Au collisions at  $\sqrt{s_{\text{NN}}} = 3$  GeV (FXT)
  - Rapidity dependence of hyper-nuclei  $v_1$  is measured
  - $v_1$  of hyper-nuclei shows mass number scaling

## Outlook

- ★ Stay tuned for more exciting results on (hyper-)nuclei from BES II energies

**Thank you**