

Thermal production of Sexaquark in relativistic heavy Ion Collisions

Tuesday 14 June 2022 17:13 (1 minute)

We consider the probability of thermal production of $uuddss$ states with small radii $r = 0.1 - 0.4$ fm and small masses $m < 2$ GeV in Pb+Pb collisions at $\sqrt{s}=2.76$ TeV. We use thermal model which was tuned to AGS, RHIC, LHC data and predicts $T = 150 - 170$ MeV.

Hereby we compare the production rates for Sexaquarks with different masses and different radii.

We found that sexaquarks are produced with relatively high rates for both 0.1 and 0.4 fm radii and for masses of 1700 and 1960 MeV.

We estimate ratios of Sexaquarks $uuddss$ to hadrons (protons, kaons, Lambda) and deuterons in Pb+Pb collisions at LHC.

At $T=170$ MeV the ratio of thermal Sexaquark with mass 1960 MeV to thermal deuteron is about 0.45.

Present via

Primary authors: TARANENKO, Arkadiy (NRNU MEPhI); BLASCHKE, David; BRAVINA, Larisa; Dr GRINYUK, Boris (Bogolyubov Institute for Theoretical Physics); Dr ZHOU, Daicui (CCNU); ZABRODIN, Evgeny; FARRAR, Glennys (NYU); VITIUK, Oleksandr (University of Wroclaw, Institute of Theoretical Physics); IVANYTSKYI, Oleksii (University of Wroclaw); KULESHOV, Serguei (Universidad Andrés Bello (Santiago, Chile)/SAPHIR Millennium Institute of ANID, Chile); KABANA, Sonia (Instituto De Alta Investigación - Universidad de Tarapacá (CL)); SAGUN, Violetta (University of Coimbra); Dr ZHANG, xiaoming (CCNU)

Presenter: BRAVINA, Larisa

Session Classification: Poster

Track Classification: Bulk matter phenomena, QCD phase diagram, and Critical point