

The logo for SHINE features the word "SHINE" in a bold, blue, sans-serif font. The letter "H" is stylized, composed of several overlapping, slanted rectangular bars in various shades of blue, creating a sense of depth and movement. The two dots of the "SH" are solid blue circles.


**SHINE**

# Highlights from the NA61/SHINE

Szymon Pulawski  
for NA61/SHINE

# Upgraded NA61/SHINE detector

Fixed target experiment located at the CERN SPS accelerator





Readout rate increased to 1 kHz  
New Vertex Detector  
New DAQ and trigger system  
Upgraded PSD

**Large acceptance hadron spectrometer** –

coverage of the full forward hemisphere, down to  $p_T = 0$

# NA61/SHINE 2-dimensional scan

NA61/SHINE performed a 2D scan in **collision energy and system size** to study the phase diagram of strongly interacting matter







# Onset of deconfinement

Qualitatively similar energy dependence is seen in p+p, Be+Be, Ar+Sc and Pb+Pb



Magnitude of T increases with the system size



Kaons are only weakly affected by rescattering and resonance decays during the post-hydro phase (at SPS and RHIC energies).

Connected temperature of the freeze-out surface and not the early-stage fireball


## Plateau like structure visible in p+p, Be+Be and Ar+Sc Ar+Sc is higher than p+p and Be+Be



Good measure of the strangeness to entropy ratio which is different in the confined phase (hadrons) and the QGP (quarks, anti-quarks and gluons).

Probe of the onset of deconfinement.

## The enhancement recalculated based on the new $\Xi$ reference from NA61/SHINE



The strangeness enhancement factor:

$$E = \frac{2}{\langle N_W \rangle} \frac{dn/dy (A + A)}{dn/dy (p + p)}$$


Nucl. Phys. B111 (1976) 461

J. Phys. G 32 (2006) 427–442



## System size dependence






None of the models reproduces  $K^+/\pi^+$  ratio or  $T$  for whole  $\langle W \rangle$  range

PHSD: Eur.Phys.J.A 56 (2020) 9, 223, arXiv:1908.00451 and private communication;  
 SMASH: J.Phys.G 47 (2020) 6, 065101 and private communication;  
 UrQMD and HRG: Phys. Rev. C99 (2019) 3, 034909  
 SMES: Acta Phys. Polon. B46 (2015) 10, 1991 - recalculated

p+p: Eur. Phys. J. C77 (2017) 10, 671  
 Be+Be: Eur. Phys. J. C81 (2021) 1, 73  
 Ar+Sc: NA61/SHINE preliminary  
 Pb+Pb: Phys. Rev. C66, 054902 (2002)





## Search for critical point

Expected: non-monotonic behavior of CP signatures



# Multiplicity and net-charge fluctuations in p+p, Be+Be and Ar+Sc

## No structure indicating critical point



$$\begin{aligned} \kappa_1 &= \langle N \rangle \\ \kappa_2 &= \langle (\delta N)^2 \rangle = \sigma^2 \\ \kappa_3 &= \langle (\delta N)^3 \rangle = S\sigma^3 \\ \kappa_4 &= \langle (\delta N)^4 \rangle - 3\langle (\delta N)^2 \rangle^2 = K\sigma^4 \end{aligned}$$

where:

$N$  – multiplicity;  $\delta N = N - \langle N \rangle$   
 $\sigma$  – standard deviation  
 $S$  – skewness;  $K$  – kurtosis






Negatively charge  $\kappa_2/\kappa_1$ : increasing difference between small systems (p+p and Be+Be) and a heavier system (Ar+Sc) with collision energy

Net-charge  $\kappa_3/\kappa_1$ : increasing difference between Be+Be and other systems (p+p and Ar+Sc) with collision energy

$\kappa_4/\kappa_1$  : consistent values for all measured systems at given collision energy

# Proton and charge hadron intermittency in Ar+Sc and Pb+Pb collisions

**No structure indicating critical point** see the talk by N. Davis on Wednesday and the poster by T. Czopowicz





$$F_r(M) = \frac{\left\langle \frac{1}{M} \sum_{m=1}^M n_m (n_m - 1) \dots (n_m - r + 1) \right\rangle}{\left\langle \frac{1}{M} \sum_{m=1}^M n_m \right\rangle^r},$$

where  $\langle \dots \rangle$  denotes averaging over events,  $M$  the number of cells

Statistically independent points, cumulative variables  
No indication of critical point in these analyses  
(power-law scaling  $F_r(M) \sim M^{\phi_r}$ )

# Symmetric Levy HBT correlations




The Levy stability parameter  $\alpha$  describes shape of the source

3D Ising model with random external field predicts  $\alpha = 0.5 \pm 0.05$  at critical point




## Highlights from strangeness production in $p+p$

# $K^\pm$ production in inelastic $p+p$ collisions at 158 GeV/c



see the talk by P. Podlaski on Tuesday  
see the poster by A. Shukla




## New results on $K^+$ , $K^-$ (preliminary) and $K_S^0$ from high statistic $p+p$ data

$K^\pm$ : almost 20 times larger dataset than previously published results (Eur.Phys.J.C 77 (2017), 671)

$K_S^0$  mean multiplicity:  $0.162 \pm 0.001 \pm 0.011$

Model predictions deviate by up to 20% from the measurements — best predictions from EPOS 1.99.



$K^*/K^-$  or  $K^*/K^+$  → time between chemical and kinetic freeze-outs, properties of hadron gas phase

STAR, PR C71, 064902, 2005;

C. Blume, APP B43, 577, 2012


$$\left. \frac{K^*}{K} \right|_{kinetic} = \left. \frac{K^*}{K} \right|_{chemical} e^{-\frac{\Delta t}{\tau}}$$

A+A

p+p

## Mean multiplicity of $K^*(892)^0$

$\sqrt{s_{NN}}$	NA61	NA49 (PR C84, 064909, 2011)
8.8	$(35.1 \pm 1.3 \pm 3.6) \cdot 10^{-3}$	-
12.3	$(58.3 \pm 1.9 \pm 4.9) \cdot 10^{-3}$	-
17.3	$(78.44 \pm 0.38 \pm 6.0) \cdot 10^{-3}$	$(74.1 \pm 1.5 \pm 6.7) \cdot 10^{-3}$




GCE: good fit (unexpectedly!)

CE: good fit only with  $\phi$  meson excluded



# [I] production in inelastic $p+p$ collisions at 158 GeV/c

see the talk by P. Podlaski on Tuesday








The only results on  $\Xi^-$  and  $\Xi^+$  production in  $p+p$  at SPS energy

Strong suppression of  $\Xi^+$  production:  $\langle \Xi^+ \rangle / \langle \Xi^- \rangle = 0.24 \pm 0.01 \pm 0.05$

# [Ξ] production in inelastic p+p collisions – model comparison




see the talk by P. Podlaski on Tuesday

Transport models **fail** to describe the NA61/SHINE results on  $\Xi$  production in p+p collisions



# $\Xi(1530)^0$ production in inelastic $p+p$ collisions at 158 GeV/c

see the talk by P. Podlaski on Tuesday



**The only results on  $\Xi(1530)^0$  production in  $p+p$  at the SPS energy**

The second result on  $\Xi(1530)^0$  production in  $p+p$  (ALICE at 7 TeV Eur.Phys.J.C 75 (2015) 1)

Suppression of  $\bar{\Xi}(1530)^0$  production:  $\langle \bar{\Xi}(1530)^0 \rangle / \langle \Xi(1530)^0 \rangle = 0.40 \pm 0.03 \pm 0.05$





# $\Xi(1530)^0$ production in inelastic $p+p$ collisions at 158 GeV/c

see the talk by P. Podlaski on Tuesday



Eur.Phys.J.C 81 (2021) 10, 911

EPOS describes well transverse momentum and rapidity distributions of  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$

UrQMD significantly overestimates all spectra of  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$  hyperons



# HRG model in the CE formulation and $p+p$ data



Eur.Phys.J.C 81 (2021) 10, 911

Fit by different variants of the HRG model (THERMAL-FIST1.3 Comput.Phys.Commun.244 (2019)295):

- Canonical Ensemble with fixed  $\gamma_s=1$
- Canonical Ensemble with fitted strangeness saturation parameter  $\gamma_s$

Significant discrepancies of the fitted parameters

The statistical model fails when fixed  $\gamma_s$

**The fit with free  $\gamma_s$  finds  $\gamma_s = 0.434 \pm 0.028$  and reproduces the measurements well - a suppression of strange particle production in  $p+p$  collisions at CERN SPS energies**



## NA61/SHINE in 2022-2025

## Upgrade almost completed First Pb+Pb data taking in autumn 2022


What is the mechanism of open charm production?

How does the onset of deconfinement  
impact open charm production?


How does the formation of quark gluon plasma  
impact  $J/\psi$  production?

To answer these questions the mean number of charm quark pairs,  $\langle c\bar{c} \rangle$ , produced in A+A collisions has to be known. Up to now the corresponding experimental data does not exist and NA61/SHINE will perform this measurement in the near future.


$\langle c\bar{c} \rangle$  and models



$\langle c\bar{c} \rangle$  and onset of deconfinement



$\langle c\bar{c} \rangle$ ,  $\langle J/\psi \rangle$  and QGP



# Summary

- 2D scan in system size and collision energy was completed in 2017 with Xe+La data
- Analysis ongoing for p+p, Be+Be, Ar+Sc, Xe+La and Pb+Pb data
- No horn in Ar+Sc collisions
- Unexpected system size dependence:  $(p+p \approx \text{Be+Be}) \neq (\text{Ar+Sc} \neq \text{Pb+Pb})$
- No convincing indication of CP
- New and unique results on  $K^+$ ,  $K^-$ ,  $K_S^0$ ,  $K^*$ ,  $\Xi^-$ ,  $\bar{\Xi}^+$ ,  $\Xi(1530)^0$  and  $\bar{\Xi}(1530)^0$  production in p+p interactions
- NA61/SHINE program with measurements of open charm production in 2022-2025




# NA61/SHINE at SQM conference

- P. Podlaski, Tuesday 15:20, „Results on system size dependence of strangeness production in the CERN SPS energy range from NA61/SHINE”
- N. Davis , Wednesday 9:40, „Assessing critical point signatures through proton intermittency in NA61/SHINE”
- Posters:
  - T. Czopowicz, „Search for critical point in NA61/SHINE (POS-BLK-11)”
  - A. Tefelska, „Mesonic strange resonances in p+p collisions at SPS energies (POS-RES-04)”
  - A. Shukla, „Identified hadron spectra in high-statistics p+p collisions at 158 GeV/c (POS-OTH-02)”




Thank you




# Strangeness production in p+p at 158 GeV/c



Eur.Phys.J.C 80 (2020) 3, 199



Eur.Phys.J.C 80 (2020) 9, 833






**Present theoretical models do not describe the NA61/SHINE results on strange particles production in p+p interactions**

# Strangeness enhancement factors

J. Phys. G 32 (2006) 427–442


Eur.Phys.J.C 80 (2020) 9, 833



$$E_{\Xi_s} = \frac{2}{\langle N_w \rangle} \frac{dn/dy(A+A)}{dn/dy(p+p)}$$

**NA61/SHINE results give new base-line for strangeness enhancement study in SPS energy range**

## Fixed target experiment located at the CERN SPS accelerator



Beams:



- ions (Be, Ar, Xe, Pb)  
 $p_{\text{beam}} = 13A - 150A \text{ GeV}/c$
- hadrons ( $\pi$ , K, p)  
 $p_{\text{beam}} = 13 - 400 \text{ GeV}/c$
- $\sqrt{s_{NN}} = 5.1 - 16.8 (27.4) \text{ GeV}$

**Large acceptance hadron spectrometer** –  
 coverage of the full forward hemisphere, down to  $p_T = 0$

# Diagram of high-energy nuclear collisions

## Hypothetical domains of hadron-production dominated by:

- resonance creation and decays
- string creation and decays
- quark-gluon plasma formation and hadronisation







theo: Phys.Part.Nucl. 51 (2020) 3, 337-339  
exp: Phys.Rev.C 102 (2020) 1, 011901



## Transition from resonances to strings

# Transition from resonances to strings



Rates of increase of  $K^+/\pi^+$  and  $T$  change sharply in p+p collisions at SPS energies


The fitted change energy is  $\approx 7$  GeV - close to the energy of the onset of deconfinement  $\approx 8$  GeV

Models assuming change from resonances to string production mechanism show similar trend



# Exclusion plots for parameters of simple power-law model

using statistically independent points and cumulative variables



The predicted intermittency index for a system freezing out at the QCD critical endpoint corresponds to the 3-D Ising universality class, to which the phase transition is expected




## NA61/SHINE in 2021-2024

# NA61/SHINE program for 2021-2024


- What is the mechanism of open charm production?
- How does the onset of deconfinement impact open charm production?
- How does the formation of quark gluon plasma impact  $J/\psi$  production?

To answer these questions **mean number of charm quark pairs,  $\langle c\bar{c} \rangle$** , produced in A+A collisions has to be known. Up to now corresponding experimental **data does not exist** and **only NA61/SHINE can perform this measurement in the near future.**


### $\langle c\bar{c} \rangle$ and models



### $\langle c\bar{c} \rangle$ and onset of deconfinement



### $\langle c\bar{c} \rangle$ , $\langle J/\psi \rangle$ and QGP



**Foreseen NA61/SHINE resolution is sufficient to answer addressed questions**

# Detector upgrade during LS2

Construction of Vertex Detector (VD)  
for  $D^0$ ,  $\bar{D}^0$  decay reconstruction

Replacement of the TPC  
read-out electronics  
to increase data rate to 1 kHz

