

**SQM2022**

The 20th International Conference on Strangeness in Quark Matter  
13-17 June 2022 Busan, Republic of Korea



# Strange Hadron Production in Au+Au Collisions at RHIC Beam Energy Scan


Yingjie Zhou, for the STAR Collaboration  
Central China Normal University

Supported in part by



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science




# Outline

---

- **Introduction**
- **STAR Fixed Target (FXT)**
- **Results of strange hadron production**
  - $p_T$  spectra
  - Rapidity distribution
  - Yield ratio
  - Kinetic freeze-out properties
- **Summary**


# Introduction





- RHIC BES covers a wide region of baryon density
  - Look for the onset of de-confinement, phase boundary, and location of possible critical point
- STAR FXT mode  $\sqrt{s_{NN}} = (3.0 - 7.7) \text{ GeV}$ 
  - High baryon chemical potential  $\mu_B$  ( $\sim 400 \text{ MeV}$  up to  $\sim 750 \text{ MeV}$ ) allows us to study the properties of high baryon density matter
  - Strangeness can be used to study medium properties at low collision energies

# FXT setup at STAR

- Target was installed at the edge of TPC
- 260M events for Au+Au FXT at  $\sqrt{s_{NN}} = 3 \text{ GeV}$  (year 2018)
- Good mid-rapidity coverage



$V_y$  vs.  $V_x$  Distribution



**Beam pipe**




**Au-Target = 0.25mm thickness  
1% interaction probability**


Conventions:  
beam-going direction is the positive direction  
In C.M. frame,  $y_{target} = -1.045$  for the 3 GeV collisions

# Particle identification and reconstruction


TPC




TOF



- TPC ( $dE/dx$ ) and TOF ( $\beta$ ) for pion, kaon and proton identification
- Reconstruct the short-lived particle  $\Xi^-$ ,  $\phi$ ,  $\Lambda$ ,  $K_S^0$  via a hadronic decay channel
  - $\phi \rightarrow K^+K^-$ ,  $\Xi^- \rightarrow \Lambda(p\pi^-) + \pi^-$ ,  
 $K_S^0 \rightarrow \pi^+\pi^-$
  - KF Particle package is used to improve the signal significance
- The combinatorial background is reconstructed by the rotation method



# Efficiency corrected $p_T$ spectra




- Tracking efficiency and detector acceptance are estimated with GEANT simulations embedded into real events
- $\Lambda$  and  $K_S^0$  invariant yields in 0-10% centrality for various rapidity regions
- **Low  $p_T$  extrapolation: Blast-Wave function**

$$\frac{d^2N}{2\pi p_T dp_T dy} = A \int_0^R r dr m_T \times I_0\left(\frac{p_T \sinh \rho(r)}{T_{\text{kin}}}\right) K_1\left(\frac{m_T p \cosh \rho(r)}{T_{\text{kin}}}\right)$$

$T_{\text{kin}}$ : the kinetic freeze-out temperature  
 $\langle \beta_T \rangle$ : average transverse radial flow velocity  
 $n$ : the exponent of flow velocity profile,  $n=1$   
 $I_0$  and  $K_1$  are from Bjorken Hydrodynamic assumption
- Alternative fit functions are used in order to estimate the systematic uncertainty in  $dN/dy$


# Centrality and rapidity dependence of yields



Phys. Lett. B 831 (2022) 137152

- Rapidity dependent yields obtained from integrating data and Blast-wave function (or  $m_T$  exponential function) fits of spectra for the unmeasured region
  - $y_{cm}$  range will be extended by eTOF & iTPC upgrade
- UrQMD reproduces the yields of  $\Lambda$  except in the 40-60% centrality bin, but overestimates Kaons,  $\Xi^-$  and underestimates  $\phi$  mesons

# Centrality and rapidity dependence of particle ratio




UrQMD: 0-10% and 10-40% centrality are scaled up by 25%

- Enhanced production of  $\Lambda$  at mid-rapidity compared to target rapidity
- UrQMD model underpredicts those ratios but can describe the trend of the data
- Comparison with hypernuclei to light nuclei ratios help us gain insight into hypernuclei and light nuclei production mechanisms

See talk from Yuanjing Ji, Jun 14, 2022, 11:30 AM




# Strangeness production vs $\langle N_{Part} \rangle$



- **Universal centrality dependence of strangeness production**
  - Increase with centrality
  - Strangeness yield ( $K^-, K_S^0, \phi, \Lambda$ )  $\propto \langle N_{part} \rangle^\alpha$ ,  $\alpha = 1.42 \pm 0.04$
- **$\Xi^-$  seems to deviate from the scaling trend**
  - $\Xi^-$  is different from other hadrons due to its multi-strange-quark content and sub-threshold production
- **Proton has a different trend**

$$\begin{aligned}
 K_S^0 \sqrt{s_{thr}} &\sim 2.55 \text{ GeV} \\
 \Lambda \sqrt{s_{thr}} &\sim 2.55 \text{ GeV} \\
 \phi \sqrt{s_{thr}} &\sim 2.89 \text{ GeV} \\
 \Xi^- \sqrt{s_{thr}} &\sim 3.25 \text{ GeV}
 \end{aligned}$$


# Strangeness production vs $\sqrt{s_{NN}}$



- STAR FXT  $dN/dy$  consistent with the  $\Lambda$ ,  $K_S^0$  trends demonstrated by published data
- First sub-threshold  $E^-$  measurement in Au+Au collisions
- Expect more results at low energies from additional high statistics BES II data sets

Data compilation: C. Blume Prog.Part.Nucl.Phys. 66 (2011) 834-879  
 HADES: Phys.Lett.B 793 (2019) 457-463, 2019

# Particle ratios vs $\sqrt{s_{NN}}$




- Local strangeness conservation is required!  
➔ **GCE to CE transition!**
- Default UrQMD failed to describe the measurement at low energies
- Transport models with high-mass resonance decay to  $\phi$  and  $E^-$  (modified UrQMD and SMASH) can reasonably describe data at low energies

UrQMD<sup>1</sup>: the public version  
UrQMD<sup>2</sup>: the modified version

$r_c$ : correlation length, radius of the volume inside which the production of particles with open strangeness is canonically conserved

Data compilation:  
STAR: Phys. Lett. B 831 (2022) 137152, Phys. Rev. C 102 (2020) 34909  
HADES: Eur. Phys. J. A (2016) 52: 178  
UrQMD<sup>1</sup>: Prog. Part. Nucl. Phys. 41 (1998) 225-370  
UrQMD<sup>2</sup>: J. Phys. G: Nucl. Part. Phys. 43 015104  
Thermal CE: Phys. Lett. B 603, 146 (2004)

# Kinetic freeze-out properties



Phys. Rev. Lett. 108 (2012) 72301  
Phys. Rev. C 102 (2020) 34909

- Kinetic freeze-out temperature ( $T_{\text{kin}}$ ) of  $\Lambda$  is systematically higher than that of  $K_S^0$  at 3 GeV
- $T_{\text{kin}}$  of  $\Lambda$  and  $K_S^0$  at 3 GeV is lower than  $\pi, K, p$  at higher energy collisions
- ➔ Similar observations for protons and deuterons, implying different EOS at freeze-out



# Summary

- Presented measurements on strangeness production in 3 GeV Au+Au collisions
  - Precise centrality & rapidity dependence of yields
  - $\phi/K^-$  and  $\phi/\Xi^-$  show a strong effect of canonical suppression [Phys. Lett. B 831 \(2022\) 137152](#)
  - $\Lambda$  and  $K_S^0$  spectra indicate lower kinetic freeze-out temperature than  $\pi, K, p$  at higher energy collisions
- At 3 GeV, the measured  $v_2$  for all particles are negative and the NCQ scaling breaks, especially for positive charged particles [Phys. Lett. B 827 \(2022\) 137003](#)
- The suppression of  $C_4/C_2$  is consistent with fluctuations driven by baryon number conservation which indicates a hadronic interaction dominated region in the top 5% central Au+Au collisions at 3 GeV [Phys. Rev. Lett. 128 \(2022\) 202303](#)
- The freeze-out parameter ( $T_{\text{kin}}$ ) of deuteron is systematically higher than that of proton at 3 GeV, which is different from higher energies [Hui Liu, QM 2022](#)

➡ All results from 3 GeV Au+Au collisions: particle production mechanism dominated by hadronic interactions

# Outlook

- Strange hadron yields together with  $\pi, K, p$  will be used for chemical equilibrium models to determine  $T_{chem}$  and  $\mu_B$
- High statistics data in STAR BES II  $\sqrt{s_{NN}} = 3 - 27$  GeV, iTPC+eTOF
  - Extract freeze-out parameters
  - Analyze baryon correlation functions
  - Analyze hyper-nuclei production and collectivity
  - And more ...



See poster from Sameer Aslam, Jun 14, 2022, 5:10 PM