

Displaced or Invisible? ALPs from B decays at Belle II

Ruth Schäfer

Heidelberg University

November 10th 2021


LLPX:

Searching for long-lived particles at the LHC and beyond

Based on an ongoing project with T. Ferber, A. Filimonova, RS, and S. Westhoff;
RS supported by GRK 1940

Long-lived Particles

- ▶ Non-prompt decays
- ▶ Weakly coupled new physics
 - ▶ $c\tau \propto \frac{1}{\text{coupling}^2}$
- ▶ Complementary detector signatures:
 - ▶ Displaced decays
 - ▶ Missing energy
 - ▶ Other signatures



Axion-like particles

- ▶ Use as benchmark model
- ▶ Effective Model of Standard Model + ALP
- ▶ Couples in the UV to either fermions or W -bosons


$$\mathcal{L}_{\text{eff,UV}} = \sum_f \frac{c_{ff}(\mu)}{2} \frac{\partial^\mu a}{f_a} (\bar{f} \gamma_\mu \gamma_5 f) + c_{WW} \frac{a}{f_a} \frac{\alpha_2}{4\pi} W_{\mu\nu}^A \widetilde{W}^{\mu\nu,A}$$

- ▶ This talk: just c_{ff} -term


using Bauer, et al. [2012.12272]

Belle II

- ▶ Belle II is an e^+e^- -collider:
 - ▶ Small boost
 - ▶ Clean background
 - ▶ Optimised for rare B -decays
 - ▶ Full solid angle
- ▶ We produce ALP in $B^+ \rightarrow K^+ a$



Displaced decays




- ▶ The ALP decays into pair of visible particles
- ▶ Within the tracking system (CDC)

$$N_{disp} = N \times Br(B \rightarrow Ka) \times \left(e^{-\frac{d_{res}}{\gamma\beta c\tau}} - e^{-\frac{R}{\gamma\beta c\tau}} \right) \times Br(a \rightarrow f\bar{f}) \times \varepsilon$$

Displaced ALP decays

- ▶ $B^+ \rightarrow K^+ a$, $a \rightarrow \mu^+ \mu^-, e^+ e^-$ displaced
- ▶ Simulated with EvtGen data
- ▶ Assume zero background




Preliminary

Invisible decays

- ▶ ALP decays outside of detector
- ▶ The other B decays generically
- ▶ Inclusive search
- ▶ Analysis of B - and continuum backgrounds




$$N_{inv} = N \times Br(B \rightarrow Ka) \times e^{-\frac{R}{\gamma\beta c\tau}} \times \epsilon$$



Kinematic variables of the system


- ▶ The decay $B \rightarrow Ka$ is given by its final state momenta
 - ▶ $p(K), p(a)$
 - ▶ $p(a)$ is not measurable
 - ▶ p_{miss} is measurable
- ▶ We choose the following variables to describe our system:
 - ▶ Kaon transverse momentum $p_T(K)$
 - ▶ Opening angle between kaon and missing momentum $\phi_{Kp_{miss}}$
 - ▶ Reconstructed B -meson mass \hat{M}_B^2 from $p(K), p_{miss}$

Kinetic variables in our signal samples




Preliminary

Kinetic variables in our backgrounds




Preliminary

Kinetic variables in our backgrounds




Preliminary

Placing cuts on the distributions




Bounds from invisible decays



Displaced or Invisible?

Projected bounds on $BR(B^+ \rightarrow K^+ + X)$




Preliminary


Conclusion

- ▶ Belle II is a good detector for long-lived particles
- ▶ Invisible searches are stronger than displaced ones
- ▶ Displaced searches are useful for characterising LLPs
- ▶ This varies from model to model and detector to detector


Backup plots




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary




Preliminary



Preliminary



Preliminary



Preliminary