



Contribution ID: 7

Type: **not specified**

## **Constraining the properties of superdense matter with observations of low-mass X-ray binaries: the case of HETE J1900.1–2455**

*Monday, September 27, 2021 3:00 PM (35 minutes)*

New observations of a transient neutron star in a low mass X-ray binary HETE J1900.1–2455 were carried out in 2018. Despite detecting only six net photons, it was still possible to infer the effective surface temperature of the star. It turned out to be quite low,  $\approx 30 - 39$  eV (for an observer at infinity) depending on the assumed mass, radius and distance to the star. Taking into account the amount of energy deposited during the  $\approx 10$  years accretion outburst as well as previous temperature measurements in 2016, these values can either indicate that the core has a very high heat capacity or that it undergoes rapid neutrino cooling. Unfortunately, current observational data does not allow us to distinguish between these two possibilities. However, both of them suggest that a significant fraction of the core is not superfluid (superconductor). Also, our modeling shows that the star can cool even further, up to the temperature of  $\approx 15$  eV. In this case future observations (that are already scheduled) might allow us to obtain constraints on the fraction of unpaired baryons in the core.

**Primary authors:** PAGE, Dany (Universidad Nacional Autonoma de Mexico); DEGENAAR, nathalie (university of amsterdam); VAN DEN EIJDEN, J. (Anton Pannekoek Institute for Astronomy, University of Amsterdam); REYNOLDS, M. (Department of Astronomy, University of Michigan); Dr BEZNOGOV, Mikhail (National Institute for R&D in Physics and Nuclear Engineering (IFIN)); WIJNANDS, Rudy (University of Amsterdam)

**Presenter:** Dr BEZNOGOV, Mikhail (National Institute for R&D in Physics and Nuclear Engineering (IFIN))