

Summer Student Lecturs 2021

Q&A - Partical Accelerators

Michaela Schaumann

30.06.2021


What do you consider the most viable long-term future option: linear or circular accelerators?

Asked by Elias

LINAC vs. Storage Ring

- All particles pass a **LINAC** only once.
- A **storage ring** brings the particles back turn by turn.
 - They get another chance to collide again and again potentially over hours.

Curved orbit of electrons in magnet field



Charged particles radiate whenever they are accelerated.

→ energy loss strongly depends on energy and particle mass

$$\Delta E \approx \frac{1}{3\epsilon_0} \frac{q^2}{R} \left(\frac{E}{mc^2} \right)^4$$

→ for electrons at high energies, the radiation becomes too strong and we might have to go linear.

What do you consider the most viable long-term future option: linear or circular accelerators?

Asked by Elias

It depends on what you want to do!

Discover new particles vs. make precision measurements

- Hadron accelerators are discovery machines
 - higher collision energies
 - not well defined initial state
- Electron/Positron accelerators are needed for precision measurements of the newly discovered particles because they provide cleaner collision events.

Could you go into some more detail on how the damping coefficient function $K(s)$, when solving Hill's equation, is implemented in the general solution (I presume it is part of the beta function, but how exactly)?

Asked by Julian

$K(s)$ is no damping coefficient but a with position “s” variable focussing force.

- Solving the harmonic oscillator equation for each element type, provides us with **the elements' transfer matrices**.
- Those can be used to **track the particles' trajectory** through the accelerator lattice **element by element**.
- To know the particles coordinates at a position s , one needs to know all elements and their strengths (k_n) between s_0 and s .
- A **circular accelerator has a certain periodicity** (cell or circumference)
 - the **sequence of strengths is a periodic function $K(s)$**
 - from harmonic oscillator to quasi harmonic oscillator
 - described by Hill's equation that transforms $K(s)$ into $\beta(s)$.

Comparing the two solutions provides the relation between $K(s)$ and the lattice functions

Use matrix formalism:
$$\begin{pmatrix} x \\ x' \end{pmatrix} = M_{foc} \cdot \begin{pmatrix} x_0 \\ x'_0 \end{pmatrix}$$

**Focusing
Quadrupole**
$$M_{foc} = \begin{pmatrix} \cos(\sqrt{K}s) & \frac{1}{\sqrt{K}} \sin(\sqrt{K}s) \\ -\sqrt{K} \sin(\sqrt{K}s) & \cos(\sqrt{K}s) \end{pmatrix}$$

From general solution of Hill's equation after defining $\alpha(s), \gamma(s)$

$$M = \begin{pmatrix} \sqrt{\frac{\beta}{\beta_0}} (\cos \psi + \alpha_0 \sin \psi) & \sqrt{\beta \beta_0} \sin \psi \\ \frac{(\alpha_0 - \alpha) \cos \psi - (1 + \alpha \alpha_0) \sin \psi}{\sqrt{\beta \beta_0}} & \sqrt{\frac{\beta_0}{\beta}} (\cos \psi - \alpha \sin \psi) \end{pmatrix}$$

By knowing the lattice functions $\beta(s), \alpha(s), \gamma(s)$, and $\psi(s)$, we can compute the single particle trajectories between two locations without remembering the exact lattice structure and strength of each element!

Stressing the optics analogy a little bit more: Is it possible to build something like achromatic focusing magnets?


Asked by Augustin

With “chromatic” we mean a system of magnets (incl. and especially dipoles) that in the end do not show a spectrometric effect \rightarrow dispersion disappears.

Which is not possible for a single magnet.

It is however possible to find a sequence of magnets (dipoles and quads) that makes the dispersion disappear (in certain locations at least), e.g. double bend achromat, which is a typical lattice for light sources.


Triple Bend Achromat



horizontal beta-function

vertical beta-function

Double Bend Achromat



dispersion

Questions to warm up ...

Why do we always desire higher particle energy?


Resolution to study structure of matter

$\rightarrow 10^{-18}m$ at LHC


The wavelength of the probe radiation needs to be smaller than the object to resolve.

de Broglie wavelength

$$\lambda = \frac{h c}{E}$$



Particle creation



$$E = m c^2$$

Study of particles that do not exist in our natural environment, since they are too heavy or unstable.

Particle Dimensions




Atomic Radius

$$R_a \approx 10^{-10} \text{ m} = 1 \text{ \AA}$$

Proton Radius

$$R_p \approx 10^{-15} \text{ m} = 1 \text{ "Fermi"}$$

→ 10^{-18} m (LHC resolution) allows us to
"see" structures of one permille of
the proton dimension




Raster tunnel microscope

In terms of universal scale...

astronomical unit:

average distance earth-sun: 1 AE $\approx 150 \cdot 10^6$ km

distance Pluto-Sun: ≈ 40 AE




10^{-18} m on the scale of a meter

means $6 \mu\text{m}$ on the scale of our planetary system

Where is the energy limit? Why are these rings so large?

The ideal circular orbit



Lorentz Force $F_L = q v B$

Centrifugal Force $F_{centr} = \frac{\gamma m_0 v^2}{\rho}$

$$F_L = F_{centr}$$

$$\frac{p}{q} = B \rho$$

$B \rho = \text{Beam rigidity}$

$B < 16 \text{ T}$ \rightarrow build LARGE rings due to the centrifugal force.

q charge,

$p = \gamma m_0 v$ momentum,

B magn. field strength,

ρ bending radius

What limits Luminosity?

Limited by machine size and filling scheme

$$L = \frac{k N^2 f \gamma}{4 \pi \beta^* \epsilon} \cdot F$$


Defined by the injectors

Limited by aperture in the quadrupoles and Q' correction scheme

- N..... No. particles per bunch
- k..... No. bunches
- f..... revolution freq.
- g..... rel. gamma
- β^* beta-function at IPs
- ϵ norm. trans. emit


Limitation:

“Collective effects” cause beam instabilities for too high bunch intensities, too small bunch spacing, too “bright” beams.




What limits Luminosity?

$$L = \frac{kN^2 f \gamma}{4\pi \beta^* \varepsilon} \cdot F$$




When colliding with many bunches, a **crossing angle is needed** to avoid unwanted collisions. However, this **reduces the beam overlap** and therefore the luminosity. Keep as *small as possible*!

→ Limited by beam-beam effects.



Beam-Beam Effects



Beam-Beam force acts (de)focusing
 → Like a quadrupole
 → Changes the tune of each single particles







Fig. 33: Calculated tune shift due to the beam-beam interaction in LHC.



Blue: bam size around IP including beam-beam effect (focusing in this case)
Black: ideal optics without beam-beam



beam-beam destroys the elliptic phase space distribution.
 → creation of tails in the transverse particle distribution (LHeC)



observed particle losses when beams are brought into collision (LHC)