

Explaining B anomalies with Planck-safe Z'

Tom Steudtner
Technische Universität Dortmund
University of Sussex

based on 2109.06201
in collaboration with

Gudrun Hiller, Rigo Bause, Tim Höhne, Daniel Litim

Jahrestreffen der deutschen LHCb-Gruppen
TU Dortmund 08.10.2021

Various hints of new physics in $b \rightarrow s\mu\mu$ decays

$$R_{K^{(*)}} = \frac{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} \mu^+ \mu^-)}{dq^2} dq^2}{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} e^+ e^-)}{dq^2} dq^2}$$

Obs.	Region of q^2 /	Pull _{SM}
R_{K^*}	[0.045, 1.1]	2.5σ
	[1.1, 6.0]	2.5σ
R_K	[1.1, 6]	3.1σ

[LHCb collaboration: 1705.05802, 2103.11769]

Various hints of new physics in $b \rightarrow s\mu\mu$ decays

$$R_{K^{(*)}} = \frac{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} \mu^+ \mu^-)}{dq^2} dq^2}{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} e^+ e^-)}{dq^2} dq^2}$$

Obs.	Region of q^2 /	Pull _{SM}
R_{K^*}	[0.045, 1.1]	2.5σ
	[1.1, 6.0]	2.5σ
R_K	[1.1, 6]	3.1σ

[LHCb collaboration: 1705.05802, 2103.11769]

» described model-independent NP Wilson coefficients

[Bause, Gisbert, Golz, Hiller: 2109.01675]

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu)$$

$$C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

Dim.	Fit	C_9^μ	C_{10}^μ	Pull _{SM}
1d	C_9^μ	-0.83 ± 0.14	0	6.0σ
1d	$C_{10}^\mu = -C_9^\mu$	-0.41 ± 0.07	$-C_9^\mu$	6.0σ
2d	$C_{9,10}^\mu$	-0.71 ± 0.17	0.20 ± 0.13	5.9σ

Various hints of new physics in $b \rightarrow s\mu\mu$ decays

$$R_{K^{(*)}} = \frac{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} \mu^+ \mu^-)}{dq^2} dq^2}{\int_{q_{\min}^2}^{q_{\max}^2} \frac{d\mathcal{B}(B \rightarrow K^{(*)} e^+ e^-)}{dq^2} dq^2}$$

Obs.	Region of q^2 /	Pull _{SM}
R_{K^*}	[0.045, 1.1]	2.5σ
	[1.1, 6.0]	2.5σ
R_K	[1.1, 6]	3.1σ

[LHCb collaboration: 1705.05802, 2103.11769]

» described model-independent NP Wilson coefficients

[Bause, Gisbert, Golz, Hiller: 2109.01675]

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu)$$

$$C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

Dim.	Fit	C_9^μ	C_{10}^μ	Pull _{SM}
1d	C_9^μ	-0.83 ± 0.14	0	6.0σ
1d	$C_{10}^\mu = -C_9^\mu$	-0.41 ± 0.07	$-C_9^\mu$	6.0σ
2d	$C_{9,10}^\mu$	-0.71 ± 0.17	0.20 ± 0.13	5.9σ

$$C_9^{\prime\mu} \sim (\bar{s}_R \gamma_\nu b_R) (\bar{\mu} \gamma^\nu \mu)$$

$$C_{10}^{\prime\mu} \sim (\bar{s}_R \gamma_\nu b_R) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

consistent with zero

[Bause, Gisbert, Golz, Hiller: 2109.01675]

$$C_9^\mu = -1.07 \pm 0.17$$

$$C_9^{\prime\mu} = 0.27 \pm 0.32$$

$$C_{10}^\mu = 0.18 \pm 0.15$$

$$C_{10}^{\prime\mu} = -0.28 \pm 0.19$$

can generate $R_{K^*} \neq R_K$

» How to generate the NP Wilson coefficients?

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu) \quad C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

» How to generate the NP Wilson coefficients?

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu) \quad C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

» Model building: Leptoquarks? Z' models? A mix thereof? ...

» How to generate the NP Wilson coefficients?

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu) \quad C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

» Model building: Leptoquarks? Z' models? A mix thereof? ...

→ ~~light, leptophilic Z'~~


» How to generate the NP Wilson coefficients?

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu) \quad C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

» Model building: Leptoquarks? Z' models? A mix thereof? ...

→ ~~light, leptophilic Z'~~

→ heavy Z' , tree-level couplings to quarks




» How to generate the NP Wilson coefficients?

$$C_9^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \mu) \quad C_{10}^\mu \sim (\bar{s}_L \gamma_\nu b_L) (\bar{\mu} \gamma^\nu \gamma^5 \mu)$$

» Model building: Leptoquarks? Z' models? A mix thereof? ...


→ ~~light, leptophilic Z'~~

→ heavy Z' , tree-level couplings to quarks



» consistent QFT: $U(1)'$ extension of SM gauge group


» fermions have generation-dependent charges



$$C_{9,10}^\mu \sim \frac{g_L^{bs} [g_R^{\mu\mu} \pm g_L^{\mu\mu}]}{M_{Z'}^2}$$

» Direct coupling to quarks: Z' is heavy

$M_{Z'} \gtrsim 5 \text{ TeV}$ first generation quarks [CMS collaboration: 2103.02708]




$$C_{9,10}^{\mu} \sim \frac{g_L^{bs} [g_R^{\mu\mu} \pm g_L^{\mu\mu}]}{M_{Z'}^2}$$

» Direct coupling to quarks: Z' is heavy

$M_{Z'} \gtrsim 5 \text{ TeV}$ first generation quarks [CMS collaboration: 2103.02708]

» Sizable Z' couplings required to account for $C_{9,10}^{\mu}$

$$g_L^{\mu\mu} = g_4 F_{L_2} \quad g_R^{\mu\mu} = g_4 F_{E_2} \quad g_L^{bs} = g_4 V_{tb} V_{ts}^* (F_{Q_3} - F_{Q_2})$$



$$C_{9,10}^{\mu} \sim \frac{g_L^{bs} [g_R^{\mu\mu} \pm g_L^{\mu\mu}]}{M_{Z'}^2}$$

» Direct coupling to quarks: Z' is heavy

$M_{Z'} \gtrsim 5 \text{ TeV}$ first generation quarks [CMS collaboration: 2103.02708]

» Sizable Z' couplings required to account for $C_{9,10}^{\mu}$

$$g_L^{\mu\mu} = g_4 F_{L_2} \quad g_R^{\mu\mu} = g_4 F_{E_2} \quad g_L^{bs} = g_4 V_{tb} V_{ts}^* (F_{Q_3} - F_{Q_2})$$

→ Landau poles of g_4 before the Planck scale

» e.g. *minimal* model

$$M_{Z'} \gtrsim 5 \text{ TeV}$$

left muon and b-quark have $U(1)'$ charge + gauge anomaly cancellation

Landau pole $\mu_L \lesssim 10^{10} \text{ TeV}$


» e.g. *minimal* model

$$M_{Z'} \gtrsim 5 \text{ TeV}$$

left muon and b-quark have $U(1)'$ charge + gauge anomaly cancellation

Landau pole $\mu_L \lesssim 10^{10} \text{ TeV}$

» in practice:



Are all these theories excluded?


» e.g. *minimal* model

$$M_{Z'} \gtrsim 5 \text{ TeV}$$

left muon and b-quark have $U(1)'$ charge + gauge anomaly cancellation

Landau pole $\mu_L \lesssim 10^{10} \text{ TeV}$

» in practice:



Are all these theories excluded?

Landau pole has to be (re)moved!

→ **Planck safety**

What is Planck Safety?

Between EW and Planck scale:

- » no Landau poles, couplings remain finite
- » parameters remain physical
- » scalar potential is stable

What is Planck Safety?

Between EW and Planck scale:

- » no Landau poles, couplings remain finite
- » parameters remain physical
- » scalar potential is stable

→ consistent, predictive until M_{Pl}

“Asymptotic Safety until the Planck scale”

→ provides additional theory constraints

How to achieve Planck Safety (in Practice)

» SM is **not** Planck-safe – Higgs metastability !

→ remove Landau poles and restore Higgs stability

How to achieve Planck Safety (in Practice)

» SM is **not** Planck-safe – Higgs metastability !

→ remove Landau poles and restore Higgs stability

» *template* BSM sector inspired by Asymptotic Safety [Litim, Sannino, JHEP 2014]

ψ_i N_f vector-like fermions (charged) – “quark”

S_{ij} $N_f \times N_f$ matrix-like scalars (uncharged) – “meson”

How to achieve Planck Safety (in Practice)

» SM is **not** Planck-safe – Higgs metastability !

→ remove Landau poles and restore Higgs stability

» *template* BSM sector inspired by Asymptotic Safety [Litim, Sannino, JHEP 2014]

ψ_i N_f vector-like fermions (charged) – “quark”

S_{ij} $N_f \times N_f$ matrix-like scalars (uncharged) – “meson”

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.})$$

unbroken flavor symmetry

How to achieve Planck Safety (in Practice)

» SM is **not** Planck-safe – Higgs metastability !

→ remove Landau poles and restore Higgs stability

» *template* BSM sector inspired by Asymptotic Safety [Litim, Sannino, JHEP 2014]

ψ_i N_f vector-like fermions (charged) – “quark”

S_{ij} $N_f \times N_f$ matrix-like scalars (uncharged) – “meson”

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.})$$

unbroken flavor symmetry

$$\frac{d\alpha_4}{d \ln \mu} = + \# \alpha_4^2 + \# \alpha_4^3 - \# y^2 N_f^2 \alpha_4^2$$

How to achieve Planck Safety (in Practice)

» SM is **not** Planck-safe – Higgs metastability !

→ remove Landau poles and restore Higgs stability

» *template* BSM sector inspired by Asymptotic Safety [Litim, Sannino, JHEP 2014]

ψ_i N_f vector-like fermions (charged) – “quark”

S_{ij} $N_f \times N_f$ matrix-like scalars (uncharged) – “meson”

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.}) \quad \frac{d\alpha_4}{d \ln \mu} = + \# \alpha_4^2 + \# \alpha_4^3 - \# y^2 N_f^2 \alpha_4^2$$

unbroken flavor symmetry

- $N_f, N_c \rightarrow \infty$ exact UV fixed point (asymptotic safety)
- $N_f = 3$, embed in gauge group → potentially enables Planck safety

How to achieve Planck Safety (in Practice) II

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.})$$

$$\frac{d\alpha_4}{d \ln \mu} = + \# \alpha_4^2 + \# \alpha_4^3 - \# y^2 N_f^2 \alpha_4^2$$

How to achieve Planck Safety (in Practice) II

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.}) \quad \frac{d\alpha_4}{d \ln \mu} = + \# \alpha_4^2 + \# \alpha_4^3 - \# y^2 N_f^2 \alpha_4^2$$

» stabilize Higgs potential by scalar portals, sometimes additional Yukawas

$$\sim \delta \text{Tr} [S^\dagger S] (H^\dagger H)$$

How to achieve Planck Safety (in Practice) II

» tame Landau poles by Yukawa interaction

$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.}) \quad \frac{d\alpha_4}{d \ln \mu} = + \# \alpha_4^2 + \# \alpha_4^3 - \# y^2 N_f^2 \alpha_4^2$$

» stabilize Higgs potential by scalar portals, sometimes additional Yukawas

$$\sim \delta \text{Tr} [S^\dagger S] (H^\dagger H)$$

» enhances physics predictivity:

→ previous work with BSM vector-like leptons [Hiller, Hormigos-Feliu, Litim, TS: Phys.Rev.D 102 (2020) 9]

→ simultaneous explanation for $(g - 2)_{\mu, e}$ [Hiller, Hormigos-Feliu, Litim, TS: Phys.Rev.D 102 (2020) 7]

Putting it all together

- » extended gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$
- » SM matter fields Q_i, U_i, D_i, L_i, E_i and Higgs H

Putting it all together

- » extended gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$
- » SM matter fields Q_i, U_i, D_i, L_i, E_i and Higgs H
- » for models with $C'_{10}{}^\mu \neq 0$: right-handed neutrinos ν_i

Putting it all together

- » extended gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$
- » SM matter fields Q_i, U_i, D_i, L_i, E_i and Higgs H
- » for models with $C'_{10}{}^\mu \neq 0$: right-handed neutrinos ν_i
- » $U(1)'$ breaking, charged scalar ϕ

Putting it all together

- » extended gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$
- » SM matter fields Q_i, U_i, D_i, L_i, E_i and Higgs H
- » for models with $C_{10}^\mu \neq 0$: right-handed neutrinos ν_i
- » $U(1)'$ breaking, charged scalar ϕ
- » Planck-Safety sector:
 - vector-like BSM fermion ψ_i
 - uncharged 3 x 3 BSM scalar S_{ij}
- » scalar portals between H, ϕ, S_{ij}

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)
- Higgs is uncharged under $U(1)'$ \rightarrow no interference with EWSB

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)
- Higgs is uncharged under $U(1)'$ \rightarrow no interference with EWSB

» 6 gauge anomaly cancellation conditions for $U(1)'$

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)
- Higgs is uncharged under $U(1)'$ \rightarrow no interference with EWSB

» 6 gauge anomaly cancellation conditions for $U(1)'$

» condition for each component of SM Yukawa matrices

- diagonal quark Yukawas Y_{ii}^u, Y_{ii}^d compatible with $U(1)'$
- lepton and off-diagonal CKM elements are small breaking
 \rightarrow some models allow diagonal lepton and RHN Yukawas Y_{ii}^e, Y_{ii}^ν

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)
- Higgs is uncharged under $U(1)'$ \rightarrow no interference with EWSB

» 6 gauge anomaly cancellation conditions for $U(1)'$

» condition for each component of SM Yukawa matrices

- diagonal quark Yukawas Y_{ii}^u, Y_{ii}^d compatible with $U(1)'$
- lepton and off-diagonal CKM elements are small breaking
 \rightarrow some models allow diagonal lepton and RHN Yukawas Y_{ii}^e, Y_{ii}^ν

» choose CKM rotation in down-sector only

\rightarrow no FCNCs for up-type quarks

Assigning $U(1)'$ charges

» choice:

- BSM fermion ψ_i is SM singlet, but $U(1)'$ charged (invisible!)
- Higgs is uncharged under $U(1)'$ \rightarrow no interference with EWSB

» 6 gauge anomaly cancellation conditions for $U(1)'$

» condition for each component of SM Yukawa matrices

- diagonal quark Yukawas Y_{ii}^u, Y_{ii}^d compatible with $U(1)'$
- lepton and off-diagonal CKM elements are small breaking
 \rightarrow some models allow diagonal lepton and RHN Yukawas Y_{ii}^e, Y_{ii}^ν

» choose CKM rotation in down-sector only

\rightarrow no FCNCs for up-type quarks

» small gauge-kinetic at the electroweak scale mixing between $U(1)_Y \times U(1)'$

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ	
BM₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	0	1	$\frac{1}{5}$
BM₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$	
BM₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	0	1	$\frac{1}{8}$
BM₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM ₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM ₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM ₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM ₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM ₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM ₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM ₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM ₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound
- » tame Landau pole

U(1)' charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM ₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM ₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM ₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM ₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound
- » tame Landau pole
- » $U(1)'$ breaking, no additional Yukawas

$U(1)'$ charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound
- » tame Landau pole
- » $U(1)'$ breaking, no additional Yukawas
- » EWSB without $U(1)'$ sector

$U(1)'$ charges and benchmark models

Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound
- » tame Landau pole
- » $U(1)'$ breaking, no additional Yukawas
- » EWSB without $U(1)'$ sector

$$C_9^\mu \neq 0, C_{10}^\mu = 0 \begin{cases} \rightarrow \text{BM}_1 \\ \rightarrow \text{BM}_4 \end{cases}$$

$$C_9^\mu = -C_{10}^\mu \rightarrow \text{BM}_2$$

$$C_{9,10}^\mu \neq 0 \rightarrow \text{BM}_3$$

U(1)' charges and benchmark models


Model	F_{Q_i}			F_{U_i}			F_{D_i}			F_{L_i}			F_{E_i}			F_{ν_i}			F_H	F_ψ	F_ϕ
BM ₁	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	$\frac{1}{20}$	$\frac{1}{20}$	$-\frac{1}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	$-\frac{9}{10}$	$\frac{9}{10}$	0	0	0	0	1	$\frac{1}{5}$
BM ₂	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{6}$	0	1	0	0	0	1	$\frac{1}{12}$	$-\frac{1}{12}$	1	0	$\frac{11}{12}$	$\frac{1}{9}$
BM ₃	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	$-\frac{1}{8}$	$-\frac{1}{8}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{2}$	0	1	$\frac{1}{8}$
BM ₄	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	0	$\frac{1}{9}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{3}$	0	1	$\frac{1}{6}$

- » pass 6 gauge anomaly cancellation conditions
- » allow at least diagonal quark Yukawas
- » no Z' – electron couplings
- » no Kaon mixing
- » B_s mixing bound
- » tame Landau pole
- » $U(1)'$ breaking, no additional Yukawas
- » EWSB without $U(1)'$ sector


$C_9^\mu \neq 0, C_{10}^\mu = 0$ \rightarrow BM₁ no right-handed neutrinos
 \rightarrow BM₄ lighter Z'

$C_9^\mu = -C_{10}^\mu \rightarrow$ BM₂



$C_{9,10}^\mu \neq 0 \rightarrow$ BM₃





How to check for Planck-Safety





How to check for Planck-Safety




How to check for Planck-Safety




How to check for Planck-Safety




RG running




RG running



RG running




RG running




BSM Yukawa brakes Landau pole



RG running



RG running





BSM critical surface



$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.})$$

$$-\mathcal{L}_{\text{portal}} = \delta \text{Tr} [S^\dagger S] (H^\dagger H) + \tilde{\delta} (\phi^\dagger \phi)(H^\dagger H)$$

BSM critical surface



$$-\mathcal{L}_y = y (\bar{\psi}_{Li} S_{ij} \psi_{Rj} + \text{h.c.})$$

$$-\mathcal{L}_{\text{portal}} = \delta \text{Tr} [S^\dagger S] (H^\dagger H) + \tilde{\delta} (\phi^\dagger \phi)(H^\dagger H)$$

$$\alpha_y(\mu_0) \gtrsim 10^{-1.25} \dots 10^{-1}$$

bounds on $\alpha_\delta, \alpha_{\tilde{\delta}}$ vary among BMs

Phenomenology

» broad decay of Z' to invisibles $Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu$ with 65 .. 85% BR

Model	jets	b	t	e	μ	τ	$\nu_{e,\mu,\tau}$	h	$\psi_{1,2,3}$	ϕ
BM₁	0.5	0.5	0.5	0	15	15	15	0	54	0.2
BM₂	14	1.5	1.5	0	9	9	18	0	46	0.1
BM₃	5	0	0	0	4	4	8	0	79	0.1
BM₄	0	0.9	0.9	0	3	11	14	0	72	0.2

Phenomenology

» broad decay of Z' to invisibles $Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu$ with 65 .. 85% BR

Model	jets	b	t	e	μ	τ	$\nu_{e,\mu,\tau}$	h	$\psi_{1,2,3}$	ϕ
BM₁	0.5	0.5	0.5	0	15	15	15	0	54	0.2
BM₂	14	1.5	1.5	0	9	9	18	0	46	0.1
BM₃	5	0	0	0	4	4	8	0	79	0.1
BM₄	0	0.9	0.9	0	3	11	14	0	72	0.2

» can be probed & models distinguished at $\mu\mu$ collider:

$$\sigma(\mu^+\mu^- \rightarrow Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu) \approx (10^2..10^3) \sigma(\mu^+\mu^- \rightarrow Z \rightarrow \bar{\nu}\nu)^{\text{SM}}$$

Phenomenology

» broad decay of Z' to invisibles $Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu$ with 65 .. 85% BR

Model	jets	b	t	e	μ	τ	$\nu_{e,\mu,\tau}$	h	$\psi_{1,2,3}$	ϕ
BM₁	0.5	0.5	0.5	0	15	15	15	0	54	0.2
BM₂	14	1.5	1.5	0	9	9	18	0	46	0.1
BM₃	5	0	0	0	4	4	8	0	79	0.1
BM₄	0	0.9	0.9	0	3	11	14	0	72	0.2

» can be probed & models distinguished at $\mu\mu$ collider:

$$\sigma(\mu^+\mu^- \rightarrow Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu) \approx (10^2..10^3) \sigma(\mu^+\mu^- \rightarrow Z \rightarrow \bar{\nu}\nu)^{\text{SM}}$$

» only mildly enhanced $B \rightarrow K^{(*)}\bar{\nu}\nu$, consistent with SM expectation

Phenomenology

» broad decay of Z' to invisibles $Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu$ with 65 .. 85% BR

Model	jets	b	t	e	μ	τ	$\nu_{e,\mu,\tau}$	h	$\psi_{1,2,3}$	ϕ
BM₁	0.5	0.5	0.5	0	15	15	15	0	54	0.2
BM₂	14	1.5	1.5	0	9	9	18	0	46	0.1
BM₃	5	0	0	0	4	4	8	0	79	0.1
BM₄	0	0.9	0.9	0	3	11	14	0	72	0.2

» can be probed & models distinguished at $\mu\mu$ collider:

$$\sigma(\mu^+\mu^- \rightarrow Z' \rightarrow \bar{\psi}\psi, \bar{\nu}\nu) \approx (10^2..10^3) \sigma(\mu^+\mu^- \rightarrow Z \rightarrow \bar{\nu}\nu)^{\text{SM}}$$

» only mildly enhanced $B \rightarrow K^{(*)}\bar{\nu}\nu$, consistent with SM expectation

» benchmarks are consistent with LHC search [CMS collaboration: 2103.02708]

Summary

Model	μ_0	$\alpha_4(\mu_0)$	C_9^μ	C_{10}^μ	$Y_{ii}^{u,d}$	Y_{ii}^e	Y_{ii}^ν	r_{B_s}	$\mathcal{B}(Z' \rightarrow \text{inv.})$	ν_R
BM₁	5 TeV	$1.87 \cdot 10^{-2}$	-0.83	0	✓	✓	X	0.35	73%	X
BM₂	5 TeV	$5.97 \cdot 10^{-3}$	-0.41	$-C_9^\mu$	✓	X	X	0.86	64%	✓
BM₃	5 TeV	$4.60 \cdot 10^{-2}$	-0.71	+0.24	✓	X	X	0.60	87%	✓
BM₄	3 TeV	$2.46 \cdot 10^{-2}$	-0.83	0	✓	✓	✓	0.70	86%	✓

Summary

Model	μ_0	$\alpha_4(\mu_0)$	C_9^μ	C_{10}^μ	$Y_{ii}^{u,d}$	Y_{ii}^e	Y_{ii}^ν	r_{B_s}	$\mathcal{B}(Z' \rightarrow \text{inv.})$	ν_R
BM₁	5 TeV	$1.87 \cdot 10^{-2}$	-0.83	0	✓	✓	X	0.35	73%	X
BM₂	5 TeV	$5.97 \cdot 10^{-3}$	-0.41	$-C_9^\mu$	✓	X	X	0.86	64%	✓
BM₃	5 TeV	$4.60 \cdot 10^{-2}$	-0.71	+0.24	✓	X	X	0.60	87%	✓
BM₄	3 TeV	$2.46 \cdot 10^{-2}$	-0.83	0	✓	✓	✓	0.70	86%	✓

heavy Z' models that

- » explain B-anomalies in several interesting NP scenarios
- » compliant with anomaly cancellation, quark Yukawas, precision measurements
- » are predictive until M_{Pl} \rightarrow no Landau poles
- » stabilize the Higgs potential
- » can be probed at colliders
- » decay mostly to invisibles