

Give all results with three significant digit.

1 - Exercise 1

1. Give the formula of the maximal kinetic energy that can be transferred to an atomic electron by an incoming particle in the regime where $\beta\gamma$ reduced to β , expressing explicitly, in this approximation, the energy in the center of mass of the incoming particle and the atomic electron.
2. Write the Bethe and Bloch formula in this regime neglecting the atomic shell and density corrections.
3. Explain how charged low-energy heavy particles of same velocity and different mass could be separated by measuring the energy loss in a thin detector.
4. For particles (Pa) of electric charge z (in unit of e) show that if the detector is thin enough the average energy loss is given by $\Delta E_{Pa} = z^2 \Delta E_{proton}$ for protons with the same velocity of Pa .
5. Express $\beta\gamma$ as function of the kinetic energy, T , and the mass m .
6. Compute $\beta\gamma$ of α particles with $T=0.4$ GeV (take $m_\alpha = 4$ GeV).
7. Compute the energy loss in hydrogen for protons of $T= 0.1$ GeV using a valid approximation for the maximal kinetic energy that can be transferred to an atomic electron by an incoming particle and neglecting the atomic shell and density corrections (take $m_p = 1$ GeV).
8. Compute the energy loss in hydrogen for α particles with $T=0.4$ GeV (take $m_\alpha = 4$ GeV).
9. How could these results be used to identify heavy-charge particles?

2 - Exercise 2

An accelerator produces 100 MeV protons and deuterons ($m_p = 938$ MeV/ c^2).

1. Show that both particles satisfy the low-energy condition $2\gamma \frac{m_e}{m_{Pa}} \ll 1$.
2. Compute the time of flight of both particles over a 10 m distance.
4. Compute the average specific energy loss in a plastic scintillator made of CH_x (specific mass = 1.03 g cm^{-3}) for both particles, where $x = 1.1$ is the equivalent molecular H/C ratio. We will neglect the density and atomic shell corrections.
5. Compute for both particles the average energy loss ΔE in a 1-cm thick detector D1 made of CH_x .
6. What should be the approximate thickness of a second detector D2 located downstream of D1, if one wants to stop both protons and deuterons inside and measure their energies?
7. If D1 is separate from D2 by 10 m what should be the signal time difference between D1 and D2 for protons and deuterons?
8. Could you use that to separate protons from deuterons?

3 - Exercise 3

1. What is the maximum kinetic energy of delta rays produced by 10 GeV/ c K^- mesons in liquid nitrogen (at boiling point)?
2. How many delta rays with kinetic energy greater between 5 and 20 MeV are produced in 2 m? The specific mass of liquid nitrogen at boiling point is 1.3954 g/ cm^3 .
3. Can the K^- mesons of point 1. produce Cherenkov radiation in liquid nitrogen (the refractive index of liquid nitrogen is 1.205)?